

THE
MEDICAL FORMULARY:

BEING
A COLLECTION OF PRESCRIPTIONS,
DERIVED FROM
THE WRITINGS AND PRACTICE
OF
MANY OF THE MOST EMINENT PHYSICIANS
IN AMERICA AND EUROPE.

TO WHICH IS ADDED

AN APPENDIX,

CONTAINING

THE USUAL DIETETIC PREPARATIONS AND ANTIDOTES FOR POISONS.

THE WHOLE ACCOMPANIED WITH A FEW BRIEF

PHARMACEUTIC AND MEDICAL OBSERVATIONS.

BY BENJAMIN ELLIS, M. D.

LATE PROFESSOR OF MATERIA MEDICA AND PHARMACY IN THE PHILADELPHIA COLLEGE OF
PHARMACY.


"Morbos autem, non eloquentia sed remediis curari."—*Cels. De Med. Lib. I.*

SIXTH EDITION, REVISED AND EXTENDED.

BY SAMUEL GEORGE MORTON, M. D.

PHILADELPHIA:
LEA & BLANCHARD.

1842.



QV

E47m


1842



ENTERED according to the Act of Congress, in the year one thousand eight hundred and forty-one, by

LEA & BLANCHARD,

in the Clerk's Office of the District Court of the Eastern District of Pennsylvania.



GRIGGS & CO., PRINTERS.

TO

N. CHAPMAN, M. D.

PROFESSOR OF THE INSTITUTES AND PRACTICE OF PHYSIC AND
CLINICAL PRACTICE,

IN THE

UNIVERSITY OF PENNSYLVANIA,

ETC. ETC.

WHOSE TALENTS AND URBANITY

HAVE RAISED HIM TO THE FIRST RANK IN HIS PROFESSION, AND HAVE ACQUIRED FOR HIM
THE CONFIDENCE AND ESTEEM

OF A LARGE AND ENLIGHTENED COMMUNITY,

THIS WORK

IS VERY RESPECTFULLY DEDICATED

BY HIS OBLIGED FRIEND,

THE AUTHOR.

C O N T E N T S.

CLASS I. Emetics -	17
II. Cathartics	24
<i>Enemata</i>	42
III. Diuretics	45
IV. Antilithics	56
V. Diaphoretics	59
VI. Expectorants	70
<i>Inhalation</i>	84
VII. Emmenagogues	88
VIII. Anthelmintics,	95
IX. Epispastics	102
<i>Rubefacients</i>	104
<i>Baths, Fomentations, &c.</i>	119
<i>Caustics, or Escharotics</i>	123
<i>Mineral Acids</i>	124
<i>Moxa</i>	125
<i>Stimulating Ointments, &c.</i>	126
<i>Issues and Setons</i>	128
<i>Bougies</i>	130
X. Stimulants	132
XI. Narcotics	141
XII. Antispasmodics	156
XIII. Sialagogues	161
XIV. Tonics	170
<i>Alteratives</i>	188
XV. Astringents	200
<i>Injections</i>	209
XVI. Absorbents	222
APPENDIX, No. 1. Dietetics	228
II. Poisons	238
III. Endermic Medicine, &c.	248
INDEX	251

THE AUTHOR'S ADVERTISEMENT

TO THE THIRD EDITION.

IN preparing a new edition of the Medical Formulary for publication, we have endeavoured to render the work as complete as the nature of the subject will admit. The materials have been carefully revised, and almost every class of medicines augmented, by the addition of many prescriptions derived from the periodical journals, standard medical works, and our professional friends.

We are chiefly indebted to the writings of M. Majendie, for the formulæ relating to the newly discovered alkaline remedies. If we have occasionally inserted compounds which apparently belong to the pharmacopœia, it was from a conviction of their utility to the country practitioner, and also from the fact that they are not contained in the ordinary pharmacopœias.

The list of *Poisons* with their *Antidotes*, has been revised;—the Index also has been formed on a different plan from that of the first edition, and in such a way that simple medicines, and their numerous compounds, may be referred to at a glance.

We cannot close this article without expressing our sincere acknowledgments to our friend Dr. Chapman, for the very flattering manner in which he has recommended the Medical Formulary to his class in the University of Pennsylvania.

THE EDITOR'S ADVERTISEMENT

TO THE SIXTH EDITION.

HAVING been originally associated with my lamented friend, the late Dr. Ellis, in preparing this work for the press, and having superintended two editions since his death, I now submit it to the medical public in a revised and extended form. Especial care has been taken to employ the chemical nomenclature at present in use, and to add to the former series of remedial combinations, all those of recent discovery or modern application. This, however, I have endeavoured to accomplish without altering the simple arrangement or unpretending character of the original work.

S. G. MORTON.

November 1, 1841.

P R E F A C E.

THE elegant and judicious formation of prescriptions is one of the difficulties which the young practitioner in medicine is obliged to encounter. While a student, he is compelled from the circumstances under which he is placed, to confine his attention to the leading principles of the science. Consequently the minor points, (and the art of directing medicines is considered one of them,) are postponed to that period when he shall have undertaken the practical duties of his profession.

To obviate, in some measure, the inconvenience which the graduate at first experiences, the volume now offered to the public was undertaken and executed. It contains, in a condensed form, and we think advantageously arranged, many of the most important prescriptions employed in modern practice, and which, though most of them may be already recorded, are scattered through various medical works.

The arrangement framed by Professor Chapman for his Therapeutics, appearing to combine greater advantages than any other, we have taken the liberty of adopting as the basis of this Formulary.

Each class has been subdivided into Powders, Pills, and Liquids; without any attempt to preserve the different preparations of the same article together, but only with a view to convenience. A few concise rules will be found prefixed to each class, and observations have been attached to those prescriptions, which possess peculiarities either in their composition or effects.

The application of the remedies to diseases has been generally left to the judgment of the practitioner, and Therapeutical detail as much as possible avoided, as it would have been inconsistent with the nature and design of the work.

The object is to furnish models for extemporaneous formulæ, the proportions of which may be either increased or diminished, according to the age, sex, constitution or idiosyncrasy of the patient, all of which are to be taken into consideration when a proscription is directed, as well as the climate and peculiar epidemic which may be prevailing.

The prescriptions have been couched in the Latin language, and the names of the several articles will be found regularly terminated. The design was to furnish something like a Medical Grammar, to those unacquainted with the dead language; and even those versed in the classics, may occasionally find words used in prescriptions which are new to them.

A class of Dietetic preparations has been placed in the Appendix, No. I. They are written in the English language, and it is hoped that they may be found

useful in furnishing directions necessary to be given to the nurse.

In the Appendix, No. II. we have arranged those poisons which are usually taken by accident or design, with their appropriate antidotes and some practical observations.

Appendix No. III. contains some concise remarks on the mode of using remedies on the "*endermic plan*," as well as the quantities employed, and the effects produced by several of the most potent remedial agents.

The materials for the work have been collected from various sources, the writings of different authors, periodical journals, manuscripts and the contributions of our friends. It is hoped therefore, that the difficulties which have attended its progress will form some apology for the errors which it may possibly be found to contain. As a reason for not more frequently attaching the names of authors to the prescriptions, it may be observed, that they are a kind of common medical property which every physician feels himself privileged not only to employ, but modify as may best correspond with his own views. It is, therefore, difficult to say to whom many of them belong; they have been touched, and retouched, till they have lost their identity.

In conclusion, we beg leave to present our acknowledgments to some of our friends for the encouragement they have given to the undertaking. To Dr. Chapman we are particularly indebted for the kindness and

interest he has manifested in its publication, and offer to him, (the only return we can make,) our warmest thanks. It is with particular pleasure we embrace this opportunity of expressing our obligations to our friend Dr. S. G. Morton, whose intimate acquaintance with the subject, and classical knowledge, are only equalled by the kindness and industry he has displayed in facilitating the work.

There are others whose names it would gratify us to mention, but who, we know, will be more pleased by our silence. The countenance they gave us on the present occasion has only increased the debt previously incurred.

B. E.

INTRODUCTION.

BEFORE entering upon the main object of this work, it is necessary to notice a few circumstances which pertain more especially to Pharmaceutic Chemistry, yet appear to be called for in this place; and though they are familiar to every educated physician, may be useful to the student who has not yet turned his attention to prescriptions.

In the first place it is requisite to be well acquainted with the *signs of quantity*, or those symbolic characters which are used in writing prescriptions. In the ancient practice of medicine there were very many of these symbols, not only for the purpose of denoting the quantity of the ingredient, but the ingredient itself. Most of these, however, have become obsolete; and a few only are retained, as indispensable to the convenience of the practitioner. These will be found in the following *Table*, which is adopted in the Pharmacopœias for the regulation of *weights*.

<i>English Name.</i>	<i>Latin Name.</i>	<i>Symbol.</i>	<i>Contains</i>
A pound,	Libra,	℔,	12 ounces.
An ounce,	Uncia,	ʒ,	8 drachms.
A drachm,	Drachma,	ʒ,	3 scruples.
A scruple,	Scrupulus,	ʒ,	20 grains.

In the manipulation of *liquid substances*, *measures* have in most countries superseded the use of *weights*:

the *Table of measures* recommended by the London College, gives particular names and symbols to the parts forming a gallon, in order to prevent their being confounded with those denoting weights; as in the following *Table* :—

<i>English Name.</i>	<i>Latin Name.</i>	<i>Symbol.</i>	<i>Contains</i>
A gallon,	Congius,	none,	8 pints.
A pint,	Octarius,	O,	16 fluidounces.
A fluidounce,	Fluiduncia,	f 3,	8 fluidrachms.
A fluidrachm,	Fluidrachma,	f 3,	60 minims.

It has been thought unnecessary in the following pages to prefix the f to the signs of quantity used for liquids, inasmuch as it is not a custom in this country to do so, nor does the omission lead to any ambiguity in compounding medicines.

The *minims* used in the table, have been adopted by the London College in place of *drops*; and although we have not employed this measure in our prescriptions, yet we can readily perceive it possesses some important advantages over the ancient practice. The symbol (*m*.) is sometimes used by medical writers, though very improperly, to denote both minims and drops.* The size of a drop varies according to the greater or less fluidity and gravity of the liquid, and the shape of the vessel from which it is dropped; but as a general rule, we cannot do better on these occasions than use a small phial with a thin edge. Dr. Dorsey observes, that, “whenever great precision is necessary, it is easy to dilute the active medicine, and give it in form of a mixture. If sixty grains be diffused

* It will be well to recollect that ten minims of any fluid, if measured by the graduated minim measure, will amount to about fifteen drops.

in six ounces of water, for instance, each half ounce will contain five."

Whenever a *table-spoonful* of any liquid is ordered, it is considered to be equal to *half an ounce* by measure; and in the same way a *tea-spoonful* is used for a *drachm*: these measures are sufficiently accurate where no great precision is requisite.

All the *prescriptions* contained in this work, are given in the *Latin language*, without any other abbreviations than those in common use to denote the *quantity* of an ingredient. The *directions* for the administration of medicines are expressed in *English*.

Prescriptions, as usually sent to the Apothecaries, are much abbreviated, and the same course may be pursued with regard to those contained in this collection. For the information of students who have not been in the practice of writing prescriptions, a *Table of Abbreviations* is subjoined.

TABLE OF ABBREVIATIONS.

<i>Abbreviation.</i>	<i>Latin Word.</i>	<i>English Word.</i>
ââ	* Ana,	Of each.
Ad saturand.	Ad saturandum,	Until saturated.
Ad lib.	Ad libitum,	At pleasure.
Aq. tepid.	Aqua tepida,	Warm water.
Aq. ferv.	Aqua fervens,	Hot water.
Chart.	Chartula,	A small paper.
Coch.	Cochlear,	A spoonful.
Coch. mag.	Cochlear magnum,	A table-spoonful.
Coch. parv.	Cochlear parvum,	A tea-spoonful.
Colent.	Colentur,	Let them be strained.
Collyr.	Collyrium,	An eye-water.
Contus.	Contusus,	Bruised or broken.
F. vel Ft.	Fiat, vel Fiant,	Let there be made.
Fol.	Foliuræ, vel folia,	A leaf or leaves.
Garg.	Gargarysma,	A gargle.
Gr.	Granum, vel grana,	A grain or grains.
Gtt.	Gutta, vel guttæ,	A drop or drops.
Haust.	Haustus,	A draught.
Infus.	Infusio,	An infusion.
M.	Misce,	Mix.
Mass.	Massa,	A mass.
Mist.	Mistura,	A mixture.
Pil.	Pilula, vel pilulæ,	A pill or pills.
Pulv.	Pulvis, vel pulveres,	A powder or powders.
Q. S.	Quantum sufficit,	A sufficient quantity.
R.	Recipe,	Take.
Rad.	Radix,	A Root.
S.	Signa,	Write.
Ss.	Semi, vel dimidius,	The half.
Tinct.	Tinctura,	A tincture.

The *doses* throughout the book are applicable to adult age, unless the contrary is specified.

* This is not properly a Latin word, but the Greek proposition *αυτ.*



CLASS I.

E M E T I C S.

IN the administration of Emetics, a few simple rules are worthy of attention. Whatever may be the apparent necessity for emptying the stomach, we must take into consideration all the circumstances of the case, and more particularly if there exist much arterial excitement, with determination to the head; for where these symptoms are present, blood-letting should be sometimes premised.

Where *poisons* have been taken, our object is to evacuate the stomach as speedily as possible, and for this purpose we select an emetic which is prompt in its action. But where the excitability of the stomach is greatly diminished by a *Narcotic*, it is necessary to assist the emetic by the addition of some excitant. Where such an effect is caused by opium, the system may be generally rendered susceptible by the use of vinegar, mustard, or ammonia. In these cases, however, the stomach-tube is our best resource.

Where free vomiting is desirable, it is effectually solicited by plentiful draughts of tepid water or camomile tea.

The evacuation of the stomach may be greatly facilitated by tickling the fauces with a feather, or introducing the point of the finger as far back as possible, and titillating the parts.

There are a few cases in which emetics are obviously inadmissible, as in apoplexy, or where there is great

determination of blood to the head, depending upon constitutional causes; pregnancy; hernia; prolapsus uteri; active hemorrhage from the lungs and uterus, &c. If emetic medicines are ever allowable in these diseases, it is for the most part in nauseating doses only.

By the frequent use of emetics, the stomach acquires an increased susceptibility to their action; hence persons of delicate habits should use them cautiously.—When excessive vomiting occurs, it may be checked by resort to some one or more of the following resources:

Ice broken into small pieces, and allowed to dissolve in the mouth.

Lime water and milk, a table-spoonful of each, mixed cold, and given at intervals of fifteen or twenty minutes.

A drop, or even half a drop of oil of Creosote, mixed in a wine-glass of cold water, is one of the most effectual anti-emetic remedies.

Brandy and water, and clove, and green tea, conduce to the same end; and where other internal means fail in irritable constitutions, an anodyne injection will sometimes accomplish the object.

Among external appliances, mustard to the epigastrium is one of the best: also, spice-plasters, and hot fomentations of brandy and cloves.—

POWDERS.

Powder of Ipecacuanha.

R. Pulveris Ipecacuanhæ, ℥ij.

Divide it into two doses.—Signa.—One to be taken in a little molasses, or sugar and water, and followed by a draught of tepid water. If one powder does not produce the desired effect, the second to be taken in like manner.

Powder of Ipecacuanha with Tartarized Antimony.

R. Pulveris Ipecacuanhæ, ℥j.

Tartratis Antimonii et potassæ, gr. j.

Misce.—To be taken in the same manner as the former.

Ipecacuanha is supposed by some to counteract the effects of opium by a *specific principle*, and hence the above prescription is often used where that drug has been taken: a more prompt emetic, however, will in such cases, be preferable to this.

Powder of Ipecacuanha with Rhubarb.

R. Pulveris Ipecacuanhæ.

—— Rhei, āā. ℥j.

Misce, et signa.—To be taken in sirup.

More or less purging will generally follow the emetic operation of this dose.

The above are the principal combinations of ipecacuanha for the purpose of *vomiting*. It fulfils several other indications, which will be noticed under their appropriate heads.

Powder of Sulphate of Copper.

R. Sulphatis Cupri, gr. ij, vel x.

Signa.—To be given in a little sirup of water.

This medicine is seldom employed as an emetic except in those cases of poisoning where other articles have failed, or are not at hand. It should be followed by a draught of warm water. All vegetable astringent infusions are incompatible substances.

Powder of Sulphate of Zinc.

R. Sulphatis Zinci, gr. x, vel ℥ss.

Signa.—To be taken in molasses, and followed by a draught of warm water.

This medicine is very prompt in its operation; and is most generally employed in cases of poisoning. It is also used as a nauseating remedy in phthisis; and both as an emetic and tonic in dyspepsia. The stomach is said to be invigorated, rather than exhausted by its operation.

WINES, MIXTURES, &c.

Wine of Ipecacuanha with Antimony.

R. Vini Ipecacuanhæ, ℥j.

Tartratis Antimonii et Potassæ, gr. j.

Fiat Solutio.—Signa.—A tea-spoonful to be taken every ten or fifteen minutes until vomiting is induced, which may be assisted with tepid drinks.

Antimonial Wine.

R. Vini Antimonii, ℥j.

Signa.—A tea-spoonful to be taken every ten or fifteen minutes until free emesis is effected.

If antimonial wine is given for the purpose of producing speedy vomiting, we may safely give half an ounce of it at once; and in the croup of infants, twenty drops every fifteen minutes, until the medicine operates, is a safe dose.

Solution of Tartrate of Antimony.

R. Tartratis Antimonii et Potassæ, gr. vj.

Sacchari albi, ℥j.

Aquæ destillatæ, ℥iv.

Misce.—Fiat Solutio.—A table-spoonful to be given every ten or fifteen minutes, until it operates freely.

The *wine of antimony* possesses no advantage over this extemporaneous prescription; and unless we know the proportion of antimony contained in each ounce of the former, and know also that it has been recently made, the latter is decidedly preferable.

Tartar Emetic and Quinine.

R. Tartratis Antimonii et Potassæ, gr. iij.

Sulphatis Quiniæ, gr. x.

Misce et divide in partes vj. equales.—This preparation is commended by Dr. Gola as successful in intermittent fever, when quinine alone

produced no favourable result. He directs a powder to be given every two hours during the apyrexia. It causes both emesis and alvine evacuations, and prevents the recurrence of the paroxysm.

Solution of Sulphate of Zinc, &c.

R. Zinci Sulphatis, ℥iss.

Aluminis, ℥ss.

Aquæ ferventis, Oss.

Misce.—Let the solution be filtered through blotting paper.—Signa.—

A table-spoonful for adults, and a tea-spoonful for a child of six months old, (and so in proportion to the strength and age of the patient,) every morning upon an empty stomach.

This is the celebrated *solutio vitriolica* of Dr. Mosely, who principally employed it in dysentery and phthisis pulmonalis.

Mustard Draught.

R. Pulveris Sinapeos, ℥j.

Aquæ tepidæ, Oss.

To be mixed and swallowed at once.

This is a very prompt emetic, and one that is always at hand.

Emetic Lozenges of Emetina.

R. Sacchari albi, ℥ij.

Emetinæ impuræ, gr. xxxij.

Misce, et fiant trochisci singuli, gr. xvij.

One of these Lozenges taken fasting will generally produce vomiting in children. Three or four will speedily excite it in adults.

Majendie.

Solution of Emetina.

R. Emetinæ fuscæ, gr. iv.

Aquæ destillatæ, ℥ij.

Syrupi simplicis, ℥ss.

Fiat solutio.—Signa.—Give two tea-spoonsful every half hour.

Emetina is destitute of the peculiar odour and nauseous taste of Ipecacuanha, and may therefore be preferable in some cases.

It will be observed that we have introduced the impure or coloured Emetina, which is much less energetic than the perfectly white alkali. The best antidote for an over-dose, is a strong infusion of nut-galls.

One grain of Emetina will generally produce as much effect as sixteen grains of Ipecacuanha. Even this quantity of *pure* Emetina would be an unnecessary large dose; and the emetic property will be readily obtained by dissolving a grain in an ounce of water, and giving it in three or four doses at the usual intervals.

Mixture of Ipecacuanha, &c.

R. Pulveris Ipecacuanhæ, ℥ss.

Antimonii et potassæ tartratis, gr. j.

Tincturæ Scillæ, ℥j.

Aquæ destillatæ, ℥vijss.

Fiat mistura.—Signa.—Four table-spoonsful at first; afterwards two table-spoonsful every ten minutes, until vomiting is induced.

This mixture is recommended in dropsies previously to giving the digitalis.

Infusion of Ipecacuanha.

R. Pulveris ipecacuanhæ, ℥ij.

Aquæ bullientis, ℥vj.

Macerate for one hour and filter.—Of this from one to two table-spoonsful, may be given every half hour, until in excite vomiting.

Infusion of Lobelia Inflata.

R. Herbæ Lobeliæ inflatæ, ℥j.

Aquæ bullientis, Oj.

Fiat infusio.—Signa.—A small wine-glassful may be taken every half hour, until vomiting succeeds.

This native remedy, called also Indian Tobacco, is frequently exhibited in the form of saturated tincture. The dose of the latter is a table-spoonful every half hour until emesis is excited. It has gained considera-

ble celebrity in the treatment of Asthma, and is now employed by the "Steam Doctors" of the western country, as part of their treatment of all diseases. Its activity is so great that it should be administered with caution.

Sirup of Seneca and Squills.

R. Syrupi Scillæ compositi, ℥j.

Signa.—Give from ten drops to a tea-spoonful, every quarter, half, or one hour, according to the age of the patient and violence of the disease.

This is the *Hive Sirup* of Professor Coxe: it has acquired much confidence from its efficacy in croup: and should it not be sufficiently active in this violent disease, a grain of tartrate of antimony may be added to each ounce.—*See Coxe's Dispen.* p. 343.

Tobacco Poultice.

R. Foliorum Nicotianæ, ℥j.

To be beat up with sufficient water to form a cataplasm.—To be applied to the region of the stomach, when emetics cannot be administered by the mouth.

CLASS II.

CATHARTICS.

As a general rule, it may be observed of cathartics, that they are capable of fulfilling three different indications :—1st. Simply to evacuate the bowels ; 2d. To excite an increased discharge from the mucous coat of the intestines ; and 3d. To stimulate the neighbouring viscera, and cause them to pour out a greater quantity of their peculiar fluids. It seldom, or perhaps never, happens, that the same article will produce these three different effects ; in order to do this, we must combine different substances in one formula.

There are some, as magnesia, that produce their effect by absorbing the acid generated in the alimentary canal, and thus become converted into a purgative salt. There are others, as aloes, that expend their principal force upon the rectum and great intestines, and for this reason are very advantageously employed in diseases of the lower bowels, as also in those of the uterus and its appendages.

These remedies should always be given on an empty stomach : if administered immediately after a full meal, they arrest the digestive process, are liable to produce nausea, if not vomiting, and do not open the bowels with the same certainty or effect. When simple costiveness is to be removed, the evening may be selected for the exhibition of a cathartic.

If this should not prove sufficient, we can repeat the dose in the morning and at regular intervals through the day, until the effect is produced. The principle,

says Dr. Paris, which should govern our conduct in the administration of purgatives, may be briefly stated.—*The excitement is to be repeated till the requisite action is induced, yet no single excitement being such as may prove an irritant to the organ.*

But in the treatment of many diseases it is preferable to purge through the day, in order that the sleep of the patient may not be disturbed at night.

The operation of these remedies may be very much accelerated by the free exhibition of diluting drinks, such as oatmeal gruel, barley water, &c. &c.

By combination we are enabled to modify and control the action of many remedies, and some of the most drastic purgatives may be made to operate mildly, and yet certainly, by uniting several of them in the same dose.

Finally, the operation of a cathartic may often be promoted by a judicious venesection; and if spasm of the intestines be present, opium may be advantageously joined with the purgative.

POWDERS.

Powder with Calomel and Jalap.

R. Hydrargyri Chloridi mitis, gr. v. vel x.

Pulveris Jalapæ, gr. x.

Misce, et signa.—To be given in molasses or sirup.

Powder with Calomel, Jalap and Rhubarb.

R. Hydrargyri Chloridi mitis, gr. v.

Pulveris Jalapæ,

—— Rhei, āā, gr. v.

Olei Cinnamomi, gtt. j.

Misce, et signa.—To be given as the former.

Powder with Magnesia and Rhubarb.

R. Pulveris Rhei, ʒj.

Magnesiae ustæ, ʒss.

Olei Cinnamomi, gtt. j.

Misce, et signa.—To be given in sirup, or sugar and water.

The operation of magnesia may be effectually obtained by giving the patient some lemonade, or a solution of super-tartrate of potassa, as a common drink.

Powder with Calomel and Rhubarb

R. Hydrargyri Chloridi mitis,
Pulveris Rhei, āā, gr. x.
Olei Cinnamomi, gtt. j.
Misce, et signa.—To be given in molasses.

Powder with Jalap and Super-Tartrate of Potassa.

R. Super-tartratis Potassæ, ℥ij.
Pulveris Jalapæ, ℥j.
Misce.—Divide in chartulas vj.—Of these, give one every two hours in molasses or sirup.

Compound Powder of Jalap and Gamboge.

R. Super-tartratis Potassæ, ℥ij.
Pulveris Jalapæ, ℥j.
—— Gambogiæ, gr. vj.
Misce.—Divide in chartulas vj.—Signa.—To be given as directed in the preceding prescription.

The compound powder of jalap has been almost consecrated to dropsical affections; but it forms a very convenient purge in many inflammatory cases; and when we wish to increase its certainty and activity, we may conveniently add the Gamboge, as in the above formula.

Powder of Aloes, Canella Alba, &c.

R. Pulveris Aloes, ℥jss.
—— Canellæ albæ, gr. xvijj.
—— Radicis Serpentariæ virginianæ, gr. vj.
Misce.—Divide in pulveres vj.—Signa.—One every three or four hours in sirup.

This compound, the *hiera picra* of the shops, forms a very good purgative, and is usually employed as an emmenagogue.

Powder with Sulphate of Potassa, &c.

R. Potassæ sulphatis, ℥j.

Pulveris Rhei, ℥iss.

———— Floris Anthemidis, ℥j.

Tere in pulverem, et divide in doses equales vj.

Signa.—Take one twice a day in sugar and water.

In dyspepsia and a sluggish state of the bowels.

Powder of Sulphur und Super-tartrate of Potassa.

R. Sulphuris sublimati, ℥ss.

Pulveris Super-tartratis Potassæ, ℥j.

Misce, et signa.—Mix in a sufficient quantity of molasses, and give a tea-spoonful four or five times a day to children. In some diseases of the skin.

It should be always recollected that sulphur promotes insensible perspiration in a very marked degree, and hence, renders the patient more liable to be affected by the vicissitudes of climate.

Powder with Milk of Sulphur and Calcined Magnesia.

R. Sulphuris præcipitati vel Lactis Sulphuris,

Magnesiae ustæ, āā. ℥ss.

Misce, et signa.—A tea-spoonful to be taken four or five times a-day.

This preparation is better adapted to delicate stomachs than the former. Either of them forms a good aperient during an alterative course of medicine.

Sulphate of Magnesia and Tartrate of Antimony.

R. Sulphatis Magnesiae, ℥j.

Tartratis Antimonii et Potassæ, gr. j.

Misce, et signa.—To be dissolved in six ounces of water, and taken for a dose.

By combining tartar emetic with this and other purgative salts, we promote at the same time a gentle diaphoresis; which, in many slight inflammatory affections, is very desirable.

Powder with Tartrate of Soda, Tartaric Acid, &c. vulgo, Seidlitz Powders.

R. Tartratis Sodæ et Potassæ, ℥ij.

Sodæ Bicarbonatis, ℥ij.

Fiat pulvis.—Signa.—To be dissolved in half a tumbler of cold water.

R. Acici tartarici pulverizati, gr. xxv.

Signa.—Dissolve in an equal quantity of water with the above, and mix them both together. To be taken while in a state of effervescence.

This a very popular purgative, and justly so; it is a gentle aperient, and may be taken by persons whose stomachs would reject any of the ordinary saline preparations.

Phosphate of Soda.

R. Phosphatis Sodæ, ℥j.

This salt so much resembles in taste the common table salt, (muriate of soda,) that it may be conveniently taken in a bowl of broth. It is a mild purgative.

PILLS.

Pills of Calomel, Jalap, &c., (known as Griffiths' Cathartic Pills.)

R. Pulveris Jalapæ,

—— Rhei,

Saponis albi, āā. ℥ss.

Hydrargyri Chloridi mitis, gr. xxv.

Tartratis Antimonii et Potassæ, gr. iss.

Aquæ destillatæ, quantum sufficit—ut fiat massa: divide in pilulas xxv.

Signa.—Two of these pills may be taken at once, and repeated in two hours, if not sufficient.

This combination of purgatives with the tartrate of antimony, forms a most excellent anti-bilious remedy; and, in cases of fever, prepares the alimentary canal for the reception of other remedies better than a more simplified preparation.

Pills of Blue Mass, Aloes, &c.

R. Massæ ex Hydrargyro,

Pulveris Jalapæ,

—— Aloes, āā. gr. xv.

Syrupi simplicis, q. s. ut fiat massa, et divide in pilulas xij.

Signa.—Three may be taken at night, and the same dose repeated in the morning, if the first is ineffectual.

Pills of Blue Mass, Soda, &c.

R. Massæ ex Hydrargyro, gr. ix.

Pulveris Rhei,

Bicarbonatis Sodæ, āā. gr. xij.

Syrupi Rhei aromatici, q. s. ut fiat massa, et divide in pilulas, xij.

Signa.—One, twice or three times a-day, as an alternative in derangement of the liver.

Hartshorne.

Powdered Charcoal.

R. Pulveris Carbonis, ʒj.

Of this a tea-spoonful may be given in milk, molasses, or sugar and water; repeated as often as necessary to keep the bowels in a soluble state.

Charcoal occasionally accumulates in the bowels, and occasions the most distressing constipation. When it fails, therefore, to operate promptly, some other purgative should be given. It may be observed, that charcoal is exceedingly well adapted to many cases of dyspepsia, enabling the patient to digest more, and a greater variety of food, than he can without it. The safest and most effectual mode of using it is to dust it over the aliments in the same way as pepper.

Pills of Calomel, Rhubarb, and Opium.

R. Hydrargyri Chloridi mitis, gr. xij.

Pulveris Rhei, gr. xx.

———— Opii, gr. j.

Misce, et divide in pilulas viij.—Signa.—Four may be taken at once, and the remainder in one hour if the first are insufficient.

This compound may be employed when we wish the effect of calomel, and when there is pain in the bowels; if there should be much costiveness, it may be necessary to follow the pills with a dose of castor oil, or Epsom salts.

Pills of Rhubarb and Soap.

R. Pulveris Rhei, ℥j.

Saponis albi, gr. x.

Aquæ destillatæ, q. s.—ut fiant pilulæ xv.

Signa.—Three for a dose.

To remove costiveness, or produce a gentle evacuation in debilitated habits.

Pills of Aloes and Rhubarb.

R. Pulveris Rhei,

———— Aloes āā. ℥ss.

Saponis albi, quantum sufficit—ut fiant pilulæ xxv.

Signa.—Three or four of these may be taken occasionally, in habitual costiveness and in dyspepsia.

Pills of Rhubarb, with Sulphate of Iron, &c.

R. Pulveris Rhei, ℥iss.

Sulphatis Ferri, ℥ss.

Saponis albi, ℥ij.

Aquæ destillatæ, q. s.—Fiat massa in pilulus xl. dividenda.

The intention of these pills is obviously to remove costiveness, and impart tone to the bowels. They were a favourite medicine with Dr. Griffiths, late of

this city. Three or four may be taken for a dose at bed-time.

Pills with Aloes, Gentian, &c.

R. Pulveris Aloes, ʒj.

Extracti Gentianæ, ʒss.

Olei Carui, gtt. x.

Syrupi simplicis, quantum sufficit.

The mass to be divided into pills, each one weighing gr. iv.—Signa.—Two or three for a dose.

Pills of Croton Oil.

R. Olei Tiglii, gtt. j.

Micæ panis, q. s.—ut fiant pilulæ iv.

Signa.—One every hour until they operate.

This powerful remedy has lately been restored to the materia medica. One drop is a full dose, and frequently less will answer every purpose.

Pill with Aloes, Ipecacuanha, &c.

R. Pulveris Aloes, ʒss.

———— Ipecacuanhæ, gr. iv.

Saponis albi, ʒij.

Syrupi simplicis, q. s.—ut fiant pilulæ xx.

Signa.—Two or three for a dose; and to be repeated when necessary.

These pills, with almost any other of the aloetic preparations, from their specific action on the rectum, will sometimes restore suppressed hæmorrhoids.

Aperient Pills.

R. Aloes socotorinæ, ʒj.

Pulveris Rhei, ʒj.

———— Ipecacuanhæ, gr. vj.

Saponis venetæ ʒj.

To be divided into sixty-four pills.—As a simple aperient, one may be taken every night at bed-time; but where a full purgative effect is desirable, the dose is two or three.

Pills with Asafætida and Aloes.

R. Gummi Asafætidæ,
 Pulveris Aloes,
 Saponis albi, āā. ℥ss.
 Aquæ destillatæ, quantum sufficit—ut fiat massa in pilulas xxv. dividenda.

These pills are usually kept made by the apothecaries, but the physician may prefer them recent, and therefore the formula is here inserted. We may alter this prescription by substituting rhubarb for aloes, if it is deemed preferable. Three or four of them are sufficient for a dose.

Pills with Extract of Butternut, Jalap, &c.

R. Extracti Juglandis cinerei, ℥ss.
 Pulveris Jalapæ, ℥j.
 Saponis albi, gr. x.

Misce.—Fiant pilulæ xv.

Signa.—Three or four of these may be taken for a dose, and if they do not operate, two or more may be safely administered.

The extract of Butternut, either alone or in combination, is very highly recommended as a cathartic in fevers, dysentery, &c.

Pills of Blue Mass, and Colocynth.

R. Massæ ex Hydrargyro Extracti colocynthidis, āā. gr. v.
 Olei Carui, gtt. ij.
 Fiant pilulæ, ij.

These constitute an active cathartic.

Confection of Sulphur.

R. Sulphuris sublimati et loti, ℥iss.
 Confectiones Sennæ, ℥ij.
 Potassæ nitratis, ℥iij.
 Syrupi Aurantii, q. s.—ut fiat confectio.

Signa.—From one or two drachms may be taken once or twice a-day.
Useful in hæmorrhoidal affections.

Pills of Compound Extract of Colocynth and Calomel.

R. Extracti Colocynthidis compositi, gr. xlvij.

Hydrargyri chloridi mitis, gr. xij.

Misce et divide in pilulas xij.

Signa.—One, or at most two of these pills will generally act as a cathartic.

Pills of Compound Extract of Colocynth, &c.

R. Extracti Colocynthidis compositi, ℥j.

——— Jalapæ resinosi, gr. vj.

Pulveris Scammoniae compositi, gr. x.

Hydrargyri chloridi mitis, gr. x.

Tartratis Antimonii et Potassæ, gr. j.

Saponis veneti, gr. v.

Olei Cassiæ, gtt. iv.

Misce et divide in pilulas xv.

Signa.—One, two, or three to be taken at bed-time, for a cathartic.

Dr. Meigs.

Pills with Aloes and Mercurial Blue Mass.

R. Massæ ex Hydrargyro, ℥j.

Pulveris Aloes, gr. xxv.

Misce et divide in pilulas xv.—Signa.—One every two hours until they operate.

In cases of constipation depending upon deficiency of bile, these pills may be advantageously employed.

Pills of Compound Extract of Colocynth, &c. &c., commonly called Fothergill's Pills.

R. Extracti Colocynthidis compositi, ℥iss.

* Oxidi Antimonii illoti, ℥ss.

Misce et divide in pilulas xxx.—One of these pills will frequently produce a free evacuation; but two or more may be taken when required, at bed-time.

* This preparation of antimony was formerly known by the name of Calx Antimonii Illota.

Pills with Aloes, Guaiacum, &c.

R. Pulveris Guaiaci, ℥ss.

—— Aloes, gr. xxxvj.

—— Rhei, ℥ij.

Balsami Canadensis, quantum sufficit—ut fiat massa in pilulas xlvij. dividenda.

Signa.—One to be taken every three hours.

This combination forms a highly stimulating cathartic.

Electuary of Senna and Figs.

R. Pulveris Sennæ, ℥ss.

Pulpæ ficus, quantum sufficit—ut fiat electuarium.

Signa.—A piece, the size of a marble, night and morning, in costiveness from sedentary habits.

MIXTURES, INFUSIONS, &c.

Oleaginous Mixture.

R. Olei Ricini, ℥iss.

Pulveris gummi Arabici, ℥ij.

Sacchari albi, ℥j.

Misce secundum artem, et tunc adde gradatim—

Aquæ Menthæ sativæ, ℥iv.

Tincturæ Opii, gtt. xl vel l.

Signa.—A table-spoonful every hour and a half or two hours, until it purges.

It is of some consequence in preparing this mixture, to shake the oil and powders well together before adding the water, which should be poured in gradually, agitating it after each addition; or, combine the whole in a mortar.

Mixture of Oil of Turpentine, &c.

R. Olei Terebinthinæ rectificati, ℥j.

Pulveris Gummi Arabici,

—— Sacchari albi, āā. ℥ij.

Aquæ Menthæ sativæ, ℥iv.

Misce, et Signa.—A table-spoonful every two hours, until it operates.

This preparation must be made in the same manner as the oleaginous mixture.

Mixture with Magnesia, Tinct. Fœtida, &c.

- R. Magnesiae carebonatis, ℥ss.
 Tincturæ Asafœtidæ, gtt. lx.
 ——— Opii. gtt. xx.
 Sacchari albi, ℥j.
 Aquæ destillatæ, ℥j.

Misce.—Fiat mistura.—Signa.—Twenty-five drops may be given to an infant of two or four weeks old; in flatulent cholic, diarrhœa, &c.

This preparation is strongly recommended and much employed by Dr. Dewees of this city.

If the *calcined* magnesia be used, it unites with the resin of the asafœtida, forms an almost solid mass at the bottom of the vessel, and defeats the object of the prescription.

Mixture of Carb. Magnesia, &c.

- R. Magnesiae carbonatis, ℥ss.
 ——— Sulphatis, ℥iij.
 Spiritus Ammoniae aromatici, ℥j.
 Tincturæ Rhei, ℥ss.
 ——— Hyosciami, ℥ss.
 Aquæ Menthæ sativæ, ℥iv.

Fiat mistura.—Signa.—A large table-spoonful to be taken two or three times a-day.

Dr. Meigs.

Oleaginous Mixture with Oil of Turpentine.

- R. Olei Ricini, ℥iss.
 Vitellum ovi, j.
 Sacchari albi, ℥ij.
 Olei Terebinthinæ rectificati, ℥ss.
 Aquæ Menthæ sativæ, ℥iv.

Misce optime et fiat mistura.—Signa.—A large table-spoonful may be taken every hour, according to circumstances.

This preparation is much used by some of our best practitioners, in cases requiring a brisk and stimulating

cathartic. It has also been recommended as a purgative, after the effects of an over-dose of opium have been partially subdued by the proper preliminary remedies.

Mixture of Oil of Euphorbia Latyris, or Oil of Spurge.

R. Olei Euphorbiæ latyris, gtt. viij.

Gummi Acaciæ,

Sacchari albi, āā. ʒj.

Aquæ Menthæ sativæ, ʒj.

Fiat mistura.—Signa.—One half of this mixture may be given at once, and repeated in two hours if the bowels are not moved.

The oil of spurge is an old remedy, lately revived and introduced to notice by the Italians. The European oil is stated to be as mild as castor oil, and as prompt and efficacious as the croton oil, in the above doses. The oil prepared from the beans grown in this country, is said to be much more acrid, occasioning both nausea and griping.

Mixture of Croton Oil.

R. Olei Tiglii, gtt. j.

Mucilaginis acaciæ,

Aquæ destillatæ, āā. ʒj.

Misce et Signa.—To take a tea-spoonful every two hours until it operates.

Saponaceous Solution of Croton Oil.

R. Olei Tiglii, gtt. viij.

Potassæ puræ, gr. vj.

Aquæ destillatæ, ʒij.

Fiat solutio.—Signa.—From three to six drops may be given for a dose.

This is a substitute for the soap of the croton oil proposed by the French, and which is more troublesome to make.

Mr. Conevel states, that four drops of this oil rubbed on and round the umbilicus, will produce a cathartic

effect; even the odour of it has been known to excite the alimentary canal, either as an emetic or purgative.

Solution of Epsom Salts with Nitric Acid.

R. Sulphatis Magnesiae, ℥ss.

Tincturæ Jalapæ, ℥j.

Acidi nitrici, gtt. x.

Aquæ Menthæ, ℥ij.

Misc.—Signa.—To be taken for a draught. The nitric acid is said to destroy in great measure the bitter taste of the salt.

Infusion and Tincture of Senna, &c.

R. Infusionis Sennæ, ℥vj.

Tincturæ ———, ℥ij.

Mannæ, ℥ij.

Magnesiae sulphatis, ℥ss.

Aquæ Menthæ sativæ,

—— destillatæ, āā. ℥iss.

Misc.—Signa.—One half to be taken for a dose, and repeated if it does not operate. This was a very common prescription of the late Mr. Abernethy, of London.

Purgative Draught.

R. Foliorum Sennæ, ℥ij.

Sulphatis sodæ, ℥ss.

Aquæ, ℥iv.

Syrupi Rhamni, ℥j.

The water should be added boiling hot, and the infusion then cooled and strained. A few grains of Jalap may be substituted for the Sirup of Buckthorn.

Mixture of Magnesia, Rhubarb, &c.

R. Magnesiae ustæ, ℥ss.

Pulveris Rhei, gr. vj.

—— Sacchari albi, ℥j.

Essentiæ Menthæ piperitæ, gtt. vj.

Aquæ destillatæ. ℥iss.

Fiat mistura.—Signa.—A tea-spoonful to be given every two hours to children, until it operates.

This preparation is designed for the bowel complaints of children, in which it is very serviceable. Laudanum can be added according to the age and circumstances of the patient.

Solution of Sulphate of Magnesia, &c.

R. Sulphatis Magnesiae, ℥j.
 Tartratis Antimonii, gr. j.
 Succī Limonis recentis, ℥j.
 Aquæ destillatæ, ℥iij.

Misce.—Fiat solutio.—Signa.—A table-spoonful to be given every two hours until it operates.

Infusion of Senna, Sulphate of Magnesia, &c.

R. Foliorum Sennæ, ℥vj.
 Mannæ optimæ,
 Sulphatis Magnesiae, āā. ℥j.
 Seminum Cardamomi contus: ℥ij.

Misce, et signa.—To be infused in a pint of boiling water for thirty or fifty minutes, and a tea-cupful given every hour until it operates.

Infusion of Senna.

R. Foliorum Sennæ, ℥vj.
 Seminum Cardamomi contus: ℥ij.

Misce, et signa.—To be infused in a pint of boiling water for half an hour; one-half to be taken at once; if insufficient, the remainder in two or three hours.

This infusion may be longer kept if a little tinctura sennæ be added to it.

Decoction of Prunes.

R. Fructus Pruni, ℥iij.
 Aquæ, ℥iij.

Decoque ad libras duas.—Used as a mild laxative.

Infusion of Senna with Rhubarb, &c.

R. Foliorum Sennæ, ℥vj.
 Mannæ optimæ, ℥j.
 Radicis Rhei contusæ, ℥ij.
 Seminum Cardamomi contus: ℥ij.

Misce, et signa.—To be infused in a pint of boiling water for one hour. A tea-cupful given every hour till it operates.

Compound Tincture of Senna, vulgo, Warner's Cordial.

R. Tincturæ Sennæ compositæ, ℥j.

Direct a table-spoonful, mixed with double the quantity of water, as hot as it can be drunk, to be taken for a dose, in cases of violent pain of the stomach, where a cordial purgative is required; to be repeated *pro re nata*.

Decoction of Aloes, &c.

R. Extracti Glycyrrhizæ, ℥ss.

Potassæ subcarbonatis, ℥ij.

Aloes Spicatæ extracti contriti,

Myrrhæ contritæ,

Croci stigmatum, āā. ℥j.

Tincturæ Cardamomi compositæ, ℥iv.

Aquæ destillatæ, Oj.

Boil down all the ingredients, except the last, with the water, to twelve fluid ounces, and strain; then add the compound Tincture of Cardamom. The dose of this decoction is from half an ounce to an ounce and a half, and is of service, frequently in dyspepsia, and wherever a mild warm cathartic is required.

Wine of Colchicum.

R. Vini Colchici saturati, ℥ss.

Signa.—Twenty or thirty drops of this may be taken in sugar and water. In cases of rheumatism or gout.

The above preparation of the Colchicum or meadow saffron is made both of the bulb and seeds, but is most powerful when the bulb is used. The ordinary dose is twenty drops three times a-day, which may be repeated until it affects the *stomach, pulse, or bowels*, when it should be immediately discontinued, and its use resumed with caution.

Colchicum and Magnesia.

R. Magnesiæ ustæ, ℥j.

Sacchari albi,

Gummi Arabici, āā. ℥ss.

Aquæ destillatæ, ℥iv.

Vini Colchici, gtt. xl.

Fiat mistura.—Signa.—To take a table-spoonful every two hours until it operates.

This preparation is much used in gouty and rheumatic affections.

Scuddamore's Mixture of Colchicum and Epsom Salts.

R. Magnesiae Sulphatis, $\bar{\text{z}}$ j ad $\bar{\text{z}}$ ij.

Aquæ Menthae, $\bar{\text{z}}$ x.

Aceti colchici, $\bar{\text{z}}$ j ad $\bar{\text{z}}$ iss.

Syrupi Croci, $\bar{\text{z}}$ j.

Magnesiae, D vij.

Fiat mistura.—Dose, one, two or three table-spoonsful, repeated every two hours, until four or six evacuations are produced in twenty-four hours. Chiefly used in gouty paroxysms.

Pills of Veratrina.

R. Veratrinæ, gr. ss.

Mucilaginis Acaciæ, q. s.—ut fiant pilulæ vi.

Signa.—One of these may be given, and if purgative effects do not follow, three may be given in one day.

The active principle of this substance has been separated, and the name of Veratrina bestowed upon it. It is extremely poisonous, and, therefore, to be taken with great caution. Its only recommendation is, that it is more certain in its operation than the preparations noticed above.

Tincture of Veratrina.

R. Veratrinæ, gr. iv.

Alcohol, $\bar{\text{z}}$ j.

Fiat tinctura.—Signa.—This may be given in doses of ten, fifteen, twenty, or twenty-five drops, in a cup of drink. It is designed as a substitute for the Vinous Tinct. of Colchicum, in dropsies, gout, rheumatism, &c.

The Tincture of Veratrina is recommended as an external application in the diseases above mentioned. For this purpose also the ointment is proposed.

Solution of Sulphate of Veratrine.

R. Sulphatis Veratrinæ, gr. j.

Aquæ destillatæ, ℥ij.

Fiat solutio.—Dose.—A dessert-spoonful. This solution is said to be a good substitute for the Eau Medicinale of Husson, in gouty affections.

Majendie.

Ointment of Veratrina.

R. Veratrinæ puræ, gr. iv.

Adipis suillæ, ℥j.

Fiat unguentum.—Signa.—To be employed externally, in the cases above directed.

Majendie.

There is obtained from the Veratrum another alkaline substance called *Sabadillin*, which possesses the properties of Veratrina, but in a less degree. It has consequently been little used in practice.

Tincture of Aloes, &c.

R. Pulveris Aloes socotrinæ,

——— Seminum Anisi, āā. ℥ij.

Spiritus vini gallici, Oij.

Fiat tinctura.—Dose.—From one-third to half a wine-glassful, three times a-day.

The above is highly spoken of by Dr. R. Pindall in hemorrhoidal affections. He considers it almost a specific, notwithstanding the prejudice against aloes, in these affections of the rectum.

Mixture of Elaterium.

R. Extracti Elaterii, gr. j.

Spiritus ætheris nitrisi, ℥ij.

Tincturæ Scillæ,

Oxymellis Colchici, āā. ℥ss.

Syrupi Simplicis, ℥j.

Fiat mistura.—Signa.—To take a tea-spoonful three or four times a day in a little water.

The above is the original prescription of Dr. Ferriar, who recommends it in the several forms of dropsy, and particularly in Hydrothorax.

Tincture of Aloes and Canella Alba, vulgo, Tincture of Hiera Picra.

R. Tincturæ Hieræ picræ, ℥j.

A tea-spoonful may be taken every hour, or two hours, until it operates.

Electuary with Jalap, Nitrate of Potash, &c.

R. Pulveris Super-tartratis Potassæ,

———— Radicis Jalapæ,

———— Nitratis Potassæ, āā. ℥ss.

Confectionis Sennæ, ℥j.

Syrupi simplicis, q. s.—ut fiat electuarium.—Of which a bolus the size of a hazlenut may be taken three or four times a-day.

This preparation once gained considerable celebrity as a popular remedy for hemorrhoids.

Electuary with Sulphur.

R. Sulphuris sublimati, ℥ss.

Confectionis Sennæ, ℥iss.

Syrupi simplicis, q. s.—ut fiat electuarium.—A tea-spoonful night and morning for a dose, in the disease above mentioned.

ENEMATA.

Enemata, used either as auxiliaries to or substitutes for cathartics, very properly succeed that class of remedies.

They are employed to fulfill other indications, and the different preparations will be noticed under their appropriate heads. It is usual to direct three times the quantity to be given for an enema that is required for a dose *per os*. For the purpose of procuring a discharge from the bowels, this rule may generally be observed; but in some of the powerful remedies occasionally administered, as opium and tobacco, it might be attended with serious consequences.

The following are the proportions recommended by Professor Chapman in the ordinary enemata for persons of different ages:—A child between the ages of one and five years old, from ℥iv. to ℥vj. A youth of

ten or fifteen years, half a pint; and an adult, from a pint to a quart, according to circumstances.

Enema with Oil, Salt, &c.

R. Olei Ricini vel Olivarum, ℥ij.

Syrupi sacchari rubri, ℥ij.

Sodii chloridi, ℥ss.

Infusionis Lini tepidæ, Oj.

Fiat enema.—One half to be thrown up the rectum with a syringe, and the remainder in half an hour or an hour, if the first produces no evacuation.

Enema with Sulphate of Magnesia, &c.

R. Sulphatis Magnesiae, vel Sodæ, ℥ij.

Olei olivarum, ℥j.

Solutionis Amyli tepidæ, Oj.

Misce.—Fiat enema.—Signa.—To be exhibited in the manner directed above.

Enema of Colocynth.

R. Colocynthidis medullæ incisæ, ʒj.

Aquæ bullientis, ℥x.

Sodæ muriatis, ℥ss.

Mannæ optimæ, ʒj.

Boil the Colocynth in the water for ten or fifteen minutes, and strain—then add the other ingredients. To be employed in cases of obstinate constipation.

Enema of Oil of Turpentine.

R. Olei Terebinthinæ optimi, ʒjss.

Vitelli unius ovi,

Solutionis gummi Arabici, vel

Infusionis seminum Lini tepidi, Oj.

Misce.—To be administered secundum artem.

The Venice turpentine may be substituted for the oil directed in this prescription; and either of them, if required, may be given in larger quantities.

Enema of Infusion of Senna, &c.

R. Infusionis foliorum Sennæ, Oj.

Sulphatis Magnesiæ, ℥ij.

Misce.—One-half to be exhibited as an enema, and the remainder if necessary.

Enema with Tartrate of Antimony.

R. Tartratis Antimonii et Potassæ, ℥j. to ℥ij.

Solutionis gummi Arabici tepidæ, Oj.

Misce.—Exhibit as the former.

Dr. Chapman recommends this preparation as one admirably calculated to overcome constipation, which is so obstinate as to resist the ordinary remedies. It is very powerful, and requires care in its management.

Tobacco Enema.

R. Foliorum Tabaci, ʒj.

Aquæ ferventis, Oj.

Make an infusion in the usual manner, and use the half as an enema.

It must be cautiously administered, as death has followed its imprudent exhibition. In case of great prostration, the rectum should be at once washed out with copious injections of water or other simple diluents, accompanied by the internal use of stimulating drinks, and the application of a mustard plaster over the region of the stomach.

CLASS III.

DIURETICS.

THE rules for the administration of this class of remedies are few and simple.

Where there is much arterial excitement, a judicious use of the lancet should precede the employment of diuretics.

The patient must be kept cool, and every precaution taken against perspiration; for these two secretions are well known to be antagonizing functions.

In order to ensure the full effects of diuretics, diluting drinks should be freely given, and especially those containing some diuretic in solution.

POWDERS AND PILLS.

Powder of Carbonate of Potash.

R. Carbonatis potassæ, ℥ss.

This powder may be most agreeably taken in a bottle of seltzer water, and repeated three or four times a-day.

The same quantity of carbonate of soda may be administered in the same manner.

Powder of Squill, &c.

R. Pulveris scillæ, gr. xij.

—— Nitratis Potassæ, ℥j.

Fiant pulveris vj.—Signa.—One to be given twice a-day in sugar and water.

Some aromatic, such as cinnamon, may be added to this preparation, if the stomach should be disposed to reject it.

Powders of Uva ursi, and Soda.

R. Pulveris Uvæ ursi, ℥iss.

Sodæ Bicarbonatis, ℥j.

Misce, et divide in chartulas xij.—Signa.—One three times a-day in sugar and water.

Useful in Nephritic complaints, and other affections of the bladder.

Pills of Digitalis, Calomel, Opium, &c.

R. Pulveris Digitalis, gr. xij.

Hydrargyri chloridi mitis, gr. iv.

Opii, gr. iv.

Confectionis rosæ, q. s.

Ut fiant pilulæ xij.—Signa.—One to be taken every eight hours.

Useful in Hydrothorax and Ascites, depending on visceral obstruction.

Pills of Balsam of Copaiva.

R. Balsami Copaivæ, ℥ij.

Magnesiæ ustæ gr. vij.

Misce in mortario.—

Copaiva requires for its solidification one-seventeenth its weight of perfectly pure magnesia. It will not, however, always become solid in these proportions, as the balsam suffers changes by age. The mixture thickens in about six or eight hours, and acquires greater solidity by time. The mass may be formed into pills of four grains each, of which two may be taken three or four times in the day. They answer

all purposes of the liquid balsam administered by the mouth.

Powders of Ergot and Cubebs.

R. Secalis cornuti, ℥ij.

Pulveris Cubebæ, ʒj.

————, Cinnamomi, ʒss.

Sacchari purificati, ʒj.

Divide into eight powders, of which one may be given three or four times a-day. In Leucorrhœa and Gleet. *Dr. Ryan.*

Pills of Extract of Dandelion.

R. Extracti Taxacaci, ʒss.

Pulveris Glycyrrhizæ, q. s.

Ut fiant pilulæ x.—One to be given morning, noon, and night. In Dropsy, and diseases of the urinary apparatus.

Extract of Dandelion and Blue Mass.

R. Extracti Taxacaci, ʒss.

Massæ ex Hydrargyro, gr. v.

Pulveris uvæ ursi, q. s.

Ut fiant pilulæ x, et signa; one to be taken morning, noon, and night. In Dropsy complicated with disease of the liver.

Pills of Squill and Calomel, &c.

R. Hydrargyri chloridi mitis.

Pulveris Scillæ, āā. gr. xij.

Conservæ rosarum, q. s.—ut fiant pilulæ xij.—Signa.—One night and morning, in dropsies depending upon visceral derangement.

This preparation is one of the most celebrated of the diuretics; and where disease of the liver causes the effusion, is decidedly one of the best. When the calomel affects the mouth, it may be omitted, and the squill only continued.

Pills of Sub-Carbonate of Soda.

R. Sub-carbonatis Sodæ exsiccata, ʒj.

Saponis albi, ℥ij.

Aquæ fontanæ, q. s.—Ut fiant pilulæ xxx.—Three to be taken three times a-day. Chiefly useful in calculous affections.

Pills of Calomel, Digitalis, &c.

R. Pulveris Scillæ exsiccatae, gr. iv,

———— Myrrhæ, ʒj.

———— Digitalis foliorum, gr. x.

Hydrargyri chloridi mitis, gr. vj.

Simul tere et adde—

Asafœtidæ, ʒss.

Extracti Gentianæ, q. s.

Fiat massa, in pilulas xv. dividenda.—Signa.—One night and morn-
ing. *Paris.*

Turpentine Pills.

R. Terebinthinæ albæ, ʒj.

Divide in pilulas xv.—Signa.—One four or five times a-day.

MIXTURES, INFUSIONS, &c.

Mixture with Tincture Cantharides, &c.

R. Tincturæ Cantharidum,

Spiritus ætheris nitrici, aa. gtt. lx.

Sacchari albi, ʒj.

Aquæ Menthæ sativæ, ʒij.

Fiat mistura.—Signa.—A tea-spoonful may be taken every three or
four hours; desisting, if symptoms of strangury should supervene.

Mixture of the Spirits of Nitre, &c.

R. Spiritus ætheris nitrici, ʒj.

Tincturæ Scillæ, ʒij.

Fiat mistura—of which give a tea-spoonful five or six times in the
day, with the following infusion:—

R. Baccarum Juniperi, contus. ʒj.

Potassæ super-tartratis, ʒss.

Aquæ bullientis, Oj.

Fiat infusio.—A tea-cupful for a dose with the above mixture.

Solution of Cream of Tartar.

R. Potassæ super-tartratis, ʒj.

Aquæ destillatæ, Oij.

Fiat solutio.—Signa.—To be freely drunk through the day.

Mixture of Acetate of Ammonia, &c.

R. Liquoris Ammoniaë acetatis, ℥ij.

Potassæ acetatis, ʒij.

Misce, et signa.—A table-spoonful every three hours for a dose.

Infusion of Parsley.

R. Apii petroselini, ʒj.

Aquæ bullientis, Oj.

Fiat infusio, et cola.—Signa.—A tea-cupful may be taken with a tea-spoonful of spiritus nitri dulcis, every three or four hours during the day.

A popular remedy in the strangury arising from blisters.

Mixture of Tincture of Jalap, &c.

R. Tincturæ Jalapæ, ℥iij.

Aceti Scillæ, ʒj.

Aquæ Menthæ piperitæ, ʒj.

Fiat haustus.—To be taken three times a-day.

Infusion of Juniper Berries, &c. in Cider.

R. Baccæ Juniperi contusæ,

Seminum Sinapæos,

Radiciſ Zingiberis, āā. ʒss.

——— Armoraciæ contusæ,

——— Apii petroselini, āā. ʒj.

Succi expressi et fermenti pomorum, Oij.

Fiat infusio.—Of which a wine-glassful may be taken three or four times a-day.

The above compound is recommended by Dr. Parrish, in those cases of general dropsy which admit of stimulation, and in which the urinary secretion is difficult to promote. He thinks he has found it of signal advantage. The cider should be *old* and *sound*.

Oil of Juniper Berries.

R. Olei Juniperi optimi, ℥ss.

Signa.—Five drops to be taken every three hours in sugar, gradually increasing the dose.

Oil of Juniper, &c.

R. Olei Juniperi, gtt. viij.

Sacchari albi,

Gummi arabici, āā. ℥iss.

Aquæ destillatæ, ℥iv.

Misce et signa.—A table-spoonful to be taken every hour or two hours.

Infusion of Juniper Berries, &c.

R. Baccæ Juniperi contusæ, ℥iss.

Super-tartratis potassæ, ℥j.

Aquæ bullientis, Oj.

Fiat infusio.—Of this a tea-cupful may be taken four or six times a-day; adding to each a tea-spoonful of the sweet spirits of nitre. In general and local dropsy.

Mixture of Colchicum, Squills, &c.

R. Oxymellis Colchici,

——— Scillæ,

Tincturæ Nicotianæ, āā. ℥ss.

M. Signa.—Take a tea-spoonful in a little water four times a-day.

If requisite, the Spiritus ætheris nitrici may be added in the same quantity. This forms the *diuretic drops* of Dr. Ferriar.

Infusion of Digitalis, &c.

R. Foliorum Digitalis, ℥j.

Aquæ bullientis, Oss.

Macerate for three hours, then strain.—Dose for an adult, a table-spoonful three times a-day.

The prostrating effects of Digitalis should always be borne in mind by the practitioner: and the rule that

when it acts upon the *stomach*, the *bowels*, the *pulse*, or the *kidneys*, it should be discontinued, cannot, when we consider the power of the article, be too scrupulously observed.

Mixture of Digitalis, Potash, &c.

R. Infusionis Digitalis, ℥iv.

Tincturæ Digitalis, ℥i.

Acetatis Potassæ, ℥j.

Tincturæ Opii, gtt. x.

Fiat mistura.—Signa.—A table-spoonful three or four times a-day.

Mixture of Spirits of Turpentine, &c.

R. Olei Terebinthinæ, gtt. C.

Pulveris gummi Arabici,

———— Sacchari albi, āā. ℥j.

Aquæ Menthæ sativæ, ℥iv.

Fiat mistura.—Signa.—A table-spoonful four or five times a-day, aided by some diuretic drink.

This remedy is used in nephritic complaints with great benefit; and in the passage of calculi down the ureters often affords great relief.

Diuretic Liniment.

R. Tincturæ Scillæ,

———— Digitalis, āā. ℥ij.

Misce.—Employed with good effect as a liniment in symptomatic anasarca.

M. Rullier.

Decoction of Cainca.

R. Radicis Caincæ, ℥ij.

Aquæ, Oiss.

To be gradually reduced by heat to one half. Dose—a table-spoonful three times a-day. *In Dropsy.*

This root is derived from the *Chiococca anguifuga* a Brazilian plant, which has of late been extensively and efficaciously employed in Europe. It occasions watery evacuations and diuresis, whence its use in dropsy.—See *Dunglison, New Remedies*, p. 108.

Mixture of Carbonate of Potash, Squills, &c.

R. Carbonatis Potassæ, ℥j.

Succi limonis, q. s. ad saturandum,

Tincturæ Scillæ, ℥ss.

———— Opii, gtt. xxx.

Aquæ Cinnamomi, ℥iv.

Misce.—Fiat mistura.—Signa.—A table-spoonful every two or three hours throughout the day.

The vinegar or oxymel of squills may be substituted in this preparation, if deemed preferable. We selected the tincture, as being more uniform and least liable to spontaneous changes.

Mixture of Wine of Colchicum, &c.

R. Vini Colchici, ℥ij.

Magnesiæ carbonatis, ℥j.

Aquæ Cinnamomi,

——— destillatæ, āā. ℥iij.

Misce, et signa.—A table-spoonful may be taken every three hours.

The vinum colchici is recommended by late European writers as very beneficial in that particular form of Dyspepsia depending upon duodenal irritation; observing at the same time to employ occasional laxatives.

Tincture of Colchicum and Digitalis.

R. Tincturæ seminum colchici,

———— Digitalis, āā. ℥ij.

Spiritus ætheris nitrici, ℥j.

Misce.—Dose—twenty drops on a lump of sugar. Given as a diuretic in Dropsy.

Infusion of Uva Ursi.

R. Uvæ Ursi foliorum, ℥j.

Aquæ bullientis, Oj.

Pour the water boiling hot on the leaves, and let them macerate for twelve hours. To be taken cold, from half a pint to a pint, daily. Chiefly used in gonorrhœa and irritable bladder.

Mixture of Balsam Copaiba.

R. Balsami Copaibæ,
 Spiritus ætheris nitrici, āā. ℥ss.
 Pulveris gummi Arabici,
 ——— Sacchari albi, āā. ℥j.
 Spiritus Lavendulæ compositi, ℥ij.
 Tincturæ thebaicæ, ℥j.
 Aquæ destillatæ, ℥iv.
 Fiat mistura secundum artem.—Signa.—A table-spoonful three times
 a-day.

This remedy is more especially used in gonorrhœa, and is considered by Professor Chapman as a specific in that disease. A mode, perhaps, more elegant than any other of exhibiting balsam copaiva, consists in dropping the quantity prescribed for a dose in a wine-glassful of water, and then adding to it a small quantity of the Tinc. Gentianæ comp. The balsam collects into a globule, and is surrounded by the tincture, which, leaving its taste in the mouth, covers entirely that of the nauseous balsam.

Mixture of Balsam Copaiva, Cubebs, &c.

R. Balsami Copaibæ,
 Pulveris Cubebæ,
 Mucilaginus gummi Arabici,
 Tincturæ Opii camphoratæ, āā. ℥ss.
 Sacchari albi, ℥ij.
 Aquæ destillatæ, ℥viij.
 Miscæ secundum artem et signa.—A table-spoonful every three hours
 for a dose.

We think we have found this remedy, in some obstinate cases of gonorrhœa, more efficacious than the preceding formula. The tincture of cubebs may be substituted for the powder—and the tinc. opii acetat. ℥j. for the paregoric, if desirable. Balsam copaiva is sometimes exhibited in the form of enema, for the cure of gonorrhœa, by the French physicians.

Copaiba Bolus.

R. Balsami Copaibæ, ʒij.

Gummiæ Arabicæ, ʒj.

Pulveris glycyrrhizæ, q. s.

To be made into a bolus, one-half to be taken in the morning, the other in the evening.

This formula is much in use in the Parisian Hospitals, for obstinate gonorrhœa and gleet which resist the ordinary treatment.

Pills of Copaiba, Calomel, &c.

R. Conservæ rosarum, ʒj.

Resinæ pterocarpi, ʒj.

Hydrargyri chloridi mitis, gr. xviii.

Balsami Copaibæ, ʒij.

Misce.—To be divided into pills of six grains each, and one, two, or three to be taken morning and evening, according to circumstances. They are employed against long-continued gonorrhœa or gleet, when the inflammatory symptoms have been removed, and the discharge is owing to an atonic state of the mucous membrane.

Ratier.

Compound Mixture of Copaiba, &c.

R. Balsami Copaibæ,	}	āā. ʒss.
Aquæ menthæ,		
Alcoholis rectificati,		
Syrupi simplicis,		
Spiritus ætheris nitrici,		gtt. xviii.

Fiat mistura.—To be taken during the day in four doses, followed by demulcent drinks. In chronic gonorrhœa.

Hydragogue Draught of Dr. Ferriar.

R. Gambogiæ, gr. iv.

Spiritus Ætheris nitrici, ʒj.

Tincturæ Sennæ, ʒij.

Syrupi Rhamni,

Aquæ Menthæ, āā. ʒss.

Misce.—Dr. Ferriar observes, that in cases of hydrothorax where there was imminent danger of suffocation, he has afforded immediate relief by the above draught.

Infusion of Dandelion, &c.

R. Infusionis Taraxaci, ℥iv.

Extracti Taraxaci, ℥ij.

Carbonatis Sodæ, ℥ss.

Tartratis Potassæ, ℥iij.

Tincturæ Rhei, ℥iij.

———— Hyosciami, gtt. xx.

Fiat mistura.—Signa.—One third part to be taken three times a-day

In dropsical and visceral affections.

Dr. Meigs.

Decoction of Pipsisseway.

R. Pyrolæ umbellatæ, ℥j.

Aquæ, Oij.

Macerate for twelve hours, and then gradually boil it down to one half. Dose—a wine-glassful three or four times a-day.—In Dropsy

Spirituous Decoction of Pipsisseway.

R. Pyrolæ umbellatæ, ℥j.

Aquæ, ℥xij.

Evaporate by heat to ℥vj.—Then add two ounces of Gin. Dose—a dessert-spoonful every two or three hours, with diluent drinks.—

In Dropsy accompanied by debility.

CLASS IV.

ANTILITHICS AND LITHONTRIPTICS.

ANTILITHICS are those substances which arrest or prevent the formation of calculi in the kidneys or bladder. Lithontriptics are supposed to possess the power of dissolving calculous concretions, by acting on the substance which serves as the bond of union to the saline or acid particles composing the stone.

To administer these remedies with any degree of success, requires a chemical knowledge of the ingredients which enter into the formation of the different calculi; attention must also be given to the properties of the urine.

As derangement of the digestive organs almost always co-exists with these foreign bodies in the urinary apparatus, we shall gain a very considerable advantage by combining the remedies of this class with those which have a tendency to restore the healthy action of the stomach and bowels.

When the Lithic or Uric acid abounds in the urine, the alkalies will be found to produce the greatest benefit; but where the salts are evidently of the calcareous or magnesian character, the muriatic, or some one of the mineral acids, will be the most appropriate remedy.

POWDERS.

Powders of Uva Ursi, Bark, &c.

R. Pulveris Uvæ ursi,
 ——— Cinchonæ, ʒā. ʒij.
 ——— Opii, gr. iij.

Misce, et divide in chartulas vj.—Of which one may be taken three or four times a-day.

The above is the original prescription of Dr. Ferriar, who advises the patient to drink two ounces of aqua calcis after each dose.

PILLS.

Soap Pills.

R. Saponis albi, ʒj.

Divide in pilulas xv.—Signa.—One every two hours during the day.

Pills of Sub-carbonate of Soda, dried.

R. Pilulæ sub-carbonatis Sodæ exsiccatae, xij.

One three or four times a-day.—As under Diuretics.

SOLUTIONS, MIXTURES, &c.

Solution of Bi-carbonate of Soda, &c.

R. Sodæ Bi-carbonatis, ʒj.

Infusionis Quassiae, ʒiv.

Tincturæ Calumbæ, ʒj.

Misce.—Signa.—A table-spoonful for a dose four times a-day.

Solution of Sal Æratus.

R. Bi-carbonatis Potassæ, gr. x vel xv.

Aquæ Seltzer, ʒvj.

Misce.—Draught, to be taken three or four times a-day.

Mixture of Bi-carbonate of Soda, &c. &c.

R. Sodæ bi-carbonatis, ʒj.

Misturæ amygdalæ, ʒiv.

Balsami Copaibæ, ʒij.

Tincturæ Opii, gtt. lx.

Fiat mistura secundum artem.—Signa.—A table-spoonful *pro re nata*.—

In acute pain.

In combination with all these alkaline preparations, demulcent drinks may, with benefit, be freely employed.

Lime Water, &c.

R. Carbonatis Potassæ, ℥ij.

Aquæ Calcis, Oij.

Fiat solutio.—Of which a wine-glassful may be taken every two hours, mixed with an equal quantity of new milk.

Mixture of Magnesia, &c.

R. Magnesiæ carbonatis, ℥j.

Infusionis Gentianæ, ℥vj.

Fiat mistura.—A wine-glassful to be taken three times a-day.

This remedy is recommended by Brande, of London, as superior to the alkalies, in cases where the uric acid abounds.

Mixture of Muriatic Acid in Barley Water.

R. Acidi muriatici, gtt. lx.

Decocti hordei, ℥vj.

Misce.—Signa.—Dose, an ounce three or four times a-day.—In cases of stone, where the calcareous properties are most apparent.

The nitric acid has been given with the same intention, but appears to be less useful than the muriatic.

Oil of Turpentine.

R. Olei Terebinthinæ, ℥ss.

Signa.—From thirty drops to a large tea-spoonful may be taken on sugar, and repeated *pro re nata*. See also the formula given under Diuretics.

This medicine has been frequently given of late in the passage of urinary calculi down the ureters. It is stated by Dr. Physick, (than whom there can be no higher authority,) that this remedy, proportioned in its dose to the degree of suffering, affords more prompt relief than any other.

CLASS V.

DIAPHORETICS.

DURING the exhibition of this class of medicines the patient must be confined to his bed; and when diaphoresis is once excited, every precaution must be taken lest it be suddenly checked.

Venesection should generally be resorted to when there is much arterial excitement with great heat of the skin, before we attempt to give diaphoretics. Where the skin continues steadily hot and dry, all external warmth must be removed, and cold water freely applied to the temples, hands, feet, &c., while the room is to be kept cool by a free current of fresh air. Under these circumstances, cold water may be frequently and plentifully drunk by the patient. Diaphoretics given in conjunction with these general means, will act with much more certainty than if great heat of the surface is maintained by covering the patient with heavy bed-clothes. If it be desirable to produce perspiration when the heat of the skin is not above the natural standard, we shall obtain our object most readily by placing the patient in a *warm* bed, giving him *warm* drinks freely, placing his feet in *warm* water, or applying hot bottles, or bags of sand, to the different parts of his body. The bags of sand accommodate themselves to the inequalities of the surface, and are therefore an excellent means of applying heat externally. Warm bricks, covered with cloths, and

wetted with vinegar, may also be used advantageously. Employed in this way they form a steam-bath. The same effect may be produced by conducting the vapour of alcohol through a tube, under the bed-clothes of the patient.

In few words, then, we have described the manner of producing diaphoresis under opposite states of the system. In the first instance we solicit, in the second we rather extort it.

It may be proper to observe, that in the febrile affections the frequent change of linen about the patient is not only warrantable, but absolutely necessary, whilst employing this class of remedies. Where, however, we are desirous of producing what is emphatically termed "*a sweat*," we should envelope the patient in a flannel shirt or gown, and avoid, as was before observed, every cause which may suddenly check the *cutaneous discharge*, and more especially purging and diuresis.

POWDERS.

Powders with Nitrate of Potash, Calomel, &c.

R. Pulveris nitratis Potassæ, ℥j.

Tartratis Antimonii, gr. j.

Hydrargyri chloridi mitis, gr. vj.

Misce.—Fiant pulveres vj.—Signa.—One to be taken every two hours in sirup or molasses. In febrile affections.

The calomel may be omitted or employed, at the discretion of the practitioner.

Powders with Opium, Ipecacuanha, Calomel, &c.

R. Pulveris Opii, gr. iij.

—— Ipecacuanhæ, gr. vj.

Hydrargyri chloridi mitis, gr. iss.

Nitratis Potassæ, ℥ss.

Misce.—Fiant pulveres vj.—Signa.—One to be taken every three or four hours.

Powders with Guaiacum, &c.

R. Pulveris Guaiaci.

——— nitratis Potassæ, āā. ʒj.

——— Ipecacuanhæ, gr. iij.

——— gummi Opii, gr. ij.

Misce.—Fiant pulveres vj.—Signa.—One to be taken every three hours in a little jelly or molasses.

The last prescription is a stimulating diaphoretic, and is particularly serviceable in acute rheumatism, after the reduction of inflammatory symptoms.

Powders of Sulphate of Potash, Opium, &c. vulgo, Dover's Powders.

R. Pulveris Opii,

——— Ipecacuanhæ, āā, gr. vj.

Potassæ sulphatis, gr. xlvij.

Misce.—Fiant pulveres vj.—Signa.—One to be taken three times a-day.

The principal use of the salt in this preparation, appears to be to facilitate the minute division of the other two ingredients. Its crystals are extremely hard, and it should therefore be broken down and reduced to powder in a mortar, at the same time with the other articles. It is one of the happiest and most useful combinations, and has stood the test of time and experience with undiminished reputation. To promote its operation, *warm drinks* may be freely taken after each dose.

Powders with Calomel, Opium, &c.

R. Pulveris Opii, gr. iv.

Hydrargyri chloridi mitis, gr. xvj.

Pulveris Ipecacuanhæ, gr. viij.

Misce.—Divide in pulveres viij.—Signa.—One to be taken every hour or two hours.

This combination is recommended by Professor Chapman in dysentery. From its evident diaphoretic tendency, we have been induced to place it under this head.

Antimonial Powder.

R. Oxidi Antimonii cum phosphate Calcis, $\mathfrak{z}\text{j}$.

Divide in chartulas vj.—Signa.—One every two or three hours in fevers. Mixed in sirup or molasses.

This powder, which has been introduced as the succedaneum of the celebrated Fever Powder of Dr. James, has lately been pronounced inert by Mr. Richard Phillips, chemist of London. Learning that it had produced no effect when largely exhibited, he determined to examine it *chemically*, and found it to consist of peroxide of antimony, (which he says is inert,) and the phosphate of lime in mixture. Dr. Paris observes, "It will be difficult for the chemist to persuade the physician he can never have derived any benefit from the exhibition of antimonial powder."

Powders of Precipitated Sulphuret of Antimony, Calomel, &c.

R. Sulphureti Antimonii præcipitati,

Hydrargyri chloridi mitis, ââ. gr. vj .

Pulveris Guaiaci, $\mathfrak{z}\text{j}$.

Misce.—Fiant chartulæ vj.—Signa.—To be taken morning and evening, with double or even triple proportion of the ingredients if necessary. In some cutaneous diseases.

MIXTURES, &c.

Mixture of Sub-carbonate of Potash, Lemon Juice, &c. vulgo, Neutral or Saline Mixture.

R. Succì limonis recentis, $\mathfrak{z}\text{iss}$.

Carbonatis Potassæ, q. s. ad saturandum,

Sacchari albi, $\mathfrak{z}\text{j}$ vel $\mathfrak{z}\text{ij}$.

Tartratis Antimonii, gr. j.

Aquæ destillatæ vel menthæ, $\mathfrak{z}\text{iiij}$.

Misce.—A table-spoonful may be taken every two hours, in Fevers.

This preparation is one of our most elegant and agreeable diaphoretics. Its powers are decidedly increased by the addition of tartar emetic. This, however, may be omitted, and the sweet spirits of nitre substituted, if preferred.

Solution of Citrate of Potash.

R. Citratis potassæ preparatæ, ℥ij.

Aquæ destillatæ, ℥iv.

Olei limonis, gtt. ij.

Sacchari albi, ℥j.

Misce.—A table-spoonful to be taken every two hours.—This preparation is a good substitute for the preceding one, and is in several respects to be preferred.

Pills of Antimonial Powder, Calomel, &c.

R. Pulveris Antimonialis, gr. x.

Opii pulveris,

Hgdrargyri chloridi mitis, āā. gr. ij.

Confectionis rosæ, q. s.

Fiant pilulæ æquales quatuor.

Signa.—Two to be taken at bed time. In acute rheumatism.

Pills of Tartar Emetic, Camphor, &c.

R. Antimonii tartarizati, gr. ij.

Pulveris Opii, gr. vj.

Camphoræ, gr. xxxvj.

Spiritus rectificati, gtt. iij.

Conservæ rosarum, q. s.

Fiat pilulæ xij.—Signa.—One to be taken every fourth hour.

Pills of Red Sulphuret of Mercury, &c.

R. Hydrargyri sulphureti rubri,

Pulveris Serpentariæ, āā ℥ss.

Syrupi simplicis, q. s.

Fiat pilulæ xij.—Signa.—Two of these may be taken three times a day.

They are alterative as well as diaphoretic, and will be found useful in herpetic, and other obstinate cutaneous complaints.

Pills of Sulphuret of Antimony, &c.—Olim Plummer's Pills.

- R. Hydrargyri chloridi mitis,
Antimonii sulphureti, āā. ʒss.
Pulveris Guaiaci, ʒj.
Balsami Copaibæ, q. s.—ut fiant pilulæ lx.

Three of these pills may be taken night and morning.

They were recommended by Professor Plummer in leprosy, secondary syphilis affecting the skin, gleet, old ulcers, &c. &c. The decoction of Guaiacum generally accompanies their use.

Anodyne Pills.

- R. Extracti Opii, gr. x.
Antimonii sulphureti præcipitati, gr. xij.
Nitratis Potassæ, gr. xxiv.
Syrupi simplicis, q. s.
Ut fiant pilulæ vj.

Signa.—One at bed-time in rheumatic affections. To diminish pain and produce diaphoresis. *M. Recamier.*

Effervescing Draught.

- R. Carbonatis Potassæ, ʒij.
Aquæ destillatæ, ʒiv.
Misce et fiat solutio.

- R. Succī limonis recentis,
Aquæ destillatæ, āā. ʒij.

Misce, et signa.—Mix two table-spoonsful of the lemonade with one of the alkaline solution, and let it be drunk while effervescing, and repeated *pro re nata*.

This preparation differs from the neutral mixture more in the mode of exhibition than in any other important particular. It is of great value where the irritability of the stomach precludes the use of the ordinary diaphoretics.

Sweet Spirits of Nitre, with Antimonial Wine, &c.

R. Tincturæ thebaicæ, gtt. xxv.

Spiritus ætheris nitrici, ʒj.

Vini Antimonii, gtt. xx.

Aquæ destillatæ, ʒss.

Fiat mistura.—To be given at a dose, and repeated *pro re nata*.

Dr. Chapman observes in his Therapeutics—"This combination very frequently acts as a diuretic in dropsy, and has more than once, within my knowledge, removed the effusion in this disease, under circumstances the least to be expected.

Sweet Spirits of Nitre, &c.

R. Spiritus ætheris nitrici, ʒj.

Vini Antimonii, ʒij.

Misce, et signa.—A tea-spoonful may be taken every hour or two hours, in barley water, during the hot stage of intermittent and other fevers.

Water of Acetated Ammonia, &c.

R. Liquoris Ammoniæ acetatis, ʒij.

Aquæ Cinnamomi, ʒj.

Vini Antimonii, ʒj.

Aquæ destillatæ, ʒij.

Fiat mistura.—Signa.—A table-spoonful every two or three hours.

Tincture of Tolu with Wine of Antimony, &c.

R. Tincturæ Tolutani, ʒv.

Pulveris gummi Arabici, ʒiij.

Sacchari albi, ʒj.

Vini Antimonii, ʒss.

Aquæ Cinnamomi, ʒij.

Fiat mistura.—Signa.—A table-spoonful three or four times a-day.

This is a stimulating diaphoretic, and will be found serviceable in pectoral diseases accompanied with an excessive mucous discharge, which appears to depend upon debility of the lungs.

Mixture with Camphor, Antimonial Wine, &c.

R. Misturæ Camphoratæ, ℥ss.

Aquæ Ammoniacæ acetatis, ℥ss.

Vini Antimonii, gtt. xx.

Tincturæ Opii, gtt. xx.

Misce, et signa.—To be taken at bed-time, or oftener, but in the latter case diminishing the quantity of laudanum.

For the prescription for *Mistura camphorata*, see Stimulants.

Mixture with Guaiacum, &c.

R. Pulveris Guaiaci, ℥ij.

—— Nitratis Potassæ, ℥iss.

Pulveris gummi Arabici, ℥j.

Tartratis Antimonii, gr. iss.

Pulveris Glycyrrhizæ, ℥j.

Aquæ destillatæ, vel Menthæ, ℥viij.

Fiat mistura.—Signa.—A table-spoonful three or four times a-day.

This preparation, if made without the extract of liquorice, is disposed to turn green, which gives it an unpleasant appearance, and alarms the patient sometimes so much as to induce him to refrain from it until the physician arrives. It is employed principally in rheumatic cases.

Sirup or Rob Anti-syphilitic of Laffeteur.

R. Radicis Smilacis sarsaparillæ,

Arundinis phragmitis, aa. ℥xxx.

Florum Boraginis officinalis, ℥viij.

Foliorum Cassiæ sennæ,

Petalarum rosæ albæ, aa. ℥ij.

Secchari albi,

Mellis, aa. ℥vj.

Boil the sarsaparilla and marsh reed grass in nine pints of water for one hour; strain off the decoction, and pour the same quantity of water on the residuum, which is to be boiled for two hours: towards the end of the boiling add the borage flower, senna, and rose leaves, and then strain off, and to both decoctions add the sugar and honey, and boil the whole to the consistence of a sirup.

The dose for men is six table-spoonsful, and for women four, taken without addition, at six in the morning.

The sarsaparilla ptisan is then to be commenced; and is made as follows:—

Boil ℥ij. of sarsaparilla in three quarts of water, to two quarts, suffer it to infuse during the night, and strain it off in the morning.

This is to be drunk by the tumblerful, and several are to be taken during the day.

The above preparation has lately been asserted, by the New York Medical Society, to be nearly identical with the noted panacea of Swaim. It is believed that corrosive sublimate is often introduced into these nostrums at the discretion of the vender. That both the French and American have frequently salivated, is well ascertained.

Decoction of Burdock.

R. Radicis Arctii lappæ consisæ, ℥iij.

Aquæ fontanæ, Oij.

Reduce the fluid by boiling to two pints, and strain. A tea-cupful may be taken for a dose, five or six times a-day.

Recommended in syphilis, scurvy, arthritic affections, &c.

Decoction of Mezereon.

R. Corticis radicis Daphnis mezerei, ℥vj.

Aquæ fontanæ, ℥vj.

Reduce by boiling to ℥iv., and near the end of the decoction, add Radicis Glycyrrhizæ glabræ, ℥j. Strain, and give from one to four pounds per diem, as the stomach will bear it. Recommended in scrofula complicated with syphilis, &c.

Decoction of Lobelia Syphilitica.

R. Radicis Lobeliæ syphiliticæ siccatae et concisæ, ℥ss.

Aquæ fontanæ, ℥xij.

Boil down to ℥viij., and strain. Commence with half a pint daily, and increase it to two pints per diem, until free purging is induced; then omit it for three or four days, and resume and continue in this way until a cure is effected.

Applicable to secondary syphilis, and those diseases depending on the excessive use of mercury.

For a variety of decoctions of similar character, see *Swediasur on Syphilis*.

Solution of Sulphate of Magnesia with Tartar Emetic.

Give of this solution a table-spoonful every hour.

For the prescription, see Cathartics.

Solution of Citrate of Ammonia, &c.

R. Succi limonis recentis, ℥j.

Carbonatis Ammoniaë, q. s. ad saturandum,

Sacchari albi, ℥j.

Aquæ destillatæ, vel Menthæ, ℥iv.

Fiat mistura.—Signa.—A table-spoonful may be taken every two or three hours, *pro re nata*.

This preparation is seldom employed in the practice of this city. It may, however, be beneficially given, where we wish a pleasant stimulating diaphoretic.

Infusion of Boneset.

R. Eupatorii perfoliati, ℥j.

Aquæ bullientis, Oj.

Infuse for thirty minutes in a covered vessel. Then decant and take a wine-glassful every hour, till perspiration is freely produced. The infusion must be taken as hot as it can be drunk, otherwise, it may produce vomiting.

Decoction of Sarsaparilla, &c.

R. Give of the Compound Decoction of Sarsaparilla, a tea-cupful four or five times a-day.

This preparation, the formula for which is to be found in almost any Pharmacopœia, has long been celebrated in the sequelæ of syphilis, and the disease produced by the improper exhibition of mercury. It is also much employed in rheumatism. The rules laid down in the Dispensatories for the preparation of the decoction should be closely observed.

Infusion of the Asclepias Decumbens, or Pleurisy Root.

R. Radicis Asclepiadis decumbentis, ℥j.

Aquæ bullientis, Oiss.

Fiat infusio.—Signa.—A tea-cupful to be taken warm, as frequently as the stomach will bear it.

Professor Chapman observes of this plant—"As a diaphoretic, I think it is distinguished by great certainty and permanency of operation, and has this estimable property, that it produces its effects without increasing much the force of the circulation, raising the temperature of the surface, or creating inquietude and restlessness."—*Therapeutics, Vol. I. p. 347.*

Decoction of Barley with Nitrate of Potash.

* R. Decocti Hordei, Oj.

Nitratis Potassæ, ʒij.

Succi limonis, ʒj vel ʒij.

Fiat potus.—To be used warm by the patient as a common drink.

Infusion of Pipsisseway.

R. Pyrolæ umbellatæ, ℥j.

Aquæ bullientis, Oj.

Fiat infusio.—A tea-cupful every two hours.

Mixture of Muriate of Baryta, &c.

R. Muriatis Barytæ,

Muriatis Ferri, aa. ʒss.

Aquæ destillatæ,

Syrupi corticis aurantii, aa. ℥j.

Fiat mistura.—Signa.—The dose may be from twenty to thirty drops every three hours. In some serofulous diseases of obstinate character.

Hufeland.

* For Decoction of Barley, see Dietetics

CLASS VI.

EXPECTORANTS.

DURING the administration of Expectorants the patient should be kept warm, and, if the complaint be violent, in bed. The dose of all Expectorants should be repeated sufficiently often to allay the irritation as much as possible; for it is obvious that the act of coughing increases that very irritation which originally gave rise to it.

POWDERS, PILLS, &c.

Powders of Myrrh, Ipecacuanha, &c.

R. Pulveris Myrrhæ, gr. xij.

———— Ipecacuanhæ, gr. vj.

———— Potassæ nitratis, ʒss.

Misce, et divide in doses quatuor. One to be taken every fourth hour.

A stimulating expectorant.—*Paris' Pharmacologia.*

Compound Powder of Ipecacuanha.

R. Pulveris Doveri, ʒss.

Divide in chartulas iij.—Signa.—One, two or three times a-day, in sugar and water.

This remedy is introduced under the head of Diaphoretics, where perhaps it most properly belongs; but it often proves of such decided advantage as an expectorant that it also deserves an insertion here.

Powders with Nitrate of Potash, Calomel, Opium, &c.

R. Nitratis Potassæ, ʒj.

Hydrargyri chloridi mitis, gr. vj.

Pulveris Opii, gr. iij.

——— Ipecacuanhæ, gr. vj.

Misce, et divide in chartulas vj.—Signa.—One every two or three hours in sirup or molasses.

This preparation closely resembles the former, with the exception that it contains *calomel*, which is often serviceable in pleuritic cases.

It may be remarked, that all powders containing calomel, or other metallic preparations, should be given in sirup or molasses, as they sink when water is employed as the vehicle.

Pills of Sulphate of Zinc and Myrrh.

R. Sulphatis Zinci, gr. x.

Pulveris Myrrhæ, ʒiss.

Confectionis rosarum, q. s.—ut fiant pilulæ xx.

This prescription is taken from the Pharmacologia of Dr. Paris, who recommends two to be taken twice a-day.

It is principally used in pertussis, in which disease a more efficient direction would be to give a pill every two hours. Dr. Paris remarks, that the principal nostrums for whooping-cough are formed either of opium or sulphate of zinc.

Pills of Conium, Ipecacuanha, &c.

R. Extracti Conii,

Pulveris Ipecacuanhæ compositi, āā. gr. x.

Misce.—Fiant pilulæ iv.—Of which two may be taken at bed-time.—

In pulmonary irritation, attended with rheumatic and other local pains.

Pills of Tartar Emetic, &c.

R. Tartratis Antimonii et Potassæ,

Opii, āā. gr. iij.

Pulveris Tragacanthæ, gr. x.

Conservæ rosarum, q. s.—ut fiant pilulæ l.—Signa.—Two to be taken night and morning. In obstinate pulmonary catarrh.

Dr. Pariset.

Pills of Squill, Ammoniac, &c.

R. Pulveris Scillæ, gr. xxx.

Ammoniaci gummi resinæ, ʒiss.

Extracti Conii, gr. xxx.

Contunde simul, et divide massam in pilulas æquales xxx.—Of which two may be taken twice a-day. In asthma and chronic catarrh.

Pills of Squill, Calomel, &c.

R. Hydrargyri chloridi mitis, gr. iij.

Scillæ recentis,

Ammoniaci gummi resinæ, āā. ʒj.

Pulveris Ipecacuanhæ compositi, ʒj.

Confectionis rosæ, q. s.—ut fiant massa et divide in pilulas xxx.—Signa.—One three or four times a-day. In troublesome cough.

Dr. Latham.

The recent squill is greatly preferred by some physicians. It is generally given in doses double the size of those of the dried article.

Pills of Sulphate of Zinc.

R. Sulphatis Zinci, ʒss.

Conservæ rosarum, q. s.—ut fiant pilulæ xx.—One of these pills may be given three times a-day.

They are recommended by some physicians in phthisis and pertussis.

Tar Pills.

R. Picis liquidæ, ʒj.

Pulveris glycyrrhizæ, q. s.

Ut fiant pilulæ xxx vel xl.—One to be taken every three or four hours.

Pills of Calomel, Squills, &c.

R. Hydrargyri chloridi mitis, gr. xxiv.
 Pulveris Scillæ, ℥ss.
 Tartratis Antimonii, gr. vj.
 Pulveris Opii, gr. xviii.
 Fiant pilulæ xxiv.—Signa.—One every night. In catarrhal affections.

Ipecacuanha Pills.

R. Pulveris Ipecacuanhæ, gr. xv.
 Saponis albi, q. s.—ut fiant pilulæ xv.—Signa.—One three times a-day.

These pills are employed in protracted catarrh, and in some cases of incipient phthisis.

Pills of Myrrh, &c.

R. Myrrhæ gummi resinæ, ℥iss.
 Scillæ exsiccata, ℥ss.
 Extracti Hyosciami, ℥ij.
 Aquæ destillatæ, q. s.—ut fiant pilulæ xxx.—Two to be taken night and morning.

These pills are recommended by Dr. Paris, in cases of phthisis and catarrh attended with great debility of the lungs.

Lozenges with Liquorice, Opium, &c.

R. Pulveris gummi Acaciæ,
 ——— Extracti Glycyrrhizæ,
 ——— Sacchari albi, āā. ℥ij.
 ——— Opii, gr. vj.
 Olei Anisi, gtt. iv.
 Misce, et adde—
 Aquæ destillatæ, q. s.—ut fiant trochisci lx.—Signa.—One of these may be dissolved in the mouth three or four times a-day.

These are the celebrated *Cough Lozenges* of Dr. Wistar.

Emetine Lozenges.

R. Emitinæ coloratæ, gr. xxxij.

Sacchari albi, ℥ij.

Divide into Lozenges, each weighing ten grains. One of these will produce more or less nausea in a child; but if this is not desirable, they can be taken in divided portions. Adults may take five or six at intervals of two hours, without the risk of vomiting.

Cubeb Lozenges.

R. Pulveris cubebæ, ℥ij.

Balsami Tolutanæ, gr. vj.

Syrupi Balsami Peruviani,

Extracti Glycyrrhizæ, āā. ℥j.

Gummi arabici, q. s.

Mix, and divide into Lozenges of ten grains each. These are called Spitta's Lozenges, and are celebrated for the relief they afford in Coryza.

MIXTURES, SIRUPS, &c.

Mixture with Extract of Liquorice, &c., vulgo, Brown Mixture.

R. Pulveris extracti Glycyrrhizæ,

—— gummi Arabici, āā. ℥ij.

Aquæ ferventis, ℥iv.

Misce.—Fiat solutio, et adde—

Spiritus ætheris nitrici, ℥ij.

Vini Antimonii, ℥ij.

Tincturæ Opii, gtt. xl vel lx.

Signa.—A table-spoonful for a dose, *pro re nata*. In catarrh.

Mixture with Sirup of Squills, &c.

R. Syrupi Scillæ, ℥ss.

Tincturæ Opii camphoratæ, ℥ij.

Vini Antimonii, ℥j.

Pulveris gummi Arabici, ℥ss.

Aquæ destillatæ, ℥ij.

Misce, et signa.—A tea-spoonful every hour or two hours.

This is more particularly designed for children.

Compound Sirup of Squills, vulgo, Cox's Hive Sirup.

R. Syrupi Scillæ compositi, ℥j.

A tea-spoonful may be given to a child of one year old every quarter, half, or one hour, according to the violence of the disease.

Strongly recommended in pertussis, by Professor Coxe.

The following appears to be a more elegant compound.

R. Infusionis Polygalæ senegæ, ℥iv.

Syrupi Ipecacuanhæ, ℥j.

Oxymellis Scillitici, ℥iij.

Tartratis Antimonii, gr. iss.

Fiat mistura.—Signa.—A tea-spoonful *pro re nata* for children of two years of age. In croup.

Infusion of Polygala.

R. Radicis Senegæ, ℥ij.

Aquæ bullientis, ℥vj.

Syrupi Tolutanæ, ℥j.

Gummi Ammoniaci, ℥ss.

Misce.—A table-spoonful is to be given every two hours. In catarrh.

Mixture of Extract of Conium, &c.

R. Extracti Conii maculati, ℥j.

Tincturæ Opii camphoratæ, ℥ss.

Syrupi Tolutani, ℥ss.

Aquæ rosarum, ℥iv.

Misce, et signa.—From half to a whole tea-spoonful may be given once in four hours to a child one year old.

Dr. Pearson, in his work on pertussis, recommends the above as a valuable formula where a narcotic is wanted.

Mixture of Almond Emulsion, &c.

R. Misturæ Amygdalæ, ℥v.

Vini Ipecacuanhæ,

Tincturæ Scillæ, āā. ℥j.

Syrupi Tolutani, ℥vj.

Misce, et signa.—Take a table-spoonful whenever the *cough* is urgent.

Almond emulsion is prepared by depriving the kernels of sweet almonds of their skins by immersing them in warm water, (called blanching.) Then beating them with some sugar to a pulp in a mortar, and adding water gradatim until an emulsion of sufficient thickness is formed. It is then to be strained.

White Linctus.

R. Amygdalarum dulcium, no. xv.

Gummi Tragacanthæ, ʒj.

Syrupi simplicis, ʒj.

Olei Amygdali communis, ʒij.

Aquæ florum Aurant: ʒij.

— destillatæ, ʒiv.

Misce.—Signa.—A table-spoonful for a dose, frequently through the day. In catarrh.

Mistura Moschi.

See Antispasmodics.

Mixture of Oil of Amber, &c.

R. Olei Succini rectificati, gtt. lxxx.

Gummi Arabici,

Sacchari albi, āā. ʒss.

Tincturæ balsami Tolutani, ʒij.

Aquæ destillatæ, ʒiv.

Fiat mistura secundum artem.—Signa.—A table-spoonful every two or three hours. In spasmodic cough.

Mixture with Carbonate of Potash, &c.

R. Carbonatis Potassæ, ʒij.

Vini Antimonii, ʒj.

Tincturæ Opii, gtt. xl.

Spiritus Lavendulæ compositi, ʒij.

Aquæ destillatæ, ʒiv.

Fiat mistura.—And let a table-spoonful be taken every hour or two hours.

Mixture with Bi-carbonate of Soda, &c.

R. Bi-carbonatis Sodæ, gr. xij.
 Vini Ipecacuanhæ, gtt. xx.
 Tincturæ Thebaicæ, gtt. iv.
 Aquæ destillatæ, ℥j.

Misce.—Signa.—Give a tea-spoonful every two or three hours.

This preparation is designed for children, and is chiefly used in pertussis.

Solution of Carbonate of Potash with Cochineal.

R. Carbonatis Potassæ, ʒj.
 Pulveris Coccinellæ, ʒss.
 Sacchari albi, ʒj.
 Aquæ destillatæ, ℥iv.

Fiat solutio.—Dose for children a tea-spoonful every two or three hours.

This is a popular remedy in pertussis.

Mixture with Tincture of Digitalis, &c.

R. Tincturæ Digitalis, ℥ss.
 ——— Opii, gtt. lx.
 Aquæ destillatæ, ℥ijss.

Misce.—A small tea-spoonful may be taken three or four times a-day.
 In hæmoptysis and in incipient phthisis.

Solution of Asafætida, &c.

R. Gummi Asafætidæ, ʒj.
 Aquæ destillatæ, ℥iv.
 Fiat solutio, et adde—
 Tincturæ Tolutani, ℥ss.
 ——— Opii, gtt. xl vel l.

A tea-spoonful to a child every two hours, and a pap-spoonful, (or more,) as frequently to an adult.

This preparation is one of great value in pertussis and all spasmodic diseases of the chest.

Tincture of Blood Root.

R. Tincturæ Sanguinariæ Canadensis, Saturatæ, quantum placet.
Direct from 30 to 50 drops twice in the day.

This native product is one of the most powerful medicinal agents. It is violently emetic in large doses. In the quantity recommended it is said to produce diaphoresis, as well as effects analogous to digitalis.

Its active principle has been separated and is called "*sanguinarina*."

Solution of Gum Ammoniac, &c.

R. Gummi Ammoniaci, ℥j.

Aquæ destillatæ, ℥iv.

Fiat solutio, et adde—

Oxymellis Scillæ, ℥ij.

Tincturæ Opii camphoratæ, ℥ss.

A table-spoonful may be given every two hours.

This is a stimulating expectorant, and is useful in cases where debility appears to be the principal disease of the lungs.

Mixture with Milk of Asafoetida and Acetate of Ammonia.

R. Gummi Asafoetidæ, ℥ss.

Liquoris Ammoniacæ acetatis,

Aquæ Menthæ, āā. ℥ij.

Fiat mistura.—A tea-spoonful to children, and a table-spoonful to adults in coughs, *pro re nata*.

This simple solution of asafoetida is frequently employed, and may, in some cases, be found to answer every purpose of a more complicated prescription.

Mixture of Copaiba and Balsam Tolu, &c.

R. Balsami Copaibæ,

——— Tolu,

Pulveris gummi Arabici, āā. ℥ss.

Acidi sulphurici aromatici, gtt. l.

Aquæ destillatæ, ℥vj.

Fiat mistura.—Dose, a table-spoonful *pro re nata*. In chronic whooping-cough. Where there is hectic excitement, digitalis may be added.

Tincture of Tolu with Digitalis, &c.

R. Tincturæ Tolutani, ℥iss.
Acidi sulphurici aromatici,
Tincturæ Digitalis,
Vini Antimonii, āā. ℥ij.
Mellis despumati, ℥iss.
Pulveris Glycyrrhizæ, ℥ss.
Aquæ destillatæ, ℥vj.

Fiat mistura, et signa.—A table-spoonful to be given, according to circumstances. *In Tussis senilis*.

Mixture with Balsam Peru, &c.

R. Balsami Peruviani, ℥ss.
Mucilaginis Acaciæ, ℥ij,
Aquæ Cinnamomi,
—— destillatæ, āā. ℥ss.

Misce.—Signa.—To be taken at once, and repeated three or four times a-day. In cases of debilitated lungs.

The different *Balsams* have been almost proscribed from the class of expectorants, on account of their stimulating qualities. Balsam Copaiva was once employed, and has again been proposed by a modern writer, in diseases of the trachea.

The author has prescribed it with great advantage in that peculiar hoarseness which sometimes results from a violent attack of measles.

Tar Water.

R. Picis liquidæ, Oij.
Aquæ cong. j.

Mix them, and stir them together with a wooden rod for an hour, then let the tar subside, strain the water and keep it in well-corked bottles.—A pint, or even two pints, may be taken daily.

Mixture of Gum Ammoniac and Nitric Acid, called Nitric Lac Ammoniac.

R. Gummi Ammoniaci, ℥ijss.

Acidi nitrici, ℥ij.

Aquæ destillatæ, ℥viiij.

Pour the nitric acid and water together, then add the compound gradually to the gum, incorporating them well in a composition mortar. A table-spoonful may be taken three or four times a-day.

This preparation is strongly recommended in protracted catarrh, and some stages of phthisis pulmonalis, by Dr. J. Stuart, author of a treatise on *Colds*.

Mixture of Spermaceti, &c.

R. Spermat. Ceti, ℥ij.

Sacchari albi, ℥iij.

Tincturæ Opii camphoratæ, ℥ss.

Aquæ destillatæ, ℥viiij.

Rub the spermaceti and sugar together with the yelk of an egg, then gradually mix the water.—A table-spoonful for a dose, four or five times a-day. In Catarrh.

Sirup of Tolu with Belladonna, &c.

R. Extracti Belladonnæ, gr. xij.

Vini Ipecacuanhæ, ℥j.

Syrupi Senegæ, ℥ss.

Misce et signa.—To take a tea-spoonful three or four times a-day, freely using gum water acidulated with lemon juice. In Coughs.

Mucilage of Gum Arabic.

R. Gummi Acaciæ optimi, ℥ss.

Aquæ ferventis, ℥iv.

Fiat solutio.

Used in most pectoral diseases.

Infusion of Flaxseed, &c.

R. *Seminum Lini*, ʒj.
Radicis Glycyrrhizæ, ʒss.
Aquæ bullientis, Oij.

Macerate for two or three hours near the fire in a covered vessel: strain, and add lemon juice sufficient to make it agreeable.

It may be given as a common drink in catarrh.

Decoction of Iceland Moss.

R. *Lichenis Islandici*, ʒj.
Aquæ fontanæ, Oiss.
 Boil to one pint, and strain.

This remedy once acquired great celebrity in coughs; but was overrated, and lost its reputation. It contains a bitter quality, which often recommends it in dyspepsia and in phthisis. From this it may be freed if necessary, by macerating it before boiling, and throwing the first liquor away. To be taken *ad libitum* by the patient.

Prussic or Hydrocyanic Acid.

As prussic acid rises to the top of an aqueous mixture, and thereby renders the dose very uncertain, we prefer directing the patient to drop the dose each time into a solution of gum Arabic or simple sirup. It should be kept covered from the light, and not allowed to be accessible to those unacquainted with its power, as even a strong inhalation will sometimes prostrate an adult. The physician should also know after what process the acid is made that he prescribes. That of Scheele, who discovered it, affords an acid containing some water, and this may be unequal at different times. The process of Gay Lussac, yields a fluid free from admixture, and of course much more energetic than the former. Majendie directs the Prussic acid of Gay Lussac to be diluted with 8.5 times its weight of water, and this he denominates *Medicinal Prussic Acid*. This

remedy is doubtless very efficacious in some diseases, but it is to be regretted that it is liable to such inequality of strength. We are told by Majendie, that, left to itself in close vessels, it decomposes sometimes in *less than an hour*, and can scarcely be kept over fifteen days. This is to be understood only of the acid of Gay Lussac.

The Prussic acid of Brande, as kept in our shops, may be commenced in doses of a drop three times a-day in mucilago acaciæ, or sugar and water, and augmented one drop daily, until its action on the system is fully displayed.

Majendie directs the following, among other formulæ, for the exhibition of his medicinal Prussic acid.

Pectoral Mixture.

R. Acidi Hydrocyanici, ʒj.

Aquæ destillatæ, ℥j.

Sacchari albi, ʒiiss.

Fiat mistura.—Signa.—A pap-spoonful may be taken morning and evening, and gradually increased, (if necessary,) to six or eight spoonsful in the twenty-four hours—shaking the vial each time.

Prussic acid is also used externally, in the proportion of ʒij. to Oj. of water, or other fluids. In cancerous ulcers; dartre; and, injected into the vagina, in cancer of the uterus.—*See Majendie's Formulary, Edit. 5th, p. 66.*

Dr. Morton's mode of administering Prussic acid is as follows:—

“ In prescribing it I have mostly confined myself to a single prescription, which is a slight modification of that recommended by Dr. Granville of London.—Let half an ounce of gum Arabic be dissolved in seven ounces and a half of water, to which add half an ounce of sirup of tolu and twelve drops of Prussic acid prepared according to the formula of the London Pharmacopœia. Of this mixture, a table-spoonful may be

taken every three hours,—a medium dose which I prefer in commencing the use of so powerful a medicine.”—*Illustrations of Pulmonary Consumption*, p. 132.

Cyanide of Potassium.

By dissolving this substance in eight times its weight of distilled water, it becomes the hydrocyanate of potassa, and may be employed as a remedy in the same doses as Prussic acid. It is said to exercise the same power over the system as the latter, with the advantage of being a more uniform and permanent medicine.

Mixture of Cyanide of Potassium.

R. Potassii hydrocyanici medicatæ, ℥j.

Aquæ destillatæ, Oj.

Sacchari purificati, ℥iss.

Fiat solutio.—Dose—a table-spoonful night and morning.

Majendie.

Cyanide of Potassium.

R. Potassii cyanidi, gr. ss.

Aquæ lactucæ, ℥ij.

Syrup althææ, ℥j.

Fiat mistura.—Dose—a table-spoonful every two hours.—*Majendie.*

Pills of Cyanide of Potassium.

R. Potassii cyanidi,

Amyli, āā. gr. iv.

Syrupi simplicis, q. s.

Ut fiant pilulæ, viij. Dose—a pill night and morning.

The preceding preparations are chiefly administered in pectoral affections, as Asthma, Bronchitis, Phthisis, &c.

Mixture of Gum Ammoniac, &c.

R. Misturæ Ammoniaci, ℥ij.

Tincturæ Castorei, ℥ss.

Syrupi Tolutani, ℥ss.

Tincturæ Opii, gtt. v.

Aquæ Cinnamomi, ℥j.

Fiat mistura.—Signa.—A small table-spoonful every three or four hours in troublesome cough.

Dr. Meigs.

Mixture of Morphia and Tolu.

R. Oxymellis Scillæ, ℥iss.

Syrupi Tolutani, ℥ss.

Acetatis Morphiæ, gr. ij.

Misce.—Signa.—A tea-spoonful may be taken every two or three hours, *pro re nata*. Useful in Phthisis pulmonalis, and other diseases attended with troublesome cough.

Dr. Meigs.

Tolu with Morphia and Antimony.

R. Mucilaginis gummi Arabici, Oj.

Syrupi Tolutani, ℥ij.

Morphiæ sulphatis, gr. iss vel ij.

Antimonii tartarizati, gr. ij.

Misce, et signa.—A table-spoonful to be given every two hours.

Morton on Consumption, page 123.

Decoction of Seneka Root.

R. Radicis Polygalæ senegæ contusæ, ℥j.

—— Glycyrrhizæ, ℥ss.

Aquæ destillatæ, Oiss.

Boil down to a pint, and strain.

This decoction is serviceable in pectoral diseases. In croup, the author has seen it productive of great benefit.

INHALATION.

Inhalation is a means of applying remedies directly to the lungs.

The best instrument for this purpose is the *Inhaler* of Mudge. As a substitute however, the common tea-pot may be used, or a funnel inverted over a bowl.

The substances usually selected for inhalation are—

The steam of warm water:

The steam of vinegar, produced by pouring it into boiling water.

Also, Balsam Tolu, as follows:—

R. Balsami Tolutani, $\bar{3}$ j.

Aquæ bullientis, Oj.

Misce.—To be inhaled in diseases of the lungs.

Ether is also used alone, or impregnated. When saturated with cicuta, and inhaled from a wine-glass, it is said to be serviceable in protracted catarrh, pneumonia, consumption, &c. Dr. Morton advises Dr. Pearson's formula, which consists in keeping a drachm of powdered cicuta in an ounce of ether for a week. A tea-spoonful or two of this impregnated fluid may be inhaled at a time, and repeated thrice per diem.—*Morton on Consumption, page 136.*

Stramonium is an article recommended particularly in asthma and consumption. The root is cut, dried, and smoked from a common pipe.

Fumigations of *tar* and *turpentine* are also recommended from high authority. The former is said to have cured consumption. It usually is applied by placing the tar on a dish over coals, and suffering it slowly to evaporate:—but a far neater and more effectual mode is that practised by Sir Alexander Crichton, viz: “an ounce of potash is added to every pound of tar, in order that the latter may be deprived of its pyroligneous acid. The two ingredients being well mixed, should be first boiled for a few minutes in the open air, in order to disengage any impurities, and then be kept at a simmer in the room of the patient. This is effected by putting the mixture in an iron vessel, and placing the latter over a spirit lamp, or some

analogous contrivance. In this way, not only a chamber, but an entire house, is speedily pervaded by a most agreeable vapour, which, although it may at first excite some disposition to cough, both in healthy and sick persons, very soon, in a majority of cases, allays this symptom, and with it a great proportion of the patient's distress."—*Morton, ut supra.*

The fumes of *resin* are frequently employed by Dr. Parrish, of this city; he directs it to be put on a hot shovel, or coals, and allows the patient to inhale it gradually until it is ascertained whether it can be borne. To some persons it is very agreeable, while the lungs of others cannot tolerate it. He has found it beneficial in debility of the lungs, whether idiopathic or the result of disease; and also advises it in pertussis and phthisis.

In some cases of croup, the inhalation of ether or Hoffman's anodyne, will often procure the most prompt relief, and gain time for the application of general remedies.

Chlorine gas has also been much used by inhalation, not pure, however, but mixed or diluted with atmospheric air. For this purpose a little acid of any kind may be dropped gently on a mixture of chloride of lime, by which means the chlorine is gradually disengaged, and be inhaled as it rises; or, the air of a room may be charged with it in like manner. It has been much commended in Bronchitis and the kindred affections, but is getting into disuse from its manifest want of efficacy.*

* *Fumigation of Chlorine.*

℞. Muriatis Sodæ, ℥ij.

Oxidi Manganesii, ʒv.

Aquæ, ℥ij.

Acidi Sulphurici, ℥ij.

Mix the salt and manganese, and add at proper intervals the acid and water, mingled. When used in the wards of hospitals or other infected apartments, the patients should be previously removed. For the method of preparing the different chlorides used as disinfecting agents, see *Journal of the Philadelphia College of Pharmacy, Vol. I. No. 4.*

The following is Dr. Scuddamore's recipe for Iodine Inhalation :

R. Iodinæ,

Iodidi potassii, āā gr. vj.

Aquæ destillatæ, ℥v ℥vj.

Alcohol, ℥j.

Mix, and use for inhalation.

“ I usually commence with a drachm of this mixture, proceeding gradually to the extent of half an ounce, (rarely more,) putting two-thirds the dose for the first half of the time (10, 15, or 20 minutes,) and the other third for the remainder, always adding thirty minims of a saturated tincture of conium, with an increase if the cough be very irritable. Occasionally I add some saturated tincture of ipecacuanha; and when the respiration is spasmodically affected, some ætherial tincture of the lobelia inflata.”—*Scuddamore*.

CLASS VII.

EMMENAGOGUES.

EMMENAGOUGES are those medicines which promote the menstrual discharge, or restore it when entirely suppressed. Although by some authors the specific action of these medicines on the uterus has been questioned, if not denied, yet by others, whose authority is equally to be trusted, it has been fully acknowledged.

It must, however, be observed, that amenorrhœa occurs under opposite states of the system; and where general *debility* appears to be the cause of the local disease, it is advisable to accompany the emmenagogue with an active course of tonics. While, on the contrary, if the obstruction be the effect of a *plethoric habit*, bleeding, cathartics, &c., are essential preliminaries to the exhibition and success of a specific remedy.

POWDERS.

Powder of Hiera Picra.

R. Pulveris Hieræ picræ, ℥j.

Divide in chartulas vj.—One three times a-day.

This article is placed under the head of cathartics, but it is a remedy so generally employed as an emmenagogue, that it appears necessary also to insert it here.

Powder of Madder.

R. Pulveris Rubiæ tinctori, ℥ss.

Signa.—A small tea-spoonful to be taken morning, noon and night, in sugar and water.

It should be remembered that there are three or four qualities of madder, which circumstance may account for the frequent failure of this remedy. Dr. Dewees frequently employs it.

Powder with Savin, &c.

R. Sabinæ exsiccatae,

Pulveris Zingiberis, āā. ℥j.

—— Potassæ sulphatis, ℥ij.

Fiant pulveres vj.—Signa.—One to be taken twice a-day.

Powder with Savin and Cantharides.

R. Pulveris Sabinæ, ℥ij.

—— Cantharidum, gr. ij.

Misce et divide in pulveres iv.—Signa.—One to be taken every night, on going to bed.

Powder of Spurred Rye.

R. Pulveris Secalis cornuti, ℥ij.

Divide in chartulas iv.—Of which one may be taken morning and night, in sugar and water.

Drs. Chapman and Dorsey have placed this remedy under the head of Emmenagogues, though we are not aware that it has ever been successfully employed as such.

Pills of Myrrh, Sulphate of Iron, &c.

R. Gummi Myrrhæ,

Sulphatis Ferri, āā. ℥ij.

Carbonatis Potassæ,

Saponis purificati, āā. ℥ss.

Fiat massa, in pilulas xl. dividenda.—Of which two may be taken morning, noon and night. In that suppression of the catamenia which is incident to phthisis pulmonalis.

Hooper's Pills.

R. Sulphatis Ferri, ℥j.

Pulveris Sennæ,

———— Jalapæ,

———— Super-tartratis Potassæ; āā. ℥ss.

———— Zingiberis, gr. xij.

Syrupi simplicis, q. s.—ut fiat massa, in pilulas xxv. dividenda.

The patient is directed to take three twice in the day, followed by twenty drops of the tinctura ferri muriatis in a glass of some bitter infusion.

This compound is sold by the apothecaries as a patent medicine.

Pills of Aloes, Sulphate of Iron, &c.

R. Sulphatis Ferri exsiccati, ℥j.

Pulveris Aloes, ℥ij.

———— Caryophylli, gr. v.

Terebinthinæ venetæ, q. s.—ut fiant pilulæ xx.—Signa.—One to be taken three times a-day.

DECOCTIONS, TINCTURES, &c.

Decoction of Seneka.

R. Radicis Senegæ contusæ, ℥j.

Corticis Aurantiarum, ij.

Aquæ destillatæ, Oj.

Let the seneka simmer over a slow fire until the fluid be reduced one-third; add the orange-peel, cover until cool, and then strain.

Dr. Chapman observes—"My rule in the administration of the medicine in these cases, (that is, of amenorrhœa,) is to give about four ounces of the decoction, more or less, during the day, according to the circumstances of the case. But at the time when the menstrual effort is expected to be made, and until the discharge is actually induced, I push the dose as far as the stomach will allow, having given as much as two ounces every hour."—*Therapeutics*, Vol. II. p. 38.

Infusion of Ergot.

R. Secalis Cornuti contusi, ℥ij.

Aquæ bullientis, ℥vj.

Fiat infusio.—To be given by spoonsful, in protracted parturition occasioned by deficiency of contractile power in the uterus.

For the same purpose the powder may also be given in doses of ℥j., as well as to assist in arresting uterine hemorrhage, and to cause the expulsion of the placenta.

Decoction of Ergot.

R. Secalis cornuti, ℥j.

Aquæ, Oiiiss.

Boil gradually down to two pints, and give a table-spoonful every fifteen minutes. Used in the same circumstances as the infusion.

Tincture of Ergot.

R. Secalis cornuti, ℥iiss.

Spiritus tenuioris, Oj.

Macerate for fourteen days, and strain. This is the formula used in Guy's Hospital. Dose twenty drops to a dram.

Sirup of Ergot.

R. Secalis cornuti, gr. xx.

Extracti Opii, gr. $\frac{3}{4}$.

Syrupi simplicis, ℥viij.

M.—Dose an ounce (two table-spoonsful) *pro re nata*, in engorgement of the uterus.

Lisfranc.

Decoction of Madder.

R. Pulveris Rubiæ tinctori, ℥j.

Aquæ bullientis, Oj.

Simmer for fifteen minutes—and add—

Caryophylli contusi, ℥j.

Signa.—When cool, strain and give a wine-glassful every three hours, a short time before the expected return of the menstrual discharge.—

See Decoction on Females, p. 74.

Mixture of Camphor, &c.

R. Camphoræ, ℥j.

Spiritus vini rectificati, q. s.—ut fiat pulvis: dein adde—

Pulveris gummi Arabici, ℥j.

Sacchari albi, q. s.

Aquæ Cinnamomi, ℥j.

Fiat mistura.

This mixture would more properly appear under the head of Narcotics, but as Dr. Dewees considers it particularly efficacious in dysmenorrhœa, we have given it a place with Emmenagogues. He directs “one-half the above draught to be given the instant pain is experienced, and if not relieved in an hour or two, the remainder to be given.”—*Syst. Med. p. 158.*

Tincture of Hellebore.

R. Tincturæ Hellebori nigri, ℥j.

Signa.—A tea-spoonful three times a-day, increasing the quantity as the stomach will admit of it.

Tincture of Hellebore, Myrrh, &c.

R. Tincturæ Hellebori nigri, ℥ss.

—— Myrrhæ, ℥j.

—— Cantharidum, ℥ij.

Fiat mistura.—The patient may take thirty drops, three times a-day, in a little sugar and water.

Tincture of Savin, &c.

R. Tincturæ Sabinæ compositæ, ℥j.

—— Hellebori nigri, ℥ss.

—— Castorei, ℥ij.

Fiat mistura.—Signa.—To take forty drops three times a-day, in any vehicle.

Tincture of Guaiacum, (Dewees.)

R. Pulveris Guaiaci, ℥iv.

Carbonatis Sodæ vel Potassæ, ℥iss.

Pulveris Pimentæ, ℥j.

Alcoholis diluti, ℥j.

Misce.—“Digest for a few days.—The volatile spirit of ammonia to be added *pro re nata*, in the proportion of a drachm or two, to every four ounces of tincture.”

The above formula is different from that adopted in the Pharmacopœias, for which reason we have inserted it here. Dr. Dewees informs us, that after nearly thirty-five years' experience with this tincture, he has never known it to fail in restoring the *suppressed catamenia*, where the suppression was an *idiopathic disease*. He gives it morning, noon and night, in doses of a tea-spoonful in a little sweetened milk; and sometimes in a little Madeira, Sherry, or Teneriffe wine. It is occasionally necessary to continue the medicine for some weeks, on which occasion, the dose must be judiciously augmented.—*Syst. Med. p. 149.*

Volatile Tincture of Guaiacum, &c.

R. Tincturæ Guaiaci volatilæ, ℥j.

Balsami Copaibæ, ℥ss.

Misce.—Signa.—A tea-spoonful two or three times a-day.

In cases of amenorrhœa connected with general debility.

Tincture of Iodine.

R. Iodini optimi, ℥ij.

Spiritus vini rectificati, ℥j.

Spiritus Lavendulæ compositi, ℥ij.

Fiat tinctura.—Signa.—From ten to twenty drops twice a-day, gradually increasing it to forty drops.

The above we are informed by M. Durand, apothecary of this place, is a saturated tincture. Dr. Bell has used it beneficially in amenorrhœa, gonorrhœa, leucorrhœa, &c.

From its great activity, it requires to be cautiously administered.

Solution of Iodide of Potassa.

R. Potassii Iodidi, ℥j.

Aquæ destillatæ, ℥j.

Fiat solutio.—Signa.—Thirty drops of this solution is directed by Dr. Bell in the above cases.—*See N. A. M. and S. Journal, Oct. 1828.*

Injection of Aqua Ammonix, &c.

R. Aquæ Ammonix, gtt. xx.

Lactis recentis, ℥i.

Misce.—Signa.—This quantity to be thrown into the vagina three or four times a-day in cases of amenorrhœa.

Tincture of Cantharides, (Dewees.)

This tincture, as prepared from the formula of Dr. Dewees, differs from that kept in the shops in strength only. Ten drops may be taken three times a-day, and gradually increased; and, if necessary, until strangury is induced.

Mixture of Muriate of Iron, &c.

R. Tincturæ Muriatis Ferri,

———— Aloes compositæ, āā. ℥ss.

———— Castorei, ℥ij.

Fiat mistura.—Signa.—Take a tea-spoonful three times a-day in a wine-glassful of camomile tea.

Tincture of Hiera Picra.

R. Tincturæ Hieræ Picræ, ℥j.

Signa.—A tea-spoonful may be taken three times a-day in sugar and water.

Compound Tincture of Aloes.

R. Tincturæ Aloes et Myrrhæ, ℥j.

Dose.—A tea-spoonful *pro re nata*.

CLASS VIII.

ANTHELMINTICS.

ANTHELMINTICS are medicines employed to destroy and bring away worms from the alimentary canal. They are also termed Vermifuges.

The only unequivocal proof of the existence of worms in the alimentary canal is the ejection of some of them *per anum*, or, as sometimes happens, by the mouth. When, in such cases vermifuges are resorted to, they must be persevered in for several days, with the occasional intervention of an active cathartic.

POWDERS.

Calomel with Gamboge.

R. Hydrargyri Chloridi mitis, gr. v.

Pulveris Gambogiæ, gr. vj vel x.

Misce.—For an adult.—To be taken in sirup early in the morning.

Gamboge is sometimes given in doses of fifteen or twenty grains for the tape-worm, and is highly spoken of in such cases.

Calomel with Pink-Root.

R. Hydrargyri chloride mitis, gr. iv.

Pulveris Spigeliæ, gr. x.

Misce.—Signa.—To be taken two mornings in succession, and on the afternoon of the second day, followed by a dose of castor oil or senna and manna.—For children upwards of four years old.

Pink-Root with Savin. &c.

R. Pulveris Spigeliæ,
 ——— Sennæ, aa. ℥ij.
 ——— Sabinæ, gr. xij.

Fiant chartulæ vj.—Signa.—One every morning till three are taken, then give a dose of castor oil; if the effect is not produced, the remainder may be exhibited after the same manner.

Common Salt with Cochineal.

R. Sodii Chloridi, ℥iij.
 Coccinellæ pulverizatæ, gr. xv.

Fiant doses vj.—Signa.—One every morning, following the last dose with some purgative medicine.

Dr. Rush speaks in high terms of this remedy.

Carbonate of Iron.

R. Carbonatis Ferri, ℥j.

Divide in chartulas iij.—One powder to be taken before breakfast.

Of this remedy, also, the late Professor Rush had a very favourable opinion. He gave it in doses of from 5 grs. to ʒss. to children of from one year old to ten, and sometimes in half ounce doses, with safety and success. It is chiefly used in tape-worm.—*See Observ. and Inquiries, p. 243.*

Powder of Male Fern.

Give of the powder of the Male Fern Root from one to three drachms, in a large cupful of any distilled water, early in the morning. If the stomach should reject it, the powder must be repeated again, as soon as the sickness has gone off, in the same quantity. When it has been upon the stomach about two hours, a strong cathartic bolus made of calomel and gamboge, (eight or ten grs. of the last and five or six grs. of the first,) must be taken, and worked off with a draught of green tea every now and then: and its operation may be still farther promoted, if necessary, by half an ounce, or ʒvj. of salts, dissolved in warm water. Till the worm comes away, which

generally happens on the same day, the patient is to take nothing but broth. If, however, the worm should not be discharged, either wholly or in part, the powder is to be repeated with the same regimen, the next day. In the tape-worm.—*Nouffer*.

It has been suggested, and with some probability, that the purge directed to follow this remedy is that on which the greatest reliance is to be placed.

Tin Electuary.

R. Pulveris Stanni, ʒvj vel ʒj.

* Sacchari liquidi, ʒiv.

Misce.—Two or three table-spoonsful may be given early in the morning; to be repeated until the whole be taken, and then followed by an active cathartic.

The efficacy of this remedy is said to be increased by combination with sulphur, by which means sulphuretted hydrogen would be generated, and the worm destroyed by this offensive gas. Larger doses than those mentioned are sometimes prescribed.

Cowhage Electuary.

R. Spicularum Dolichi, ʒj.

Mellis liquidi, q. s.—ut fiat electuarium.—Dose, a tea-spoonful to a child two or three years old; to be given before breakfast in the mornings of three successive days, and then followed by a brisk cathartic. In *Lumbrici*.

Vermifuge Bolus.

R. Hydrargyri Chloridi mitis, gr. ij.

Seminum Artemisiæ santonicæ, gr. viij.

Camphoræ, gr. vj.

Syrupi, q. s.—ut fiat bolus.

To be repeated *pro re nata*—for children of four or six years of age.

Ratier's Formulary.

* Molasses. Any sirup will, of course, answer the same purpose.

INFUSIONS, DECOCTIONS, &c.

Infusion of Pink-Root, &c., vulgo, Worm-Tea.

R. Radicis Spigeliæ Marilandicæ, ʒss.

Sennæ foliorum, ʒij.

Mannæ optimæ, ʒj,

Seminum Fœniculi, ʒij vel ʒss.

Aquæ bullientis, Oj.

Fiat infusio.—Signa.—A wine-glassful may be given to a child of two years old, three or four times a-day.

Pills of Venice Turpentine, &c.

R. Terebinthinæ venetæ, ʒj.

Extracti Hyosciami, gr. iv.

Hydrargyri Chloridi mitis, gr. viij.

Fiat massa, et divide in pilulas singulas gr. ij.

Of these four are to be taken every three hours, and during the entire period, the patient is to take only weak broth, tea, or coffee.

They are strongly recommended by some of the German practitioners in tape-worm.

Infusion of Pink-Root.

R. Radicis Spigelia, ʒj.

Aquæ bullientis, Oj.

Fiat infusio.—Of which a child may take from one to two table-spoonsful at a dose; an adult from ʒj to ʒiv.

The spigelia sometimes affects children violently; hence the necessity of caution in its use.

Decoction of the Geoffrea Bark.

R. Corticis Geoffrææ, ʒj.

Aquæ Oij.

To be boiled over a slow fire until it assumes the colour of amber or Madeira wine.—Dose for adults four table-spoonsful; for children two or three years old, one table-spoonful; and so in proportion to

the age and strength of the patient.—The dose of the extract of this medicine is stated by Dr. Chapman to be three grains, and of the powder, ʒss.

We are not aware that the remedy has been much employed in this city. Cold water is to be avoided while using this decoction; if, however, any unpleasant consequences follow its exhibition, and particularly where an overdose has been given, our best resource is *Oleum Ricini*, together with tepid water acidulated with lemon juice, or some other vegetable acid.

Mixture of Worm-seed Oil.

R. Olei Chenopodii, ʒj.

Sacchari albi,

Gummi Arabici, āā. ʒiss.

Misce, secundum artem; tunc adde—

Aquæ Menthæ sativæ, ʒijss.

A tea-spoonful of this preparation may be given four times a-day for two days in succession, and then purged off with castor oil or salts.—

For children of three to eight years old.

This is a vermifuge of considerable reputation in this country. It is generally used in *Lumbrici* and *Ascarides*; though we have heard of its having been successfully given for *Tenia*, but in much larger doses.

Spirit of Turpentine.

Is recommended for *Tape-worm*, in doses of ʒss to ʒij. daily, taken in honey.

Dr. Chapman speaks highly of this remedy; and the safety of such doses seems to be now fully established.

Olive Oil and Ammonia.

R. Olei Olivarum, ʒviij.

Spiritus Ammoniaë aromatici, ʒij.

Dose, three spoonful night and morning.

This remedy is highly spoken of by some authors, and may be given in much larger quantities; if it should not purge, it will be necessary to follow it with some aloetic or other cathartic preparation.

Decoction of the Bark of Pomegranate.

R. Corticis radicis Granati, ʒij.

Aquæ destillatæ, Oiss.

To be boiled away one-half, and given in doses of an ounce several times in the day.

“The efficacy of the bark of the root of the Pomegranate,” says Dr. Ainslie, (Mat. Med. of Hindoostan,) “as a remedy for the tape-worm, is now well established in India.”—*Paris*, vol. ii. p. 173.

Aloetic Enema.

R. Gummi Aloes, ʒij.

Lactis recentis, Oj.

Fiat enema.—The half of this may be given to a child eight or ten years old. In Ascarides.

Spirits of Turpentine with Hartshorn.

R. Olei Cornu Cervi fœtidi, ʒss.

— Terebinthinæ, ʒiiss.

Misce, et signa.—A tea-spoonful to be given every morning. In tænia. *M. Chabert.*

Spirit of Turpentine, Ether, &c.

R. Olei Terebinthinæ, ʒss.

Ætheris sulphurici, ʒij.

Pulveris gummi Arabici, ʒss.

Aquæ Chamæmelæ destillatæ, ʒxvj.

Misce, et signa.—Two table-spoonsful to be taken morning and evening. In tænia.

Dr. Frank.

Enema of Camphor, &c.

R. Camphoræ, ʒj.

Olei olivarum, ʒij.

Misce pro enemate.

This preparation is particularly adapted to *Ascarides*, and should be freely thrown up the rectum for three or four successive nights. No remedy is more effectual than this in relieving the intolerable itching caused by those worms.

Enemata are the most certain of all remedies in *Ascarides*; they may be made of various substances, such as solution of asafœtida, of common salt, hepar sulphuris, pure olive oil, soap, &c. &c.

CLASS IX.

EPISPASTICS.

“BY the ancients, all the external applications which redden or inflame the skin, were called *Epispastice*, and these were designated according to the several degrees of effect:—the slightest, entitled *phænigmoi*; the next, *sinapismi*; the more active, *vesicatorii*; and the strongest, *caustici*.”

We shall, however, as heretofore, follow the arrangement adopted by Dr. Chapman, and divide them into Rubefacients, Blisters, and Caustics; and with him too, we shall first speak of *Blisters*.

As a general rule, blisters should remain on the patient ten or twelve hours, in order to ensure their full effect. There are individuals, however, whose skins are so delicate, that a shorter application will answer every purpose; in children, it is seldom, and perhaps never, necessary to leave them on longer than four or six hours. When they are applied to the scalp, twenty-four hours is almost universally required; and Dr. Chapman directs the hair to be shaved, if possible, some hours before the application is made.

The usual dressing, when a blister is taken off, is basilicon ointment, or *ceratum simplex*, according to the urgency of the case, or the irritability of the system. In order to keep it open, the weak epispastic, or savin ointment, will be found sufficient.

In cases of acute disease, these remedies ought never to precede those means which have a tendency to reduce the system. It is generally admitted, that otherwise employed, they do harm, though there are individuals of acknowledged experience who say, "they can bring a patient to the blistering point at pleasure;" and in pleurisy, and some other acute diseases, apply them soon after a free bleeding. The blister should be placed as near the affected part as possible. In fevers they are usually put on the inside of the forearms or legs, back of the neck, or between the shoulders.

Where the system is very prostrate, they should be applied to the inside of the thighs, or the last two places mentioned: for when the circulation is very languid in the extremities, they seldom draw well, and are even liable to produce gangrene by exhausting the little vitality that remains.

Where, from peculiarity of constitution, *strangury* is produced by them, we must take them off after a few hours, and bathe the part with sweet oil or some such mild application—if this prove insufficient, resort must be had to diluent drinks, such as flaxseed tea or barley water, with the sweet spirits of nitre; camphor or parsley tea, or bladders half filled with warm water and applied to the pubes. An opium suppository or injection will be found very advantageous. A poultice of ground flaxseed or starch is often very serviceable. Prevention, however, is better than cure; and in irritable constitutions, or in persons disposed to strangury, the blistering ointment, before being spread, should be rubbed up with three or four grains of opium and three or four times as much camphor, before it is spread. This preventive seldom fails. Blisters are rendered much less irritating by covering them with a fine gauze, which, if it be pressed in close contact with the plaster, will not prevent or materially retard its operation.

It is said by some writers, that if a blister be sprinkled with powdered camphor, previous to its application, strangury will never result: this precaution cannot

always be relied on, though it will be well to give it a trial where strangury is anticipated.

Epispastic Ointment.

R. Ceræ flavæ,

Resinæ,

Axungię Porcinæ, āā. ʒvj.

Melt over a slow fire, and when at the temperature of 98° add slowly—

Pulveris Cantharidum, ʒvijss.

Misce, et fiat unguentum.

If the cantharides be united with the melted plaster before it is partially cooled, they are liable to be injured by the heat.

In employing this ointment, the margin of the blister should be covered with adhesive plaster, in order to confine it to its place, and prevent its creating unnecessary irritation. Another mode is to draw straps across it diagonally, which plan we generally prefer; it prevents the blister from rising in the middle, and thereby partially defeating its object.

It is mentioned in Paris' Pharmacologia, that the flies may be deprived of their power of producing strangury by first boiling them in water.

RUBEFACIENTS.

The substances called Rubefacients not unfrequently vesicate on very delicate skins. The term, therefore, is relative. They are employed for the same purpose as Blisters, and are often good substitutes for them.

Plaster of Mercurial Ointment, Camphor, &c.

R. Unguenti Hydrargyri, ʒj.

Olei Terebinthinæ,

Camphoræ, āā. ʒij.

Cerati simplicis, ʒj.

Fiat unguentum secundum artem. Used as a rubefacient over the hypochondriac region, when blisters are precluded.

Tartar Emetic Plaster.

This may be made by sprinkling the powdered Tartrate of Antimony on the surface of a Burgundy pitch or adhesive plaster, or on one made by spreading Basilicon, or simple ointment, on leather or linen. In either way the margin should be left clean.

This remedy requires a longer time to produce its effect than the ordinary blister; the irritation is of a peculiar and very painful kind.

Tartar Emetic Ointment.

R. Tartratis Antimonii, ʒj.

Axungiae Porcinæ, ʒj.

Misce secundum artem.

This ointment, when applied to the skin, is a slow but certain means of exciting an eruption. We have given the formula of Dr. Hannay, of Glasgow, in Scotland. Dr. Jenner recommended a much larger proportion of the antimony.

Tartar Emetic and Corrosive Sublimate.

R. Tartratis Antimonii, ʒj.

Chloridi Hydrargyri, gr. v.

Aquæ destillatæ, ʒj.

Spiritus Lavandulæ compositi, ʒj.

Solve sales in aqua, dein adde spiritum.

This solution is applied by wetting the fingers with it, and rubbing it on the part. The rubbing should be continued for five or ten minutes, and after some hours a pustular eruption will appear.—Dr. Hannay's essay, "*De Tartratis Antimonii Usu Externo*," p. 26.

It has been extolled in pertussis, dyspnœa, cough, pains in the chest, &c., and in the removal of incipient tumours.

It has been ascertained by the illustrious Jenner and others. that Tartar Emetic employed for an *undue*

length of time externally, will produce violent effects upon the system.

We find in a note in Beck's Medical Jurisprudence, vol. ii. p. 283, that Dr. Francis mentions two cases where Tartar Emetic Ointment, excessively used, produced effects similar to those at times arising from the outward application of arsenic. Majendie and Brodie ascertained by experiments upon animals, that inserted into wounds its effects resemble those produced by internal exhibition.

Granville's Ammoniated Counter-irritant, or Antidynous Lotions.

These preparations act with great celerity, and are now in common use. They are formed of *three ingredients*, mixed in certain proportions, according to the urgency of the case.

1. The first ingredient is the *strongest liquor of ammonia* (liquor ammoniæ fortissimus) which is made by saturating a given quantity of distilled water, contained in a glass receiver surrounded by ice, with ammoniacal gas, obtained in the usual way from a mixture of equal parts of hydrochlorate of ammonia and recently slacked lime, both reduced to a fine powder. The water, by this process, may be made to take up nearly eight hundred times its bulk of ammoniated gas, and is three times the strength of the liquor ammoniæ of the London Pharmacopeia.

2. The second ingredient is distilled spirits of Rosemary, prepared by taking two ℥s of the tips or small leaves of fresh Rosemary, and eight pints of alcohol, which are infused together for twenty-four hours in a covered vessel, and after adding as much water as will first prevent the empyreumatic smell, distil over *seven* pints.

3. The third ingredient is spirit of camphor, made as follows: To four oz. of pure camphor, add two pints of alcohol, so as to dissolve the camphor, which solution should be filtered.

Of the preceding ingredients, Dr. Granville prepares two ammoniated lotions, differing in strength; viz.

The Milder Ammoniated Lotion.

(Lotio ammoniata mitior.)

R. Liquoris ammoniæ fortissimi, 3j.

Spiritus rosmarini, 3vj.

—— camphoræ, 3ij.

Misce

The Stronger Ammoniated Lotion.

(Lotio ammoniata fortior.)

R. Liquoris ammoniæ fortissimi, ℥x.

Spiritus rosemarinæ, ℥ss.

—— camphoræ, ℥ij.

Misce.*

As to the method of applying these counter-irritants, we shall use the words of their inventor, Dr. Granville. "First impregnate with the colourless and transparent liquid, either a piece of linen folded six or seven times to the size of the part intended to be covered, or a piece of thick and coarse flannel; and, secondly, lay either of them on the spot, pressing with the hand at the same time, very steadily and firmly, the said linen or flannel, over which there should be placed a thick towel, doubled several times, so that not only the evaporation of the lotion may be impeded, but the hand employed in pressing the application to the part may not suffer damage from any evaporation, or from contact with the liquid. In some parts of the body more convenient than the others, the readiest and most effectual mode of pressing down the application is, by tying over it a towel or thick bandage; but to this mode there is the objection that we cannot, under certain circumstances, infect the part as quickly and as often as required, so as to judge, from the effect of the application, when to stop, or how long to persevere in using it."†

When applied to the temples, forehead, jaws, or chin, great care must be taken to prevent the effluvia from affecting the eyes and nostrils. The skin should be examined in a minute or two after the application, in order to see the effect produced. When the pain is severe after the lotion is removed, simple diluent washes, and warm water is perhaps the best, will speedily allay it. Vesication will generally take place in from two to five minutes, and even in a minute in some persons: but where cauterization is requisite, ten, or twelve minutes may be necessary. The irritated or denuded surface may be dressed according to existing indications, with a simple poultice, tepid fomentation, cerate or irritating ointments.

The *Pommade Ammoniacale* of M. Goudret, which belongs to this series of counter-irritants, is made as follows:

* Dunglison, *New remedies*, p. 504.

† GRANVILLE. Counter-irritation, its Principles and Practice, p. 39.

R. Adipis, ʒvij.

Olei amygdalarum, ʒiss.

Ammonia liquidæ (25°) ʒvj.

Melt the lard, mix the oil with it, and pour them into a wide-mouthed vial with a glass stopper; then add the ammonia, close the bottle, mix the contents together by shaking, and keep the mixture in a cool place. If the skin is to be excited, perspiration to be re-established, or some sub-cutaneous engorgement to be dissipated, light and hasty practices accomplish these objects. If a rubefacient effect is desirable, spread the ointment thickly on linen, and apply it for one or two minutes. For vesication, continue the application for five or six minutes; and by a farther continuance, absolute cauterization is speedily produced.

These powerful preparations are used in a great variety of diseases, and in all cases when speedy and powerful counter-irritation is demanded; as in Neuralgia and rheumatic affections generally, both acute and chronic; spasms, convulsions, gout, paralysis, inflammatory diseases of internal organs, diseases of the heart, sprains, &c. &c.

Instantaneous Vesication.

The French practitioners have a process for raising an instant blister. A piece of linen or paper being cut to the requisite size, is immersed in spirits of wine or brandy, or even cologne water, or spirits of camphor, whichever is most readily obtained. It is then laid on the part to be blistered, taking care that the moisture does not wet the surrounding surface. The flame of a lighted taper is then applied quickly over the surface of the paper, so as to cause ignition, which is almost instantaneous. At the conclusion of this operation, the cuticle is found detached and raised from the true skin beneath.

Ointment of Iodine and Calomel.

R. Iodini, partem j.

Hydrargyri chloridi mitis, partem iss.

Axungia, partes xlvij.

The iodine to be first united to the lard, and then the calomel added.

MM. Plane and Souberain, by whom this ointment is proposed, state that these are the only proportions that can be used, without leading to decompositions that will defeat our intentions.

Ointment of Protoiodide of Mercury, &c.

R. Protoiodidi Hydrargyri, gr. vj.
 Acetatis Morphæ, gr. viij.
 Axungiæ, ʒj.
 Misce, fiat unguentum.

Recommended by the French physicians in glandular swellings. The morphia is believed by them to prevent the unpleasant effects of iodine when used in the shape of ointment.

Pelletan.

Ointment of Protoiodide of Mercury.

R. Protoiodidi Hydrargyri, ʒj.
 Axungiæ, ʒiss.
 Fiat unguentum.—Used to accelerate the cicatrization of inveterate venereal ulcers.

An ointment may be made in the same proportions of the *deutoiodide of mercury*. Of course to be used with more caution.

Ointment of Iodate of Zinc.

R. Iodatis Zinci, ʒj.
 Axungiæ, ʒj.
 Fiat unguentum.—A drachm to be rubbed on the tumour twice a-day.

Dr. Ure recommends the above as a substitute for the ointment of hydriodate of potassa in glandular swellings.

Ointment of Hydriodate of Potash.

R. Hydriodatis Potassæ, ʒj.
 Cerati simplicis, ʒj.
 Fiat unguentum.

To be applied to scrofulous swellings, where the skin is unbroken.

Iodine Ointment.

R. Deuto-iodureti hydrargyri, gr. xv.

Adipis preparatæ, ℥ij.

Olei limonis, gtt. xx.

Fiat unguentum.

Employed as a curative application to syphilitic tubercles, chronic ulceration and other sequellæ of venereal. Its great activity requires caution in its use.

Ratier.

Simple Iodine Ointment.

R. Iodinæ, gr. iij.

Adipis, ℥ij.

Fiat unguentum.

Used as the preceding preparation.

Iodine Liniment.

R. Linamenti saponis compositi, ℥j.

Tincturæ Iodinæ, ℥j.

Misce.

Ointment of Iodide of Potassium.

R. Potassæ iodidi, ℥ss.

Adipis, ℥iss.

Ut unguentum.—For Tetter and the kindred eruptions.

Lugol's Ointment of Ioduretted Iodide of Potassium.

R. Iodinæ, gr. xij.

Potassæ iodidi, ℥iv.

Adipis, ℥ij.

Fiat unguentum.—Used in scrofulous ophthalmia, ulcers, &c.

Ointment of Ioduretted Iodide of Potassium with Opium.

R. Iodinæ, gr. xvj.

Potassæ iodidi, ℥j.

Tincturæ opii, ℥ij.

Adipis, ℥ij.

Fiat unguentum. Applied to scrofulous and other ulcers.—*Lugol.*

Ointment of Iodide of Arsenic.

R. Iodidi arsenici, gr. iij.

Adipis preparatæ, ℥j.

Fiat unguentum.

Employed in Lepra, Impetigo, and other cutaneous affections.

Ointment of Cyanide of Mercury.

R. Hydrargyri Cyanidi, gr. xij.

Adipis, ℥j.

Fiat unguentum.

Ointment of Iodide of Sulphur.

R. Iodureti Sulphuris, partes 5.

Adipis, partes 96.

Fiat unguentum.—In Diseases of the Skin.

Magendie.

Ointment of Bromium.

R. Adipis, ℥j.

Hydrobromatis Potassæ, gr. xxxiv.

Fiat unguentum.

Decoction of Turpentine and Cantharides.

R. Pulveris Cantharidum, ℥i.

Olei Terebinthinæ, ℥viiij.

Boil them together for three hours, then filter.

This is the process usually followed by our apothecaries, but we conceive thirty minutes' boiling would be quite sufficient, and the cantharides be less likely to be injured by the heat. It was first made by Dr. Hartshorne of this city, who has used it with great success in the typhoid state of fever. It is merely rubbed over the surface of the body.

Kentish Ointment.

R. Unguenti Resinæ flavæ, ℥ij.

Olei Terebinthinæ, ℥ij. vel ℥ss.

Fiat unguentum.

This preparation has long been celebrated in burns and scalds. We must be careful to avoid touching the sound skin, for turpentine occasionally inflames it severely.

Turpentine Liniment.

- R. Olei Terebinthinæ, ℥ij.
 — Olivarum, ℥ij.
 Tincturæ Camphoræ, ℥j.
 Aquæ Ammoniæ, ℥j.
 Fiat linimentum.

Creosote Ointment.

- R. Creosotis, gtt. xv.
 Cerati simplicis, ℥j.
 Olei cinnamomi, gtt. vj.
 Fiat unguentum.—In ulcers of a chronic character, and in cutaneous eruptions.

Digestive Ointment of the French Hospitals.

- R. Unguenti terebinthinæ comp. ℥ij.
 Vitellum ovi j.
 Olei olivarum, ℥ij.
 Misce.—Used in dressing indolent wounds and ulcers. *Ratier.*

Sulphur Ointment.

- R. Adipis suilli, ℥bj.
 Sulphuris sublimati, ℥viij.
 Fiat unguentum.—Used in friction—three to six drachms at a time.

Mercurial Cerate.

- R. Unguenti Hydrargyri, ℥iv.
 Cerati simplicis, ℥x.
 Fiat unguentum.—A dressing for venereal ulcers. *Ratier.*

Ointment of Belladonna.

- R. Extracti Belladonnæ, ℥ij.
 Aquæ destillatæ, ℥ij.
 Adipis Suillæ, ℥ij.
 Misce.—Employed by M. Chaussier as a direct application to the neck of the uterus, in case of rigidity of that part. *Ratier.*

Liniment with Camphor, &c.

R. Spiritus Camphorati, ℥iij.

Acidi acetici fortis, ℥j.

Fiat embrocatio.—To be applied to bruises, rheumatic parts, &c.

Soap Liniment, &c.

R. Linimenti saponis, ℥ij.

Aquæ Ammoniæ, ℥j.

Tincturæ Opii, ℥ss.

Fiat linimentum.—To be used as above.

Liniment with Ammonia and Olive Oil, vulgo, Volatile Liniment.

R. Olei Olivarum,

Aquæ Ammoniæ, aa. ℥j.

Misce.—This preparation is frequently employed in inflammation of the throat and other similar cases.

Liniment of Ammonia and Tartarized Antimony.

R. Linimenti Ammoniæ, ℥j.

Tartratis antimonii, ℥j.

Misce.—This is often applied to indolent tumours, and other swellings.

Liniment of Cantharides, &c.

R. Camphoræ, ℥iij.

Solve in decocti Cantharidis cum terebinthina, ℥ss.—et adde—

Unguenti Hydrargyri fortis,

———— simplicis, aa. ℥j.

Fiat linimentum.—This preparation might be employed in cases of low fever, where it is desirable to rouse the system, and at the same time affect it with mercury; also in tumours.

Lotion of Cyanide of Potassium.

R. Potassii cyanidi, gr. xij.

Misturæ amygdalarum, ℥vj.

Fiat lotio.—Used in Lichen, and other chronic eruptions.

Solution of Caustic Potash.

R. Potassæ puræ, ℥iss.

Aquæ destillatæ, ℥ij.

Fiat solutio.—Signa.—To be applied to the spine in cases of tetanus.

Dr. Hartshorne speaks in high terms of the beneficial effects of this solution in the above disease. The best mode of applying it, so as merely to produce a rubefacient action, is to tie a piece of sponge to a stick or common table fork, and use it as a brush to apply the solution from the nape of the neck to the sacrum. When the solution is tolerably strong, the effect is powerful.

Compound Opiate Liniment.

R. Tincturæ Opii,

Spiritus Ætheris Sulphurici,

——— Camphoræ, aa. ℥ij.

Fiat embrocatio.—To be applied to parts affected with rheumatism.

It is very useful in odontalgia, pain in the region of the ear, &c.

Opiate Liniment.

R. Olei olivarum, ℥ij.

Tinctura Opii, ℥ij.

Aquæ acetatis plumbi, ℥ss.

Misce.

Embrocation of Acetate of Ammonia.

R. Ammoniæ carbonatis, ℥ij.

Acidi acetici, Oij. vel q. s. ad saturandum:

Spiritus tenuis, Oiss.

Mix the ammonia with the vinegar, and after effervescence ceases, add the spirit.

To be applied to rheumatic affections; or mixed with linseed meal, it may be used as a poultice to chronic inflammations of the joints, &c.

Compound Liniment of Tar.

R. Petrolei Barbadosensis, ℥j.

Camphoræ, ℥ss.

Alcoholis, gtt. xl.

Fiat linimentum.—Powder the camphor by adding the spirit, and then mix the whole in a mortar.

To be applied round the joints in cases of a chronic kind, and especially where thickening of the ligaments is suspected.

Creosote Liniment.

R. Creosoti, gtt. x.

Olei Olivarum, ℥j.

Fiat linimentum.—In Herpes, &c.

Creosote Lotion.

R. Creosoti, gtt. x.

Aceti, ℥ij.

Aquæ destillatæ, ℥ij.

Fiat lotio.—Applied in phagadenic ulceration.

Camphorated Oil.

R. Camphoræ, ℥j.

Olei Olivarum, ℥iv.

Tere, et fiat solutio; tunc adde,

Olei Terebinthinæ, ℥iss.

To be applied by rubbing with a piece of flannel; in deep-seated rheumatic pains.

Epithem with Tincture of Cantharides.

R. Pulveris Cantharidum, ℥j.

Spiritus vini rectificati, ℥iv.

Digest with a gentle heat for two days.—Strain and add—

Camphoræ, ℥ss.

To be rubbed on the parts affected. In paralysis.

Epithem with Camphor and Carbonate of Ammonia.

- R. Spiritus Camphorati, ℥iv.
 Carbonatis Ammoniaë, ℥ss.
 Olei Juniperi,
 — Succini, āā. ʒij.

Misce.—To be employed in the same cases as the former.

Liniment of Oil of Amber, &c.

- R. Olei Succini rectificati, ʒij.
 — Olivarum, ℥ss.
 Tincturæ Opii, ʒij.
 Spiritus vini Gallici, ℥iij.

Fiat linimentum.—Signa.—To be rubbed between the shoulders. In pertussis.

Cayenne Pepper and Alcohol Dilute.

- R. Pulveris Capsici, ʒij.
 Spiritus vini diluti, ʒviiij.

Misce.—This preparation forms an excellent rubefacient in cases of low fever. Dr. Chapman recommends it strongly in sciatica, and other rheumatic affections.

In dyspepsia, and some diseases of the head attended with *cold feet*, socks dusted with Cayenne pepper are very useful.

CATAPLASMS, &c.

Cataplasm of Pepper, &c.

- R. Semen Sinapeos contus: ℥ss.
 Piperis nigri,
 Pulveris Zingiberis, āā. ʒj.
 Syrupi simplicis, q. s. ut fiat cataplasma.

To be applied to the region of the stomach in severe pain or sickness; or, to the soles of the feet when revulsion is desirable.

Spice Plaster.

- R. Pulveris Caryophyllorum,
 ——— Cinnamomi,
 ——— Capsici annui, āā. ℥ss.

Misce, et adde—

Farinæ,

Vini Gallici, āā. q. s.—ut fiat cataplasma.

Let the plaster, thus prepared, be made hot, and applied to the epigastric region.

Dr. Parrish was accustomed to speak in the highest terms of this plaster in cholera.

Mustard Cataplasm.

- R. Pulveris Sinapeos, ℥ij.
 Acidi acetici tepidi, q. s.—ut fiat cataplasma.

The uses of the mustard poultice are too well known to require description. We will only observe, that it is frequently made by adding an equal quantity of wheat or rye flour; this should always be done for children, or other persons of delicate skins. It cannot, generally, be borne longer than a half hour or hour.

Iodine Cataplasm.

- R. Tincturæ Iodinæ, ℥ss.
 Pulveris lini, ℥j.
 Farinæ avenæ, ℥iij.
 Aquæ q. s.—ut fiat cataplasma.

Applied to indolent ulcers, (especially those of a scrofulous character) and to tumours of the same class.

Garlic Cataplasm.

The Garlic Cataplasm is employed with the same intention as the mustard, though it is much less active. It may be made by bruising the cloves of garlic, and mixing them in a common poultice of bread and milk. Onions may be substituted for garlic.

Cataplasm of Common Salt, &c.

R. Pulveris Lini,

Micæ panis singulorum partes æquales, Aquæ Sodæ muriatis saturatæ, q. s.—ut fiat cataplasma.

To be applied to indolent strumous swellings, and enlargements of the joints. When it produces irritation or inflammation, the common bread and milk poultice must be substituted; and the other re-applied when the skin will admit of it.

Cataplasm of Yest.

R. Farinæ, ℥j.

Cerevisiæ fermenti, Oss.

Misce.—This cataplasm is useful in stimulating foul ulcers and gangrenous wounds, and removing the unpleasant odour. It should be made with fresh yeast, and applied warm.

Cataplasm of Charcoal.

This cataplasm is made by stirring into the common bread and milk poultice as much finely powdered charcoal as it will bear. It is a very useful application to offensive ulcers; removing in a short time the fætor, and disposing them to a more healthy aspect.

Cataplasms are made of carrots or turnips, by simply boiling them to softness, and beating them to a pulpy mass. Both of these are said to furnish a good poultice for cancerous and other painful sores.

Cloves.

Quilt some powdered cloves in flannel, and wring them out in hot brandy. Thus prepared, they form an excellent external application in cases of colic, and great irritability of the stomach, as it occurs in cholera.

Emollient Cataplasm.

R. Farinæ seminum Lini,

————— Hordei, āā. partes æquales.

Aquæ vel lactis, q. s.—ut fiat cataplasma.

The milk or water may be warmed if desirable. Applied to inflammatory tumours.

Antiseptic Cataplasm.

R. Cataplasomatis emollientis, ℥viij.

Pulveris Kino,

——— Camphoræ, āā. ʒij.

Fiat cataplasma.—Applied to wounds and ulcers that have assumed an unhealthy action. *Hotel Dieu.*

Resolvent Cataplasm.

R. Cataplasomatis emollientis, ℥iv.

Acetatis plumbi, ℥j.

Muriatis ammoniæ, ʒss.

Fiat cataplasma.—Applicable to tumours that require a gentle excitement. *Ratier's Formulary.*

Warm Strengthening Plaster.

R. Picis Burgundici, ℥v.

Unguenti Cantharidum, ℥j.

Fiat emplastrum.—This plaster is applied in the same way, and used in all the cases for which Burgundy Pitch is prescribed. It is irritating, and on some persons will draw a blister.

BATHS, FOMENTATIONS, &c.

The ordinary application of cold, tepid and shower baths is too well known to require any particular notice. The hot bath should be of a temperature of 93 to 96 degrees of Fahrenheit. It is designed to alleviate general or local irritation, and to induce repose and sleep. It is especially adapted to fevers with deficient capillary circulation, rheumatism, herpetic eruptions, paralytic affections, &c. The lower temperature is best adapted to cutaneous complaints, and to bring on relaxation of the skin during febrile irritation: the warmer temperature is suited to paralysis; and, as a general rule, more heat is desirable to act upon deep seated than on superficial diseases.

Vapour Baths are produced by the evaporation of boiling water in a suitable apparatus. They are used, like ordinary warm baths, in rheumatic and cutaneous

affectons and in every case wherein profuse perspiration is desirable. They are avoided in persons labouring under pulmonary or cardiac disease, and in cases of exhaustion from any cause.

Local Baths are admirably adapted to parts which are swelled and painful, and may consist of tepid water only, or their strength may be increased by the addition of saline, emollient, narcotic or stimulant substances.

Sulphurous Water, or Artificial Barèges Bath.

R. Sulphureti Potassæ, ℥j.

Aquæ, cong. xxx.

Mix for a bath. In cutaneous eruptions.

Sulphureo-Gelatinous Bath.

R. Sulphureti Potassæ, ℥ij vel ℥iv.

Aquæ, cong. xxx.

Add to this solution,

Ichthyocollæ, ℥j. vel ℥ij in Aquæ bullientis solutæ, ℥x.

This bath, according to Dr. Green, (*Diseases of the Skin*, p. 240.)

“is preferable to the artificial Barèges Bath, as it is neither irritating nor apt to occasion feverishness, which the common sulphureous water bath is. A cheaper and not less efficacious gelatine may be procured by dissolving from a pound and a half to two pounds of parchment-clippings in water by long boiling, or by using a neat’s or a calf’s foot for the purpose.”

Emollient Bath.

To an ordinary tepid water bath, add a large cupful of thick gruel or paste, and mix it well with the water. These baths are very efficacious in prurigo, eczema, lichen and impetigo.

Nitro-muriatic Acid Bath.

R. Acidi nitrici, ℥ij.

Acidi muriatici, ℥j.

Mix, and add to the water of a tepid bath, which should then be about as sour as distilled vinegar.

Sublimate Bath.

R. Hydrargyri bichloridi, ℥ij vel ℥j.

Aquæ tepidæ, cong. xxx.

Mix for a bath. Prescribed by the French physicians in syphilitic affections, when it is desirable to avoid the action of the medicine on the stomach. This plan is seldom resorted to, on account of the great risk of employing corrosive sublimate in such large quantity.

Alkaline Bath.

R. Potassæ carbonatis, ℥iv vel ℥viij.

Aquæ tepidæ, cong. xxx.

Mix for a bath. Very useful in promoting desquamation from the skin, and in allaying itching in various forms of Prurigo.

Artificial Harrowgate Bath.

R. Sodæ muriatis, ℥ij.

Magnesiæ sulphatis, ℥iij.

Potassæ sulphureti, ℥j.

Aquæ, cong. xxx.

Mix for a bath.

Fomentation is a yet more restricted mode of applying fluids to the surface of the body, which is done by means of a sponge, flannel or soft cloth. To be efficacious they should be often repeated, and not allowed to get cold.

Stimulant Fomentation.

R. Vini rubri, Oij.

Mellis, ℥iv.

Misce.—This is an active fomentation, and is applied to parts in a low state of vitality. *Ratier.*

Saponaceous Fomentation.

R. Spiritus vini gallici, Oij.

Saponis, ℥j.

Misce.—A resolvent in indolent tumours, and in sprains. *Ratier.*

Narcotic Fomentation.

R. Gum Opii, ℥ij.

Aquæ, Oj.

Solve.—Adapted to painful affections of the joints, as chronic rheumatism, &c. *Ratier.*

Narcotic Fomentation.

R. Foliorum Dulcamaræ, ℥ij.

Capsularum papaveris, ℥ij.

Aquæ ferventis, Oj.

Fiat infusum.—Employed as a soothing application to irritable surfaces.

Mustard Fomentation.

R. Farinæ sinapeos, ℥iv.

Aquæ ferventis, Oj.

Misce.—By wetting cloths in this fluid and applying them to the skin, irritation may be prolonged to any requisite degree.

Tobacco Fomentation.

R. Foliorum tabaci, ℥ij.

Aquæ ferventis, Oj.

Misce.—Employed in Psora, but is apt to produce nausea and vomiting.

Fumigations are for the most part used to purify infected air by absorbing or otherwise counteracting deleterious gases. They are also employed in diseases of the skin, and may be sometimes substituted for a local bath. The ordinary process for fumigating apartments with chlorine, has been already described. (Page 86.)

Fumigations of sulphur, cinnabar, and mercury are also in common use, for medicinal purposes, and are readily applied with a proper apparatus.

For *Inhalations*, see Expectorants, p. 84.

CAUSTICS, OR ESCHAROTICS.

Caustics or cauteries are divided into the actual and potential. The first consists of fire itself; and the second of those substances which destroy the living solids, either by excessive stimulation, or, as some have supposed, by producing a chemical decomposition.

Common Caustic with Opium.

R. Potassæ cum Calce, ʒij.

Pulveris Opii, ʒss.

Saponis mollis, q. s.

Misce.—This preparation is employed for the same purpose as the common caustic.

Potassa.

The *Causticum commune acerrimum* is the pure Potash: when mixed with lime, its activity is lessened, and it is then called *Causticum commune mitius*. For the best mode of applying this remedy, see the remarks on Issues. As this caustic spreads rapidly and eats deeply, care should be taken not to leave it on too long, when in the neighbourhood of large blood-vessels. When, however, we wish to arrest its effects, we may neutralize it at once by a little vinegar.

Lunar Caustic.

This caustic is the *Nitras Argenti*, deprived of its water of crystallization, fused and cast into moulds. Its mode of application is universally known. It is free from the inconveniences of the former.

Burnt Alum.

The *Alumen ustum* is made by depriving this salt of its water of crystallization by heat. It is applied in powder to the fungous granulation of ulcers; and, mixed with an equal part of powdered sugar, is used to remove specks or *nebulæ* of the cornea.

MINERAL ACIDS.

The mineral acids are all escharotics, though seldom employed, because of the difficulty of managing fluids when used in this way. *Nitric Acid* diluted is sometimes applied to extensive foul ulcers, in warm weather, to destroy the animalculæ which are apt to be generated in such sores.

Solution of Muriate of Ammonia, &c.

R. Muriatis Ammonia, ℥j.

Aceti destillati, ℥ij.

Aquæ fontanæ, ℥iv.

Fiat solutio.—To be applied to venereal warts.

Bell on Venereal.

White Oxide of Arsenic, or Arsenious Acid.

R. Acidi Arseniosi albi, ℥j.

Aquæ fontanæ, ℥ij.

Fiat solutio.

This solution is occasionally applied to cancerous sores: it was formerly used in the shape of powder or paste, as an empirical remedy. But it is now seldom employed, from the circumstance of its producing constitutional irritation. Sir Astley Cooper mentions cases where, from its absorption when applied to open sores, partial if not complete paralysis of the limbs was the consequence.

Ointment of Arsenic.

R. Acidi Arseniosi pulveris, ℥j.

Adipis Suillæ,

Cerati cetacei, āā. ℥vj.

Melt the cerate and lard over a slow fire, and then pour it on the white oxide of arsenic, and triturate it in a glass mortar until it is *intimately* united.—*Pharm. Chirurg.*—For cancers, &c.

The same care is required in its use as in the solution.

Dubois's Arsenical Powder.

R. Oxydi arsenici albi, ℥ss.

Sulphureti Hydrargyri rubri, ʒj.

Resinæ pterocarpi, ʒss.

Fiat pulvis.—Chiefly used in the French hospitals as a caustic to cancerous affections.

Ratier.

MOXA.

This is another mode of using the actual cautery; and, as it is engaging the attention of physicians, we insert the usual mode of applying it.—The Chinese employ this remedy by means of the Mugwort. They separate this soft lanuginous substance by beating, and form it into a cone; the base of this is attached, by being wetted, to the skin, and the point set on fire.

Dr. Boyle recommends dipping cotton in a strong solution of nitrate of potash, and enclosing pieces of this thoroughly dried in paper cylinders, about one inch in diameter, and another inch in height. Dr. Burne employs wood in a state of dry-rot, touch-wood. This only requires to be shaped properly, and burns, he says, with certainty and steadiness, and is preferable to the other modes suggested;—some of which, as common cotton, or lint, requiring the use of the bellows, and when they have been dipped in nitre, demanding care to prevent the scintillations from injuring the surrounding parts.

The cylinders or cones of touch-wood may be confined to the skin by a little adhesive plaster affixed to the edges.

It is proposed to place pieces of cold iron in the neighbourhood of the part while the moxa is burning, in order to render the sensations more vague, and therefore less painful.

Moxa is used in diseases of the spine, paralysis, &c. &c.

STIMULATING OINTMENTS, &c.

Ointment of Red Precipitate.

R. Hydrargyri oxydi rubri, ℥j.

Terebinthinæ venetæ, ℥j.

Fiat unguentum.

This preparation may be applied to indolent sores, and is said, by some practitioners, to be a specific in itch.

White Precipitate.

R. Hydrargyri precipitati albi, gr. xv.

Nitratis Potassæ, ℥ss.

Florum Sulphuris, ℥j.

Bene teranturet adde—

Axungię Porcinæ, ℥ij.

Fiat unguentum.

Dr. Chapman remarks, that this ointment has long been celebrated in the cure of itch.

Solution of Phosphoric Acid.

R. Acidi phosphorici, ℥j.

Aquæ destillatæ, ℥viiij.

Fiat solutio.

Dr. Lentin, a German Physician, has employed this solution on compresses applied to ulcers situated over carious bones, and he imagines with decided benefit.—*Spillan's Supplement to the British Pharmacopœias.*

Corrosive Sublimate and Lime-Water.

R. Chloridi Hydrargyri corrosivi, gr. j vel iij.

Aquæ Calcis, ℥j.

Fiat solutio.—*Signa.*—The yellow wash.

For ordinary purposes one grain to the ounce will be sufficient. It is employed in phagedænic and venereal ulcers.

Calomel and Lime-Water.

R. Hydrargyri Chloridi mitis, ʒj.

Aquæ Calcis, ʒiv.

Misce.—Signa.—The black wash.

This prescription is used in the same cases as the other. The red-oxide in that, and the black-oxide of mercury in this, are precipitated by the lime-water. They are sometimes administered in the form of injections, in gleet and obstinate gonorrhœa.

Anti-psoric Lotion of Dupuytren.

R. Sulphureti potassæ, ʒiv.

Aquæ, Oj.

Acidi sulphurici, ʒiv.

Misce.—To be applied two or three times a-day on the parts covered with pustules, and with this treatment the simple bath is conjoined.

Lotion of Myrrh.

R. Tincturæ Myrrhæ,

Liquoris Calci, āā. ʒj.

Misce.

This lotion is recommended by Dr. Kirkland for taking down fungous flesh. In cases of fungous brain, he found it extremely serviceable applied to the excrecence on dossils of lint.

Powder of Verdigris and Calomel.

R. Acetatis Cupri,

Hydrargyri chloridi mitis, āā. ʒj.

Fiat pulvis subtilissimus.—For chancre and indolent ulcers.

Powdered Rhubarb, &c.

R. Pulveris Rhei,

——— Ipacacuanhæ, aa. ʒj.

Fiat pulvis.—To be dusted on sarcomatous and warty tumours.

Rhubarb alone forms an excellent application to indolent ulcers.

Powder of Savin, &c.

R. Pulveris Sabinæ,

——— Acetatis Cupri, aa. ʒj.

Fiat pulvis.—To be dusted on sores accompanied with fungus excrescences.

Ointment with Carbonate of Ammonia, &c.

R. Carbonatis Ammoniæ, ʒss.

Unguenti simplicis, ʒss.

Fiat unguentum.—To be used in indolent ulcers of a scrofulous character.

ISSUES AND SETONS.

These remedies are employed as substitutes for blisters, where a permanent discharge is desired. We therefore insert the best rules for their employment.

There are three kinds of issues in common use, by either of which we may procure a discharge of purulent matter for any length of time that may be required. They are applied in different situations, according to circumstances, and some one of the following parts is generally chosen; the nape of the neck; the hollow of the deltoid muscle, and between the shoulders or ribs; in the inferior extremities they are most conveniently formed in the inside of the leg, either above or below the knee.

1. *Blister Issue.*

To form this issue it is merely necessary to apply a blister to any part in the usual manner; the skin being thus removed, the discharge should be continued by means of stimulating ointments, as Unguentum Cantharidum, Ung. Basilici, Ung. Sabinæ, etc. These may be alternated with mild liniments, in order to prevent the ulceration and consequent discharge from becoming excessive.

2. *The Pea Issue.*

There are two modes of forming this issue; the first is made by means of a lancet; the part is merely taken in the thumb and finger, and an incision made by passing the lancet through the skin. The wound should be kept open by means of orange peas, the common pea, a bean, or a bit of orris root.

The following extracts from the *Surgeon's Vade Mecum*, will best exemplify the manner of forming the issue with caustic:—

“The part is first to be covered with a piece of adhesive plaster, in the centre of which a circular hole has been cut of the size of the intended issue; in this hole the caustic, *kali purum*, made into a paste with soap, is to be placed. The whole is then to be covered with another plaster, secured by a bandage, and suffered to remain for *ten or twelve* hours. Generally in two or three days an eschar begins to separate, when the opening should be filled with the substance made choice of.” These substances are enumerated above.

3. *The Seton.*

To form this artificial discharge, take the seton needle, to which is attached a skein of silk, the latter being partially anointed with some oleaginous substance. The needle is to be passed completely through the part chosen for the operation, after which it is removed, and the ends of the silk left hanging from the wound. It should be dressed every day, or twice a-day, with some mild ointment; and if this be not sufficient to keep up the discharge, the more stimulating unguents may be used, as mentioned when speaking of the blister issue.

BOUGIES.

Although directions for forming bougies may appear out of place under this head, yet as country physicians may often be obliged to make these articles, we can not forbear inserting the formulæ, and we see no more appropriate place in the work.

R. Ceræ albæ, ℥xvj.

Terebinthinæ venetæ, ℥iss.

Olei olivarum, ℥i.

Melt the wax and turpentine over a slow fire, and then add the oil.

Bougie, with Red Lead.

R. Olivarum Olei, Oj.

Ceræ flavæ, ℥x.

Oxidi Plumbi rubri, ℥xv.

Boil over a slow fire until the minium is perfectly dissolved, which will be in about four or six hours.

Dr. Swediaur's Bougie.

R. Ceræ flavæ, ℥bj.

Cetacei ceti, ℥iij.

Plumbi acetatis, ℥v.

These are to be boiled like the former until the composition is of a proper consistence for forming bougies.

Sharp's Bougie.

R. Diachyli cum pice Burgundica, ℥ij.

Argenti vivi, ℥j.

Antimonii sulphureti ppt: ℥ss.

The quicksilver to be previously dissolved in balsam of sulphur or in honey, and added to the plaster when melted in a moderate heat.

Bell on Venereal.

In order to form a bougie, a slip of linen, an inch in width at one end, and an inch and a quarter at the other, and from ten to twelve inches long, must be dipped into the melted composition, taking care that it be evenly applied; it is then to be carefully folded up

into the proper longitudinal shape, and rolled firmly between two marble slabs, till perfectly smooth and even.—*Pharmacop. Chirurgica*.

The caustic bougie is made by simply rolling in the centre of the wax bougie, at the extremity, a small piece of lunar or common caustic. The former is more manageable, and therefore generally preferred. Bougies are sometimes dipped in Basilicon or mercurial ointment, oil of turpentine, &c., in order to render them more stimulating when employed in the treatment of gleet, &c.

CLASS X.

STIMULANTS.

IN the use of stimulants, it is for the most part advisable to begin with small doses, and augment them as circumstances may require. There are cases, however, in which it is necessary to give them plentifully *ab initio*.

It will be often requisite to change not only the stimulating substances, but the part of the body to which they are applied; thus when the stomach fails, the rectum and skin may be acted upon with advantage.

PILLS, POWDERS, &c.

Pills of Camphor, &c.

R. Pulveris Lyttæ, gr. xvij.

——— Opii,

——— Camphoræ, āā. gr. xxxvj.

Conservæ rosarum, q. s.

Ut fiant pilulæ xxxvj.—Of which one or two may be taken every night on going to bed.

Designed for impotency arising from general debility.

Pills of Deuto-iodide of Mercury.

R. Deuto-iodidi Hydrargyri, gr. j.

Extracti Juniperis vel Glycyrrhizæ, gr. xij.

Divide in pilulas viij.—Signa.—Two in the morning and two at night; dose to be augmented gradually to four in the morning and four at night.

Pills of proto-iodide of mercury may be made in the same proportions and given in the same doses. In *scrofula*.

Pills of Turpentine, &c.

R. Pulveris Guaiaci, ʒj.

Terebinthinæ venetæ, q. s.

Ut fiant pilulæ xv.—Signa.—One three times a-day.

These pills have a powerful effect upon the urinary organs. They are employed in gleet and læucorrhœa, and frequently produce the most obstinate strangury.

Pills of Cayenne Pepper.

R. Pulveris Capsici, ʒj.

Micæ panis,

Aquæ destillatæ, āā. q. s.

Ut fiant pilulæ xij.—Signa.—One three or four times a-day. In debility of the stomach, resulting from intemperance.

Aromatic Pills.

R. Carbonatis Ammoniæ,

Capsici,

Caryophylli,

Macis, āā. ʒj.

Olei Carui, gtt. v.

Extracti Gentianæ, gr. xij.

Syrupi simplicis, q. s.

Ut fiant pilulæ xx.—Signa.—One every two hours in gout of the stomach.

Dr. Parrish.

Tincture of Cantharides, &c.

R. Radicis Rhei contusæ ʒiss.

Guaiaci, gr. xlv.

Gummi Laccæ, ʒss.

Cantharidum contusarum, ʒj.

Spiritus vini rectificati, ʒxij.

Fiat infusio, et cola. Dose from thirty to fifty drops, night and morning, in a draught of water. In gleets.

From the active nature of these articles, we would recommend caution in the use of them.

Mixture of Carbonate of Ammonia, &c.

R. Carbonatis Ammoniaë, ℥iss.

Pulveris sacchari albi,

—— gummi Acaciaë, aa. ℥iss.

Spiritus Lavendulæ compositi, ℥ij.

Aquæ destillatæ, vel Menthæ, ℥iv.

Fiat mistura.—Signa.—A table-spoonful for a dose, every one or two hours.—And, if desirable, Tinct. Opii, gtt. xl vel l. may be added to the mixture.

Draught with Valerian and Carbonate of Ammonia.

R. Pulveris Valerianæ, ℥j.

Carbonatis Ammoniaë, gr. xv.

Aquæ Cinnamomi, ℥ij.

Misce.—Fiat haustus.—To be taken every fourth hour. Suited to cases of nervous headach, and depression of spirits.

Mixture of Cubebs, &c.

R. Pulveris Cubebarum, ℥ij.

Carbonatis Sodæ, ℥ss.

Mucilaginis Acaciaë, ℥vj.

Aquæ Menthæ, ℥vj.

Fiat mistura.—Signa.—Take a table-spoonful every hour and a-half.

Mixture of Oxide of Bismuth, Cubebs, &c.

R. Pulveris Cubebarum, ℥ij.

Sub-nitratis Bismuthi, ℥ss.

Mucilaginis Acaciaë, ℥ss.

Syrupi simplicis, ℥vj.

Aquæ destillatæ, ℥vj.

Fiat mistura.—Signa.—Take two table-spoonsful three times a-day.

We have extracted these prescriptions from a paper of John Fosbroke, surgeon, Chitttenham; published in the Medical Recorder, April, 1825.

They are recommended, with other similar compounds, in various affections of the mucous membranes:—viz. of the bowels, bladder, trachea, &c.

There is not much general confidence reposed in cubebs in this country. Perhaps this skepticism as to their efficacy may not be well founded.

Camphorated Enema.

R. Enematis communis, ℥ij.

Camphoræ, ʒij.

Vitellum unius ovi.

Misce.—Signa.—One fourth part to be thrown up the rectum in the treatment of putrid fevers, *pro re nata*. M. Chaussier.

R. *Acetatis Ammonia*.—*Quantum placet*.

This article, it is said, will dissipate inebriety in less than five minutes. The dose may be safely ten grains.

Antiseptic Pills.

R. Camphoræ,

Nitratis potassæ,

Pulveris Gummi Arabicæ, aa. ʒj.

Mix, and divide into pills of four grains each.—Dose two pills every hour or two hours, when a tendency to gangrene exists.

Draught with Powdered Ginger, &c.

R. Pulveris Zingiberis, gr. xv.

Carbonatis Ammonia, gr. viij.

Spiritus Cinnamomi, ʒij.

Aquæ, ʒiiss.

Misce.—Signa.—To be taken at once. In cases of gout in the stomach.

Camphor Mixture.

R. Camphoræ, ℥j.
 Pulveris gummi Arabici,
 ——— Sacchari albi, āā. ℥iiss.
 Tincturæ Opii, gtt. xl.
 Aquæ Menthæ, ℥iv.

Fiat mistura.—Signa.—A table-spoonful to be given every two hours.

In reducing camphor to powder, (which must be done before it can be formed into a mixture,) it is necessary to add to it a few drops of alcohol.

Camphor in Milk.

R. Camphoræ, ℥j.
 Lactis bullientis, ℥iv.

Fiat solutio.—This preparation may be taken in the manner as directed in the former.

Camphor Julep.

R. Camphoræ, ℥j.
 Aquæ bullientis, ℥viij.

Fiat mistura.—This preparation should be set aside in a covered vessel for half an hour, and then strained.—Signa.—A table-spoonful for a dose *pro re nata*.

Camphor with Magnesia.

R. Camphoræ pulverizatæ, ℥j.
 Magnesiæ carbonatis, ℥ij.
 Aquæ destillatæ, Oj.

Fiat mistura.

This prescription answers the same purpose as the former. It is considered a more elegant preparation, and is given in the same doses. It should be suffered to settle; and the clear liquor, which is slightly impregnated with the camphor, decanted for use.

Camphor with Myrrh.

R. Camphoræ pulverizatae, ℥j.

Pulveris gummi Myrrhæ, ℥ss.

—— Sacchari albi, ℥ij.

Aquæ destillatæ, ℥vj

Misce.—Signa.—A table-spoonful to be given every two hours.

Mixture with Oil of Turpentine.

R. Olei Terebinthinæ, gtt. cxx.

Pulveris gummi Arabici,

—— Sacchari albi, āā. ℥ij.

Tincturæ Opii, gtt. lx.

—— Lavendulæ compositæ, ℥ij.

Aquæ Menthæ sativæ, ℥v.

Fiat mistura.—Signa.—A table-spoonful for a dose every two hours.

This remedy has gained great reputation in some of the low forms of fever. The author once employed it in the dose prescribed above, in passive hemorrhage from the bowels, with complete success; and in hemorrhage from the bladder, resulting from low fever, he knows it to have been given with equal advantage.

Linctus with Oil of Turpentine.

R. Olei Terebinthinæ, ℥ij.

Mellis optimi, ℥j.

Fiat linctus—Dose a tea-spoonful night and morning, with a draught of warm tea. In Sciatica.

Emulsion of Phosphorus.

R. Phosphori puri, gr. ij.

Mucilaginis Arabici, q. s.

Let these articles be carefully and effectually triturated in a mortar, and add gradually as much distilled water as will make an emulsion of ℥vj. then add—

Syrupi simplicis, ℥j.

Liquoris Hoffmani mineralis, gtt. xxx.

Fiat mistura.—Of which the usual dose is a table-spoonful every two hours.

This preparation was originally made by the celebrated Hufeland, and is now taken from Dr. Chapman's Therapeutics, p. 176, vol. 2.

The violence of the remedy, and its very great power, are generally known to physicians. We need hardly repeat the necessity there is for the greatest caution in its administration.

It is sometimes given in sweet oil, but this is said to be extremely nauseous. A saturated solution of ether has also been employed, and is perhaps as neat a mode as any other. This solution contains about eight grains to the ounce.

Mixture of Camphor and Guaiacum.

R. Tincturæ Opii Camphoratæ,

——— Guaiaci ammoniatæ, aa. $\overline{3}$ j.

Fiat mistura, et signa.—Two tea-spoonsful every two hours. In misplaced and retrocedent gout.

Aromatic Spirit of Hartshorn.

R. Spiritus Ammoniæ aromatici, $\overline{3}$ ij.

Signa.—Twenty drops to be given in half a wine-glassful of water, *pro re nata*. In sudden exhaustion from any cause.

Mustard Seed.

R. Seminum Sinapeos albi, $\overline{3}$ j.

Signa—A tea-spoonful of the unbruised seeds two or three times a-day.

They are said to prove beneficial in costive habits, where the constipation depends upon a torpid state of the muscular coat of the intestines.—*Paris on Diet*, p. 187.

Infusion of Sassafras, &c.

R. Corticis Sassafragis, $\overline{3}$ ss.

Radici Glycyrrhizæ, $\overline{3}$ ij.

Aquæ bullientis, Oj.

Fiat infusio.—Signa.—One half may be taken in the morning, and the other half in the evening.

Recommended highly by the Germans in many diseases of a scrofulous character.—*Hufeland's Treatise*.

Mustard Whey.

R. Lactis vaccinae, ℥j.

Semen Sinapeos contus: ℥j.

Simmer until the caseous part separates, then strain, and add—

Vini albi, ℥vj.

Signa.—A draught of this whey may be taken every hour or two in low fever, and debilitated stomach.

Wine Whey.

R. Lactis vaccinae, Oss.

Vini albi, (Madeira,) ℥j vel ℥ij.

Boil the milk and then add the wine.

There are few mild stimulants more employed, or more useful than wine whey. Dr. Chapman speaks of it as having particularly a kindred action with carbonate of ammonia. The dose to be regulated by the circumstances of the case; from four ounces to ten during the day.

Infusion of Cayenne Pepper.

R. Pulveris Capsici annui, ℥iss.

Aquæ bullientis, Oss.

Fiat infusio.—**Signa.**—A table-spoonful for a dose in severe colic, or gout in the stomach.

For the external use of it, see Rubefacients.

Infusion of Cloves.

R. Caryophylli contusi, ℥j.

Aquæ bullientis, Oss.

Fiat infusio.—**Signa.**—To be employed in the same cases, and in similar doses as the former.

This preparation is also very useful as an adjunct to bark in intermittent fever.

An Infusion of Ginger,

Is also an excellent stomachic, and may be employed with great benefit in debility of the stomach and bowels.

Mixture with Wine, &c.

R. Vitellum Ovi j.

Olei Cinnamomi, gtt. xx.

Misce, et adde—

Vini albi, (Madeira,)

Aquæ Cinnamomi, aa. ℥ij.

—— destillatæ, ℥ij.

Sacchari albi, ℥ij.

Signa.—Three spoonsful for a dose. During convalescence from low fevers.

CLASS XI.

NARCOTICS.

NARCOTICS have been denominated *Sedatives*, from their power of diminishing action; *Anodynes*, from their capability of alleviating pain; and *Hypnotics*, or *Soporifics*, from their power of inducing sleep. A few words only will be necessary as general rules for their administration, and any peculiarities that individuals of the class may possess will be noticed under the appropriate prescriptions. It is well known they possess the power of fulfilling two indications; namely, of exciting and depressing the powers of the system.

To obtain the first of these results, it is necessary to begin with small doses, and repeat them frequently; while, in order to arrive at the second, a full dose must be given, and not resorted to again for a considerable length of time. We should also bear in mind that the system very soon becomes habituated to their action, and the doses must be continually enlarged if it be found necessary to maintain their impression. The difference between the susceptibility to their action when the body is in a morbid or healthy condition, should not be forgotten.

PILLS.

R. Pulveris Opii, gr. xij.

Saponis albi, gr. iij.

Misce et divide in pilulas, xij.—Of which one is the ordinary dose to produce sleep.

In equal weights of powdered and soft opium, the former is much the more powerful, the latter containing a considerable proportion of water. This should be recollected when we are directing them in the form of pills.

An opium pill that has been made long enough to become perfectly hard and dry, will be retained by an irritable stomach when a fresh one would be rejected.

Pills of Lupulin.

R. Lupulinæ puræ, ʒss.

Contunde in mortario, et divide in pilulas x.

Signa.—From two to four pills may be given for a dose.

This substance, we are told by Dr. Ives, contains all the active properties of the hop. It is slightly narcotic, and will prove serviceable in irritable states of the stomach.

The lupulin may be administered in powder, with two parts of white sugar, if it be deemed a preferable mode.

Pills of Acetate of Morphia.

R. Acetatis morphiæ, gr. iij.

Conservæ rosarum, q. s., ut fiant pilulæ, xij.

Signa.—One may be given for a dose, repeated *pro re nata*.

One grain of either of the salts of morphia, is usually estimated as equal to four of opium.

The Sulphate and Muriate of Morphia are prepared in the same way, and given in the same dose.

Pills of Lactucarium.

R. Lactucarii optimi, gr. xij.

Divide in pilulas, vj.

Signa.—One for a dose, to be repeated in two hours, if sleep be not procured.

The inspissated juice of the garden lettuce is said to act as a direct sedative, lessening the frequency of the pulse, diminishing the natural heat, and procuring sleep, without inducing the subsequent distressing effects which follow the use of opium.

The Thridace of the French appears to be prepared by a different process from that employed in the preparation of the British Lactucarium. The former is not so highly esteemed in this country as the latter.

Pills of Hyosciamus and Ipecacuanha.

R. Extracti Hyosciami nigri, gr. x.

Pulveris Ipecacuanhæ, gr. v.

Misce et divide in pilulas x.—Signa.—One every half hour until relief is procured.

These pills are recommended by Dr. Paris, in his work on diet, in cases of flatulence. They are designed to calm the irritability of the bowels.

Pills of Opium, Hyosciamus, &c.

R. Pulveris Opii, gr. iv.

Extracti Hyosciami,

———— Conii, āā, gr. xv.

Fiat massa in pilulas vj. dividenda.—One every night where an anodyne is required.

Pills of Extract of Hyosciamus.

R. Extracti Hyosciami nigri, ʒj.

Divide in pilulas x.—Of which one may be given every four hours, increasing the dose as required.

This medicine is employed as a substitute for opium, and is said not to constipate the bowels.

Pills of Extract of Stramonium.

R. Extracti Stramonii, ʒss.

Divide in pilulas xv.—Signa.—One three times a-day.

The powdered leaves and seeds are also employed in similar doses. *Vide Expectorants.*

Opiate Confection.

R. Confectionis Opii, ʒss.

To be given for a dose.—One grain of opium is contained in forty-three grains of the Edinburgh confection.

Pills of Extract of Conium.

R. Extracti Conii, ʒj.

Pulveris foliorum Conii, q. s.; ut fiat massa, et in pilulas xx dividenda.

One of these pills night and morning, may be taken at the commencement, gradually increasing them to two, three, four, or more, as the patient can bear them.

The efficacy of narcotic plants is influenced by soil, climate, exposure, and cultivation; and the extracts by the season when the plants are gathered, and the mode of preparing them. Physicians would do well therefore to make themselves acquainted with the quality of the article they prescribe, inasmuch as there may be a fatal difference between two parcels of different ages, modes of preparation, &c.—*Vide Journal of the Philadelphia College of Pharmacy.—On Narcotic Extracts, Vol. II.*

Pills of Camphor.

R. Gummi camphoræ, ʒss.

Gummi Arabici,

Alcoholis, āā. q. s.

Ut fiant pilulæ xv.—Dose.—A pill every two or three hours.

Pills of Opium, Digitalis, &c.

R. Pulveris Opii,

———— Digitalis, āā. gr. vj.

Conservæ rosarum, q. s., ut fiant pilulæ xij.

Signa.—One to be taken every four hours. In Asthma, &c.

Pills of Extract of Conium and Calomel.

R. Extracti Conii, ℥ij.

Hydrargyri chloridi mitis, gr. xv.

Misce.—Fiant pilulæ xv.—Signa.—One three times a-day. In pseudo-syphilitic affections.

Pills of Powdered Nux Vomica.

R. Pulveris Nucis vomicæ, ʒss.

Conservæ rosarum, q. s., ut fiant pilulæ x.

Signa.—One to be given two or three times a-day, closely watching the effects. In paralysis.

Pills of the Resin of Nux Vomica.

R. Extracti Nucis vomicæ resinosi, gr. x.

Divide in pilulas xv.—One pill twice a-day, gradually augmenting until tetanic effects are produced.

Pills of Strychnia.

R. Strychniæ, gr. ij.

Conservæ rosarum, q. s., ut fiant pilulæ xxiv.

Of which one pill may be given once or twice a-day, according to the idiosyncrasy of the patient.

This latter medicine is still more powerful than the former two: for their mode of preparation, manner of exhibition, and peculiar effects, see Majendie's Formulary, page 82. It is only necessary in this place to observe, that great nicety is requisite in dividing the mass into pills of equal strength.

Tincture of Strychnia.

R. Alcohol, ʒj.

Strychniæ, gr. iij.

Fiat tinctura.—This tincture to be given by drops, from six to twenty-four, in draughts or drinks.

Draught of Strychnia.

R. Strychniæ puræ, gr. j.

Sacchari albi, ℥ij.

Acidi acetici, gtt. ij.

Aquæ destillatæ, ℥ij.

Fiat solutio.—Signa.—A tea-spoonful morning and evening.

Strychnia forms with acids a variety of salts, which generally are very soluble, and for this reason much more active than the alkali uncombined. These may be resorted to when the system becomes habituated to the action of strychnia. The commencing dose of the salts, should not be greater than that of the alkaloid. It should be borne in mind that this substance will be increased by all acid drinks, such as lemonade, both in solubility and power.—*See Majendie's Formulary, Edit. 5, p. 12.*

MIXTURES, TINCTURES, &c.

Opiate Mixture.

R. Pulveris Opii, ℥ss.

Sacchari albi, ℥j.

Aquæ Cinnamomi, ℥vj.

Fiat mistura.—Signa.—A table-spoonful may be taken every two hours, so that the whole be consumed in twenty-four. In cases of tetanus and colica pictonum.

Hemlock Mixture.

R. Succi Conii spissati, ℥ss.

Syrupi Papaveris albi, ℥j.

Aquæ destillatæ, ℥vij.

Misce.—A table-spoonful three or four times a-day for a dose.

Tincture of Resin of Nux Vomica.

R. Extracti Nucis vomicæ exsiccati, gr. iij.

Spiritus vini rectificati, ℥j.

Fiat tinctura.—This tincture to be given in doses of fifteen or twenty drops, in any kind of drink.

This preparation, and the pills made of the resin of *nux vomica*, have been successfully used in cases of general and local paralysis. *Nux vomica* is also applied by friction in paralytic and other nervous diseases. The following is Majendie's formula for this purpose:

R. Tincturæ nucis vomicæ, ℥j.

Aquæ Ammoniæ (ammoniaque concentrée,) ℥ij.

Misce.—

Opiate Collyrium.

R. Pulveris Opii, gr. j.

Camphoræ, gr. v.

Mucilaginis Acaciæ, ℥j.

Misce, et fiat collyrium.

Anodyne Draught.

R. Tincturæ Opii, gtt. xv vel xxv.

Syrupi Papaveris, ℥ij.

Spiritus Cinnamomi, ℥j.

Aquæ destillatæ, ℥iss.

Misce.—Signa.—Anodyne draught. Laudanum is very disagreeable to some persons; combined as above, the taste may be disguised.

As the regulation of doses for children is, to the young practitioner, one of the most difficult parts of prescribing; and as laudanum is an article so universally employed in their diseases, we subjoin the following graduated table from Dr. Dorsey's Syllabus, 1817, page 36.

For a child at birth, or within a month after	$\frac{1}{2}$ to 1 drop.
Under a year old - - - -	$\frac{1}{2}$ to 2 or 3
From one to two years of age - - - -	1 to 5
From two to five - - - - -	2 to 8
From five to ten - - - - -	5 to 15
From ten to fifteen - - - - -	10 to 20
At fifteen years - - - - -	15 to 20
For an adult - - - - -	25 to 30

It is important in the employment of this remedy, that it should be a *saturated* tincture, and perfectly transparent. For these reasons it should be obtained from the apothecary in small quantities, as it is liable to deposit part of the opium; hence the liquid near the surface is weaker, and causes that in the bottom of the phial to be very strong, if it be shaken before using. Infants have been destroyed by the exhibition of *thick laudanum*. Physicians, therefore, should always examine the phial, if any be in the possession of a family, before trusting the nurse to give a dose to the child.

Solution of Acetate of Morphia.

R. Acetatis Morphiæ, gr. xvj.

Aquæ destillatæ, ℥vj.

Acidi acetici diluti, ℥ij.

Misce.—The dose is from ten to twenty drops.

Sirup of Acetate of Morphia.

R. Acetatis Morphiæ, gr. iv.

Syrupi simplicis, ℥bj.

Misce.—Signa.—Two tea-spoonsful may be taken for a dose, and repeated *pro re nata*.

The sirup of the sulphate of morphia may be made in the same proportions, and given in the same doses. The latter salt has one superiority over the former, namely, that the acetate when dissolved in water is decomposed, two salts resulting, the one soluble, the other not. From this objection the sulphate is exempt.

Majendie states that, by alternating these new alkaline remedies, we may prolong the effect without increasing the dose. If this observation be correct it is well worth remembering.—*Majendie's Formulary*, p. 24

Solution of Sulphate of Morphia.

R. Morphiæ sulphatis, gr. ij.

Aquæ destillatæ, ℥ij.

Fiat solutio.—Dose.—A tea-spoonful, repeated according to circumstances. Twice that quantity is nearly equivalent to a grain of opium or twenty-five or thirty drops of laudanum.

The muriate and nitrate of morphia possess no advantages over the sulphate and acetate.

Solution of Citate of Morphia.

R. Morphinæ puræ, gr. xvj.

Acidi citrici crystallizati, gr. viij.

Aquæ destillatæ, ℥j.

Tincturæ coccinellæ, q. s.

Ut fiat solutio.—Dose five to twenty drops, *pro re nata*. *Majendie*.

The French chemists have also discovered in opium a principle allied in its effects to morphia, and called *Codéine*. In dose of a single grain it proves highly anodyne, but distresses the stomach in larger doses. Its salts are given in somewhat smaller quantities.—*Majendie, 9vme, Ed. p. 80.*

A preparation of opium, known as *McMunn's Elixir*, has been for several years much in vogue in this country. The mode of preparation is a secret with the proprietor; but the medicine has certainly some advantages over all others of its class. For example, it rarely affects the head, and causes little or no constipation, but acts freely on the urinary organs and skin. It is given in the same dose as laudanum.

Tincture of Lupulin.

R. Lupulinæ puræ contusæ, ℥j.

Alcohol, ℥ij.

Digest for six days in a close vessel. Strain, filter, and add a sufficient quantity of alcohol to make ℥iij. of the tincture. Of this from ℥ss. to ℥ij. may be taken for a dose.—See *Pharmacologia*, art. *Humulus*.

Tincture of Hops.

R. Tincturæ Lupulorum, ℥j.

Signa.—Of this, a tea-spoonful may be taken as a narcotic in cases where opium is inadmissible.

Hops are also given in the form of infusion; for which see tonics. Enclosed in a bag, they have acquired some celebrity as a pillow. Their soporific powers, however, when thus employed, are very equivocal.

Camphor and Hoffman's Anodyne.

R. Aquæ camphoræ, ℥iv.

Liquor Hoffmanni, ℥ij.

Misce.—Dose.—A dessert-spoonful every hour or two hours. In nervous affections, and the sleeplessness of fever.

Infusion of Camphor.

R. Camphoræ gummi, ℥j.

Aquæ bullientis, ℥viij.

Pour the hot water on the gum, and allow it to stand until cold: it should be then kept in a well-stopped phial. Dose.—A table-spoonful every hour or two hours. In febrile pervigilium and nervous irritability.

Tincture of Opium and Oil of Turpentine.

R. Tincturæ Opii, ℥j.

Olei Terebinthinæ, ℥ss.

Misce.—Signa.—A tea-spoonful every hour, may be given in delirium tremens.

The turpentine may be added or omitted at the discretion of the physician, and the quantity of laudanum in some cases may be doubled. We have given two tea-spoonsful every hour until ℥ij. of laudanum were taken, with success. We do not believe in the doctrine, however, that opium in this disease may be administered almost *ad libitum* without danger. On the contrary, we are well assured that death has been caused by the excessive use of this narcotic in delirium tremens.

Denarcotized Laudanum.

R. Tincturæ Opii sine Narcotina.

This preparation, now common in all our shops, was proposed by Robiquet. It is said to relieve pain and induce sleep, without producing the ordinary sequelæ of opium or laudanum.

Black Drop.

R. Tincturæ Opii acetatæ, ℥i.

From six to eight drops of this preparation may be given to an adult for a dose. It is said to be three times the strength of laudanum, but has obtained celebrity from the fact, that it equally promotes the ends of laudanum without affecting the head and stomach. Where it cannot be had, it is stated by Dr. Chapman, that by giving the alcoholic tincture of opium in vinegar, we obtain nearly the same results.

Elixir Paregoric.

R. Tincturæ Opii camphoratæ.

A tea-spoonful of this may be given to an adult for a dose. It is principally employed as an expectorant.

Wine of Opium.

The dose of the Vinum Opii is the same as the spirituous tincture; though we believe it is seldom employed, except as an external application to chronic inflammation of the eyes.

Tincture of Digitalis.

R. Tincturæ Digitalis, ℥ij.

Of which ten drops may be given at first twice or three times a-day, increasing the quantity as the system becomes accustomed to it. The peculiar character of this remedy should induce caution in giving it.

For its particular applications and effects, see Expectorants and Diuretics.

Infusion and Tincture of Digitalis.

Vide Diuretics and Expectorants.

Infusion of Conium.

R. Foliorum Conii, ʒss.

Aquæ bullientis, Oj.

Fiat infusio et cola.—Signa.—To be applied as a wash to cancerous and scirrhus ulcers.

A cataplasm is also employed for the same purpose, made by mixing the powdered leaves with bread and water poultice.

Solution of the Extract of Belladonna.

R. Extracti Belladonnæ, ʒj.

Aquæ destillatæ, ʒiij.

Fiat solutio.—Signa.—To be applied to the eye in cases of cataract, for the purpose of dilating the pupil and exposing the lens.

The extract of belladonna is occasionally given in doses of a grain three or four times a-day.

Ointment of Belladonna.

R. Extracti Belladonnæ, ʒij.

Aquæ destillatæ, ʒij.

Adipis suilli, ʒij.

Fiat unguentum.—To be applied to the neck of the uterus in rigidity of that part, occasioned by disease, or previous to delivery. It is said to act on the same principle as it does in dilating the iris. The application to be made for some time by means of a syringe with a wide mouth.

M. Chaussier.

Infusion of Hyosciamus Niger.

R. Foliorum Hyosciami nigri, ʒss.

Aquæ bullientis, Oj.

Fiat infusio.—Employed as an external application to cancerous, scrofulous, and other indolent ulcers.

Decoction of Bitter Sweet.

R. Dulcamaræ stipitum, ʒj.

Aquæ destillatæ, Ojss.

Boil to a pint and strain.

Dr. Chapman directs a wine-glassful to be taken frequently, and gradually increased, until a pint is consumed daily, in chronic eruptions, &c. The same preparation is employed as a wash.

Decoction of Stramonium in Milk.

R. Foliorum Stramonii, ʒij.

Lactis recentis, Oiss.

Boil to a pint and strain. Apply to gouty, rheumatic, and other painful swellings.

Solution of Extract of Belladonna.

R. Extracti Belladonnæ, gr. iij.

Aquæ Cinnamomi, ʒj.

Fiat solutio.—Signa.—Two or three drops twice or thrice a-day, to a child under a year old, and an additional drop for every additional year.

Belladonna has been recently used in Germany as a preventive of scarlatina. The extract, if given for several days, produces a scarlet eruption, and affection of the fauces, analogous to those of scarlatina; and it is pretended that this factitious disease is sufficient to prevent the access of the real one.

Opium Enema.

R. Pulveris Opii, gr. ij.

Mucilaginis Acaciæ, ʒss.

Lactis tepifacti, ʒij.

Misce pro enemate.

Opium Enema.

R. Pulveris Opii, gr. iij vel iv.

Axungię Porcinæ, ʒj.

Melt them together with a gentle heat, and when of a proper warmth, inject the mixture with a small syringe.

Solution of Opium in Lime Water, &c.

R. Extracti Opii, gr. j.

Aquæ Calcis,

Olei Amygdalæ dulcis, āā. ʒiij.

Fiat mistura.—Recommended by Dr. Sibergundi of Dorsten, in the treatment of sore nipples. It is to be applied to the parts on dossils of lint.

Laudanum Enema.

R. Tincturæ Thebaicæ, ʒj.

Infusionis Lini, ʒij.

Fiat enema.

It is generally observed that three times the quantity of a remedy may be thrown into the rectum that may be taken into the stomach. As regards opium, however, we believe it to be too much, particularly in children. Indeed, it is an unsettled point in therapeutics, whether remedies act most powerfully upon the rectum or the stomach.

Suppository of Opium.

R. Pulveris Opii, gr. ij.

Saponis albi, gr. iv.

Misce bene.—Introduce into the rectum, when an enema cannot be administered.

Plaster of Opium, &c.

R. Opii puri,

Camphoræ, āā. ʒss.

Emplastri Lithargyri, ʒij.

Melt the plaster, and having previously powdered the other articles, mix them gradually, stirring it. For local pains.

Plaster of Opium.

R. Pulveris Opii,

——— Camphoræ,

——— Saponis albi, āā. ʒj.

Tincturæ Thebaicæ, q. s.—ut fiat emplastrum.

This plaster is recommended by Dr. Dewees, and is employed in the same case as the above. Another, and the most common plan, is to spread Theriaca Andromachi* on leather of the size required, and cover it with powdered opium, sprinkling a little laudanum to promote the adhesion of the opium.

* The original Theriaca Andromachi is composed of sixty-one ingredients, pulverized and rubbed up with honey into an electuary. A preparation bearing the same name, but used for the same purposes, is now made in a much less elaborate way. The word *Theriaca* is, at the present day, applied to any coarse sirup, as treacle or molasses.

CLASS XII.

ANTISPASMODICS.

ANTISPASMODICS are those medicines which allay spasm, and compose the irregular actions of the nervous system. Some of the most powerful of them are derived from the class of Narcotics.

Pills of Asafætida and Soap.

R. Gummi Asafætidæ, ʒj

Saponis albi, gr. x.

Aquæ fontis, q. s., ut fiant pilulæ xx.

Two or four of these pills may be taken for a dose and repeated *pro re nata*.

Pilulæ Asafætidæ cum aloe, et

————— cum Rheo,

May be given in the same doses as those made with soap only.

Opium Pills.

As under Narcotics.

Pills with Castor and Salt of Amber.

R. Castorei, ʒj.

Acidi Succinici, ʒss.

Extracti Gentianæ, q. s., ut fiant pilulæ xxiv.

Dose three pills night and morning.—In hysteria.

Calomel Pills.

As under Sialagogues.

Pills of Musk, Camphor, &c.

R. Moschi optimi, ℥ss.

Camphoræ, ʒj.

Gummi Ammoniæ, ʒij.

Opii purificati, gr. iv.

Misce, et fiant pilulæ singulæ, gr. iv.

Signa.—The dose may be four or five in the twenty-four hours. In nervous diseases.

M. A. Richard.

Bolus with Valerian and Rust of Iron.

R. Pulveris radices Valerianæ, ʒj.

Carbonatis præcipitatis Ferri, ℥ss.

Mucilaginis Acaciæ, q. s., ut fiat bolus.

One three times a-day.

Bolus with Musk and Camphor.

R. Camphoræ, gr. v.

Moschi, gr. xv.

Syrupi simplicis, q. s., ut fiat bolus.

Signa.—To be taken at once.

This is a strong dose, but can be readily modified by the practitioner.

To confine the aroma, musk should always be kept in phials, whether in the form of pills or otherwise.

Bolus with Musk and Carbonate of Ammonia.

R. Moschi,

Carbonatis Ammoniæ, āā. ℥ss.

Conservæ rosarum, q. s., ut fiat bolus.

To be taken every third hour.

Recommended in mortification accompanied with spasmodic action.

Musk Mixture.

R. Moschi optimi, ℥ij.

Sacchari albi, ℥j.

Pulveris gummi Arabici, ℥ij.

Aquæ destillatæ, ℥vj.

Fiat mistura.—Signa.—A table-spoonful to be given every hour or two hours.

Mixture with Musk, &c.

R. Misturæ Moschatæ, ℥vj.

Tincturæ Opii camphoratæ, ℥ss.

———— Velerianæ ammoniatæ, ℥j.

Misce.—Signa.—A tea-spoonful three or four times a-day. For children in pertussis.

Tincture of Asafætida, Castor, &c.

R. Tincturæ Asafætidæ,

———— Castorei, āā. ℥ss.

Aquæ Ammoniacæ, ℥ij.

Misce.—A tea-spoonful may be taken in a glass of water for a dose.

Hoffman's Anodyne and Laudanum.

R. Liquoris Anodynæ Hoffmāni, ℥iij.

Tincturæ Thebaicæ, gtt. lxxx.

Aquæ Cinnamomi, ℥vj.

Fiat mistura.—Signa.—A table-spoonful every one or two hours. In hysteria, &c.

Mixture with Asafætida, &c.

R. Asafætidæ, ℥j.

Aquæ Menthæ piperitæ, ℥iss.

Fiat solutio, et adde—

Tincturæ Valerianæ ammoniatæ, ℥ij.

———— Castorei, ℥iij.

Ætheris sulphurici, ℥j.

Signa.—A table-spoonful for a dose every second hour. In hysteria.

Infusion of Valerian.

R. Valerianæ radiceis, ℥j.

Aquæ bullientis, Oj.

Put them into a covered vessel and let them stand until cold.—Dose a wine-glass full every two or three hours.

Tincture of Valerian and Hoffman's Anodyne.

R. Liquoris Hoffmani Anodynæ,

Tincturæ Valerianæ, āā. ℥i.

Misce.—Signa.—A tea-spoonful for a dose, *pro re nata*, shaking the phial each time.

Mixture with Ether and Laudanum.

R. Ætheris Sulphurici, ℥ij.

Sacchari albi,

Gummi Acaciæ, āā. ℥iss.

Tincturæ Thebaicæ, gtt. lx.

Aquæ Cinnamomi, ℥ij.

Misce.—Signa.—A tea-spoonful every hour. In cramp of the stomach, &c.

Solution of Boracic Acid, &c.

R. Syrupi simplicis, ℥j.

Acidi boracici, ℥j.

Infusionis Tiliæ vel fœniculi, ℥iv.

Misce.—Signa.—A small table-spoonful three or four times a-day.

The above is employed by the French physicians in cerebral affections. *Ratier.*

Asafœtidæ Enema.

R. Asafœtidæ, ℥ij.

Decocti Avenæ, ℥x. vel ℥xij.

Misce pro enemate.—Tinc. Opii may be added *pro re nata*. In hysteria, spasmodic colic, &c.

Musk Clyster.

R. Moschi, gr. xij.

Sacchari purificati, ℥ij.

Gummi Acaciæ contriti, ℥jss.

Alcoholis ammoniati, gtt. xxx.

Infusionis Lini, ℥iv.

Fiat enema.—For children seized with convulsions.

Confection of Opium, &c.

R. Confectionis Opii, 3vj.

Tincturæ Camphoræ, 3x.

Aceti destillati, ℥ij.

Misce.—To be applied as a plaster warm to the region of the stomach.

In vomitings, &c.

Opiate Plaster.

As under Narcotics.

Hemlock Plaster.

As under Narcotics.

Tobacco Clyster.

As under Cathartics.

CLASS XIII.

SIALAGOGUES.

THOSE medicines which increase the salivary secretion are denominated Sialagogues. In forming these into a *class* in this place, we trespass somewhat upon the arrangement of Dr. Chapman, but merely from a motive of convenience; for we agree with him that such a class is better expunged from the *Materia Medica*, at the same time that it appeared necessary to introduce it here, in order to place the mercurial preparations in stronger relief than could otherwise be done.

Pills of Calomel.

R. Hydrargyri Chloridi mitis, gr. xij.

Conservæ rosarum, q. s., fiant pilulæ xij.

Signa.—One morning and evening, and if necessary until salivation is excited.

Pills of Corrosive Sublimate.

R. Chloridi corrosivi Hydrargyri,

——— Ammoniæ, āā. gr. v.

Aquæ destillatæ, gtt. xxx vel xl.

Conservæ rosarum, ʒj.

Pulveris Glycyrrhizæ, q. s., ut fiant pilulæ xl.

Of which, give one pill three or four times a-day. In lues venerea.

Dupuytren's Antisyphilitic Pills.

R. Deuto-chloridi hydrargyri, gr. ss.

Extracti cinchonæ, gr. x.

——— Opii, gr. ss.

Pulveris cinchonæ, q. s.

Ut fiant pilulæ ij.—The two pills may be taken, one in the morning and the other at night; but as each contains a quarter of a grain of sublimate, their action must be closely watched. M. Dupuytren was partial to these pills in the treatment of Syphilis. He thought the opium and cinchona promoted and beneficially modified the action of the sublimate.

Pills with Corrosive Sublimate and Hemlock.

R. Chloridi corrosivi Hydrargyri, gr. vj

Solvantur in—

Aquâ destillatâ, q. s. et adde—

Succi Conii spissati, ℥j. cum

Pulvere Conii, q. s., ut fiat massa, in pilulas xlviij dividenda.

These pills contain the same quantity of mercury as the former, and may be given in the same number, and for the same purpose.

Pills of Acetate of Mercury.

R. Acetatis Hydrargyri,

Mannæ optimæ,

Pulveris Acaciæ, āā. ʒj.

Aquæ rosarum, q. s., ut fiant pilulæ, xx.

Of which let three be taken every night, till a gentle salivation is excited.

This preparation is said to be the mildest of the mercurial salts, and is less apt to disturb the bowels; but it is believed to be less efficacious in arresting the venereal disease.

The effect of the mercurial preparations upon children is sometimes very serious, and even fatal. So powerful upon them is the action of this medicine, that a profuse, or even gentle salivation, will sometimes

produce mortification and destruction to the jaw, cheek, and lip. This, however, does not often happen, but is such a shocking occurrence, that the gums should be frequently and carefully examined whilst they are taking mercury. It is, nevertheless, a fact, that a disease precisely similar in all its features occasionally attacks children who are badly clothed and fed, though no mercury has been given to them.

Solution of Cyanide of Mercury.

R. Hydrargyri cyanidi, gr. viij.

Aquæ destillatæ, ℥j.

Fiat solutio.—This is the *Liqueur Anti-Syphilitique* of Chaussier, who prefers it to corrosive sublimate on account of its greater solubility and more speedy action. By giving a tea-spoonful of the solution we give one-eighth of a grain of the cyanide, which may be repeated morning, noon and night, watching its effects.

Pills of Iodide of Mercury.

R. Iodidi hydrargyri, gr. v.

Confectionis rosæ, q. s.

Ut fiant pilulæ, xxx.

Each pill will contain the sixth of a grain, which may be given three times a-day. Double the quantity (two pills) is sometimes administered at first, but care should be taken not to push it too far.

Compound Pills of Proto-Iodide of Mercury.

R. Proto-iodidi Hydrargyri, gr. iij.

Extracti Opii, gr. ij.

Lactucarii, gr. xij.

Extracti guiaci, gr. xxiv.

Misce, et fiant pilulæ, xxiv.—Given by M. Ricord in the syphilis of children. He directs one pill for a child six months old, and two, three, or four pills for older children.

Dunghlison, New Rem. p. 280.

Pills of Proto-Iodide of Mercury.

R. Proto-iodidi Hydrargyri, gr. j.

Extracti Juniperi, gr. xij.

Pulveris Glycyrrhizæ, q. s.

Ut fiant pilulæ, viij.—In adult patients Majendie directs two to be given morning and evening, to be increased to three or four.

Dunghlison, ut supra.

Pills with Calomel, Camphor, &c.

R. Hydrargyri chloridi mitis,

Camphoræ, aa. ʒj.

Pulveris Opii, gr. xij.

Syrupi simplicis, q. s.—ut fiant pilulæ xx.

Signa.—One or two to be taken morning, noon and night. In cases where calomel by itself causes purging.

Blue Mercurial Pills.

R. Massæ ex Hydrargyro, ʒiss.

Divide in pilulas, xij.

Signa.—One may be taken three or four times a-day, according to circumstances.

If they should produce any effect upon the bowels, not desired, Pulv. Opii. gr. ss. may be added. The mass directed in the American Pharmacopœia contains one grain of mercury in every two and a half grains. One grain is contained in four of the Edinburgh—and in three of the London and Dublin.

Powder of Mercury and Chalk with Ipecacuanha.

R. Hydrargyri cum creta, ʒj.

Pulveris Ipecacuanhæ, ʒss.

Misce.—Fiant chartulæ, vj.

Signa.—One night and morning in sirup.

Recommended by Dr. Paris, in those cases of dyspepsia in which the biliary secretion is impaired. He prefers it under these circumstances to every other form of mercury.

Compound Powder of Sulphate of Mercury.

R. Hydrargyri flavi sulphatis, gr. j.

Pulveris Asari, ʒiss.

Misce et divide in chartulas, viij.

Signa.—One of these may be used as an Errhine, morning and evening. In Gutta Serena, Ozæna, and after the removal of Polypi.

Pills of the Red Oxide of Mercury.

R. Oxidi Hydrargyri rubri, gr. iij.

Pulveris Opii, gr. j.

Olei Caryophilli, gtt. ij.

Fiant pilulæ iij.—Signa.—One pill every night for a week. In venereal cases.

This preparation was formerly used by the celebrated J. Hunter; but from its harshness it is now generally superseded by others.

Mixture with Muriate of Mercury.

R. Chloridi Hydrargyri corrosivi, gr. ij.

Aquæ destillatæ, ℥vj.

Spiritus Cinnamomi,

Syrupi simplicis, aa. ℥j.

Misce.—Dose.—One or two large spoonsful, twice or thrice a-day. In venereal cases.

Mercurial Solution.

R. Chloridi Hydrargyri corrosivi, gr. xvj.

Alcoholis, ℥ss.

Aquæ destillatæ, Oj.

Misce.—This is the celebrated *Liquor of Van Swieten*, who directs half an ounce of it to be taken morning and evening, in milk, solution of gum or sirup. Its use requires caution. In venereal affections.

Spirituuous Solution of Corrosive Sublimate.

R. Chloridi corrosivi Hydrargyri, gr. ij.

Spiritus vini tenuoris, ℥iv.

Solve.—Of which a table-spoonful, night and morning, may be taken in a draught of decoction of Sarsaparilla.

The muriate of mercury is decidedly preferred by some physicians in the treatment of syphilis. John Hunter was of opinion, however, that the disease was more liable to recur, after its use, than when some of the other preparations had been employed.

Strong Mercurial Ointment.

R. Unguenti Hydrargyri fortis, ℥j.

One drachm of this ointment to be rubbed on the inside of the thighs, before the fire, every night.

It is by some physicians recommended to employ the warm bath before using this ointment, while others consider it useless. We think the practice may occasionally be beneficial. The hands of the person rubbing it on, if it be an attendant, should be enveloped in a bladder, or gloves made of oiled silk.

Strong Mercurial Ointment with Camphor.

R. Unguenti Hydrargyri fortis, ℥j.

Camphoræ, ℥j.

Misce.—To be employed as the former.

Masticatory with Pellitory, &c.

R. Pulveris Radicis Pyrethri,

Resinæ Mastichi, āā. ℥j.

Let these be dissolved together by a moderate application of heat: then divide the mass into two equal parts. One of these lumps may be chewed at pleasure. In toothach and paralytic affections of the tongue.

Mixture of Pellitory, Opium, &c.

R. Radicis Pyrethri contusæ, ℥ss.

Aceti destillati, ℥vj.

Gummi Opii, gr. iij.

Digest in a sand bath for an hour.—Signa.—A spoonful to be held in the mouth and frequently renewed. Used in the same cases as the former.

There are several other substances, such as polygala senega, nitric acid, &c., which occasionally excite

salivation; but as this is simply one effect of these remedies, and one of little consequence, they are treated of under more appropriate heads.

Mercurial Lotion.

R. Chloridi Hydrargyri corrosivi, ʒj.

Aquæ destillatæ, ʒvj.

Solve et adde—

Muriatis ammoniæ, ʒij.

Potassæ nitratis, ʒss.

Fiat lotio.—Principally used as a wash in itch.

Good's Study of Med. vol. iv.

Antacid Tincture.

R. Guaiaci pulveris, ʒj.

Balsami Canadensis, ʒj.

Hydrargyri chloridi corrosivi, ʒj.

Olei Sassafragis, ʒij.

Spiritus Vini Rectificati, ʒviij.

Dissolve the Corrosive Sublimate in one-half the spirit, and then add to it the other ingredients.—*Signa.*—Of this from ten to twenty drops may be taken morning and evening, in wine or water.

This form of exhibiting corrosive sublimate is of ancient date; and has been lately employed by Dr. Emerson and other practitioners of this place in lues venerea, &c.

Antacid Ointment.

R. Hydrargyri Oxidi rubri, ʒss.

———— Chloridi corrosivi, ʒij.

Acetatis Plumbi, ʒj.

Ceræ Albæ, ʒiv.

Olei olivarum, ʒvj.

Olei Bergamotta gtt. xx.

Rub the two mercurial preparations with water to an impalpable powder—then dry them, and add the acetate of lead. Melt the oil and wax together, and add the other ingredients gradually—stirring the mixture.

This ointment has some celebrity in the cure of tetter, and some other obstinate cutaneous diseases.

Ointment of Red Precipitate, &c.

R. Oxidi Hydrargyri rubri, gr. x.
Sulphatis Zinci, ʒj.
Axungiae purificatae, ʒij.
Fiat unguentum.

Recommended by Professor Dupuytren in obstinate chronic ophthalmia depending on a scrofulous diathesis.

Anti-herpetic Ointment.

R. Sulphatis Hydrargyri flavæ, ʒj.
Tincturæ Opii, ʒj.
Sulphuris sublimati, ʒss.
Axungiae, ʒj.
Fiat unguentum.—This has been successfully used in herpes, alternated with the simple water bath by *M. Cullerier*.

Ointment of Deuto-ioduret of Mercury.

R. Deuto-iodureti Hydrargyri, gr. xv.
Adipis preparatae, ʒij.
Olei limonis, gtt. xx.
Fiat unguentum.

This ointment has been found highly efficacious in syphilitic tubercles, or chronic ulceration of a syphilitic character, but its great activity requires close attention. The following formula, which is much less active, may be employed with less inconvenience.

R. Proto-iodureti Hydrargyri, ʒss.
Adipis præparatae, ʒiss.
Olei, gtt. xv.

Fiat unguentum.—Syphilitic ulcerations which have resisted a variety of remedies, have cicatrized rapidly under the use of this ointment.

Ratier.

Ointment of Cyanuret of Mercury.

R. Cyanureti Hydrargyri, gr. xvj.

Adipis præparatæ, ʒj.

Essentiæ citri medicæ, gtt. xv.

Fiat unguentum.—Used in Herpes, attended by intense inflammation and itching.

CLASS XIV.

T O N I C S.

THOSE medicines are considered Tonics which impart strength to the system, without subjecting it to any preternatural excitement during their operation. They stimulate, however, in a secondary manner, by gradually increasing the force of the circulation, and by restoring the digestive and secretory functions to a more healthy state. Considerable perseverance is often necessary in giving any one of this class a fair trial; and when one fails, we may often resort with success to another: much also may sometimes be gained by alternating them.

POWDERS, &c.

Powder of Peruvian Bark.

R. Pulveris Cinchonæ optimæ.

A drachm or a tea-spoonful to be taken every hour in port-wine, water, or milk, during the intermission of fever and ague.

It is directed by some practitioners to suspend the bark till within a few hours of the expected paroxysm; and by a few others to administer it during the hot stage of the disease. We have generally found it to be the most successful plan to commence immediately after the paroxysm has ended, and give every hour a

dose of the remedy, until the expected period of its return; and if the chill does not recur, to continue it in smaller quantities for several days.

Powder of Peruvian Bark and Cloves.

R. Pulveris Cinchonæ,
Super-tartratis Potassæ, āā, ʒj.
Pulveris Caryophilli, ʒj.

Misce.—Dose.—A drachm and a half every second hour.

This compound will sometimes succeed when the bark alone fails.

Powder of Peruvian Bark and Sulphate of Magnesia.

R. Pulveris Cinchonæ,
Sulphatis Magnesiae, āā. ʒvj.

Fiat pulvis, et divide in partes æquales iv.—One to be taken every two hours, where purging is required.

Powder of Peruvian Bark, Snake Root, &c.

R. Pulveris Cinchonæ, ʒss.
——— Serpentariae, ʒj.
Sodæ Bi-carbonatis, gr. xl.

Misce.—Divide in chartulas iv.—Signa.—One to be given every two hours. In obstinate intermittents.

If the bark should purge when given alone, a little opium may be added: when it oppresses the stomach, cloves, as in a former prescription. If costiveness should follow, rhubarb, or sulphate of magnesia may be used. The alkali in the above preparation will correct acidity.

Powder of Augustura Bark.

R. Pulveris Cuspariæ febrifugæ, ʒss.

Divide in chartulas vj.—Of which one may be taken three or four times a-day, in water or any other vehicle.

This remedy, we learn, has been successfully employed in uterine hemorrhage.

Powder of carbonate of Iron.

R. Pulveris carbonatis Ferri præcipitati, ʒj.

Divide in chartulas iv.—Signa.—One every six hours in molasses. In Chorea, Neuralgia, &c.

The above remedy, we are told, has been successfully used in these very painful affections, both in England and America. The dose is sometimes carried as far as ʒss. every four or six hours, after antiphlogistic treatment, if the latter be required.

Powder of Colomba, Ginger, &c.

R. Pulveris Colombæ,

———— Carbonatis Ferri præcipitati,

———— Rhei,

———— Zingiberis, āā. ʒj.

Misce.—Fiant doses xij.—One to be taken three times a-day, in molasses.

Powder of Colomba and Tartrate of Iron.

R. Tartratis Ferri, ʒij.

Pulveris Colombæ, ʒj.

Fiant pulveres iv.—Signa.—One every three or four hours, in sirup. }

Powder with Rust of Iron, &c.

R. Sulphatis Ferri, ʒij.

Carbonatis Ferri præcipitatis, ʒiss.

Divide in pulveres xij.—Signa.—One to be taken morning, noon and night, in sirup or molasses.

Pills of Cinchona and Camphor.

R. Extracti cinchonæ, ʒj.

———— Opii, gr. j.

Gummi camphoræ, gr. xij.

Pulveris cinchonæ, q. s.

Ut fiant filulæ xij.—One to be taken three or four times a-day.

Ratier.

Powder of Prussiate of Iron and Guaiacum.

R. Ferrocyanatis Ferri,

Pulveris Guaiaci, āā. ʒj.

Misce et divide in chartulas xij.—Signa.—One three times a-day.

This prescription has been successfully employed in obstinate intermittents.

Pills of Extract of Bark, &c.

R. Extracti Cinchonæ,

————— Gentianæ, āā. ʒj.

Sulphatis Ferri, ʒss.

Pulveris Myrrhæ, ʒj.

Olei Carui, gtt. x.

Syrupi Zingiberis, q. s., ut fiant pilulæ lx.

Signa.—Three to be taken three times a-day.

Pills of Sulphate of Quinia, &c.

R. Sulphatis Quiniæ, gr. x.

Conservæ Rosarum, q. s., ut fiant pilulæ x.

Signa.—One to be given every hour in the apyrexia of intermittent fever.

There is also another principle obtained from the cinchona lancifolia, or pale bark, which possesses very nearly the same powers as the first. This is called *Cinchonia*. These preparations produce all the effects of the Peruvian bark, and have now almost entirely superseded it. The sulphate of quinia affects the chest of the consumptive patient with tightness in an equal degree with the bark; and we have several times observed it to produce a soreness of the mouth exactly resembling that resulting from mercury. The observations of some of our friends have confirmed this fact.

A substance has lately been discovered in the bark of the willow, (*Salix helix*,) which is proved not to be alkaline, but which answers all the purposes of quinia in the cure of intermittent fevers. It is called salicine,

and has been used both in France and Italy.—*Vide Journal of the Philadelphia College of Pharmacy, Vol. II. No. 3.*

Pills of Salicine.

R. Salicinæ, gr. xxiv.

Mucilaginis, q. s.

Ut fiant pilulæ viij.—A pill to be taken three or four times a-day.

Salicine is obtained from willow bark (*salix helix*.) It has been chiefly used in intermittent fever, and produces effects very similar to those of Quinia.

Pills of Ammoniuret of Copper.

R. Cupri ammoniati, ʒj.

Conservæ rosarum, q. s., ut fiant pilulæ xxx.

Signa.—One or two to be taken at bed-time. In Epilepsy.

Pills of Sulphate of Quinia and Carbonate of Ammonia.

R. Sulphatis Quiniæ, gr. xij.

Carbonatis Ammoniæ, gr. xxiv.

Misce.—Fiant pilulæ vj.—Signa.—One every hour, beginning six hours before the expected paroxysm.

These pills are large, but we have succeeded with them when quinia alone and in smaller doses had proved unavailing. They may also be combined advantageously with the oil of black pepper, or other essential oils, in cases of irritable stomach.

Febrifuge Boluses.

R. Pulveris Cinchonæ, ʒj.

Muriatis Ammoniæ,

Carbonatis Ammoniæ, āā. gr. xij.

Tartratis Antimonii, gr. xvij.

Syrupi simplicis, q. s., ut fiat massa et divide in bolos xlvij.

Signa.—One to be taken four times a-day, during the apyrexia of fever.

Hotel Dieu.

Pills of Piperina

R. Piperinæ puræ, gr. xij.

Extracti Gentianæ, q. s., ut fiant pilulæ xij.

Signa.—One every hour during the apyrexia of fever.

This is one of the new remedies lately introduced into practice. It is prepared from the black pepper, and is said to possess considerable febrifuge powers. It sometimes vomits, purges, or produces diaphoresis; and therefore requires caution in its administration. It may be given in mixture, if preferred.

Pills of Arsenic, &c.

R. Acidi arseniosi, gr. ij.

Pulveris Opii, gr. iij.

Saponis venetæ, gr. viij.

Pulveris Glycyrrhizæ, q. s., ut fiat massa in pilulas xx dividenda.

Signa.—One four times a-day, increased gradually according to circumstances. In intermittent fever and local diseases of an intermittent character.

Dr. Meigs.

Pills with the Sulphate of Copper.

R. Sulphatis Cupri, gr. iv.

Extracti Cinchonæ, gr. xxxij.

Syrupi simplicis, q. s., ut fiat massa, in pilulas xvj dividenda.

One to be taken four times a-day. In obstinate intermittents.

Pills of the Oxide of Bismuth.

R. Sub-nitratis vel oxidi Bismuthi, ʒj.

Mucilaginis gummi Acaciæ, q. s., ut fiant pilulæ xxx.

Signa.—One to be taken every two hours. In dyspepsia.

Pills of Muriate of Ammonia, Arsenic, &c.

R. Muriatis Ammonia, ʒss.

Pulveris Opii, gr. viij.

Acidi arseniosi, gr. vj.

Syrupi simplicis, q. s., ut fiant pilulæ xxxij.

Of which one may be given three times a-day. In intermittent fever.

These pills are strongly recommended by some practitioners. Great care is requisite in preparing them.

Pills of Ammoniated Iron, &c.

R. Ammoniaci Ferri, ℥j.

Pulveris Rhei, gr. viij.

Conservæ rosarum, q. s., ut fiat massa in pilulas v dividenda.

This remedy is recommended in Rachitis, and is to be continued with one pill a-day for a fortnight, and then intermitted for the same length of time and again resumed. The intention of this direction appears to be to prevent the system from becoming habituated to the medicine.

Pills of Oxide of Zinc.

R. Oxidi Zinci, ℥ij.

Conservæ rosarum, q. s., ut fiant pilulæ x.

Signa.—One three or four times a-day. In epilepsy, chorea, &c.

Dr. Chapman speaks favourably of this medicine, and recommends to begin with the above dose, and gradually to increase it.

Pills of Steel Filings.

R. Limaturæ Ferri, gr. xvj.

Extracti Gentianæ, q. s., ut fiat massa.—Divide in pilulas vj.

Signa.—One three times a-day, followed by a draught of some bitter infusion.

Pills of the Extract of Quassia.

R. Extracti Quassiae mollis, ℥j.

Pulvaris Gentianæ, q. s., ut fiant pilulæ xx.

Of which one may be taken three or four times a-day, succeeded by a wine-glassful of the infusion of quassia.

This is one of the purest bitters, and where we wish the effect of this class only, we cannot select a more appropriate preparation.

Pills of Nitrate of Silver, &c.

R. Nitratis Argenti, gr. iij.

Pulveris Opii, ʒss.

——— Camphoræ,

——— Nucis moschatæ, aa. ʒj.

Mucilaginis Acaciæ, q. s., ut fiat massa.—Divide in pilulas xlv.

Dose.—One morning and evening. In chorea and epilepsy.

Pills of the Sulphate of Iron.

R. Sulphatis Ferri, ʒj.

Extracti Gentianæ, q. s., ut fiat massa.—Divide in pilulas, xxx.

One to be taken morning, noon and night. In dyspepsia.

Pills of Nitrate of Silver.

R. Nitratis Argenti, gr. iij.

Micæ panis, ʒss.

Aquæ fontanæ, q. s., ut fiant pilulæ xij.

Signa.—One to be taken night and morning, gradually increasing them as the case may demand. The nitrate of silver should first be dissolved in a little water, and the crumbs of bread afterwards added; and a glass mortar should always be employed to compound them in.

Electuary of Cinchona, Rust of Iron, &c.

R. Pulveris Cinchonæ optimæ,

——— Carbonatis Ferri, aa. ʒj.

Balsami Copaivæ, q. s., ut fiat electuarium.

To be made (by the patient) into pills of the ordinary size. Of which four may be taken three times a-day.

Pills of the Nitro-Muriate of Gold.

R. Nitro-muriatis Auri, gr. v.

Pulveris Glycyrrhizæ, ʒiss.

Mucilaginis Acaciæ, q. s., ut fiat massa, in pilulas lxxv. dividenda.

Signa.—One or two thrice a-day. In syphilis, &c.

This metal is placed under the head of tonics by Dr. Chapman, and we have followed his example. Although once much extolled, it is now seldom used.

Pills of Ammoniuret of Copper and Extract of Quassia.

R. Cupri Ammoniaci, ʒss.

Extracti Quassiae mollis, ʒiss.

Misce, et divide in pilulas xxx.—Of which one may be taken three times a-day. In nervous diseases.

MIXTURES, &c.

Griffith's Myrrh Mixture.

R. Gummi Myrrhæ, ʒj.

Sulphatis Ferri, ʒj.

Carbonatis Potassæ, ʒj.

Sacchari Purificati, ʒij.

Aquæ destillatæ, ʒvj.

Fiat mistura.—Dose.—A table-spoonful according to circumstances. Employed as a tonic in phthisis.

Mixture of Sulphuric Acid, &c.

R. Sulphatis Ferri, gr. ij.

Acidi sulphurici, gtt. x.

Sacchari albi, ʒj.

Aquæ destillatæ, ʒj.

Misce.—Signa.—A tea-spoonful for a dose, to be repeated every two, three, or four hours. Designed for children.

Vinous Infusion of Carbonate of Iron, &c.

R. Carbonatis Ferri Præcipitatis, ʒiss.

Pulveris radices Gentianæ,

Corticis Aurantii contusæ, aa. ʒss.

Vini Lusitanici rubri, Oij.

Macerate for two or three days, and give a wine-glassful two or three times a-day. In dyspepsia arising from intemperance.

Mixture with Tincture of Bark and Citrate of Potash.

R. Succo Limonis recentis, ʒjss.

Carbonatis Potassæ, ʒj.

Tincturæ Cinchonæ, ʒj.

Aquæ Cinnamonii, ʒiij.

Misce.—Signa.—A table-spoonful every two hours. In cases in which it is desirable to promote insensible perspiration whilst taking the bark.

Infusion of Sage and Boneset.

R. Salivæ officinalis,
Eupatorii perfoliati, āā. ʒss.
Corticis Cascarillæ, ʒj.

Misce.—Infuse in three half-pints of water until cold. Dose.—A wine-glassful every three or four hours. In Hectic Fever

Decoction of Cinchona.

R. Corticis Cinchonæ contusæ, ʒj.
Aquæ frigidæ, Oj.

Boil for ten minutes—and while hot add—

Surpentariæ radicis, ʒss.

Corticis Aurantii contusæ, ʒij.

And suffer them to infuse for half an hour near the fire in a covered vessel.—Dose.—A wine-glassful every hour.

If the bark be boiled longer than ten minutes, it deposits extractive and resinous matter, becomes paler, and is less efficient. The last ingredients are directed to be infused because their aromatic properties would be dissipated by decoction.

Decoction of Bark, &c.

R. Decocti Cinchonæ, ʒvj.
Tincturæ Cinchonæ Huxhami, ʒj.
Acidi sulphurici aromatici, gtt. xl.

Misce.—A table-spoonful may be taken every one or two hours *pro re nata*.

By this preparation we obtain the extract, both aqueous and alcoholic, of the bark.

Sulphate of Quinia in Sirup.

R. Sulphatis Quiniæ, gr. xv.
Acidi sulphurici, gtt. vj.
Sirupi Zingibe vel Limonis, ʒij.

Misce.—Signa.—A large tea-spoonful may be given every hour during the apyrexia.

The French physicians employ a wine and tincture of quinia, which, however, are not superior to the compound used in this country. Both in Europe and America, and especially in Italy, the dose is carried to a much greater amount than we have directed. Eight, ten, fifteen, twenty, and thirty grains have been given by the Italians without danger and with success. Such doses many persons cannot bear, and generally the cure can be accomplished with sixteen grains in twenty-four hours.

Mixture of Sulphate of Quinia.

R. Sulphatis Quiniæ, gr. xvi.

Acidi sulphurici, gtt. v.

Sacchari albi, 3j.

Aquæ Cinnamomi, ʒijss.

Misce.—Signa.—A tea-spoonful for a dose, every hour during the apyrexia of intermittents.

By the addition of sulphuric acid, a super-sulphate of quinia is formed, which is soluble in the aqueous menstruum.

Tincture of Quinia.

R. Quininæ, ʒj.

Alcohol, ʒss.

Fiat tinctura.—Dose.—Ten to twenty drops every hour.

Ferrocyanate of Quinia.

R. Ferrocyanitis quiniæ, gr. iv.

Spiritus rectificati, ʒj.

Solve et adde—

Aquæ camphoræ, ʒvij.

Misce.—This preparation is said to be more active than the sulphate, whence it should be given in less doses: a tea-spoonful of the solution (which will contain half a grain of the salt) may be taken every

hour or two hours, according to circumstances.* Pills of the Ferrocyanate of Quinia are prepared with simple mucilage, and administered like the sulphate.

The acetate, citrate, muriate and nitrate of Quinia, have also been employed in Europe, and occasionally in this country. They possess no advantages over the ordinary preparations, and are given in much the same quantity.

Sirup of Cinchonine.

R. Sulphatis Cinchoninæ, gr. xxiv.

Syrupi simplicis, Oss.

Fiat mistura.—A table-spoonful will contain nearly a grain and a-half of the salt, which is a full dose; to be repeated *pro re nata*.

Cinchonine is also and more readily administered in pills of a grain or two grains each. In intermittent fevers.

Mixture of Red Bark in Port Wine, &c.

R. Pulveris Cinchonæ oblongifoliæ, ℥ss.

Theriaca Andromachi, ℥j.

Succi limonis, ℥ij.

Vini Lusitanicæ, ℥iv.

Misce.—Signa.—A wine-glassful every second hour during the apyrexia.

We find this prescription strongly recommended in the *N. A. Medical and Surgical Journal*, by Dr. Meigs, in intermittent fever.

Mixture with Decoction and Tincture of Bark, &c.

R. Decocti Cinchonæ, ℥ij.

Tincturæ Huxhami, ℥j.

Pulveris Cinchonæ, ℥ij.

Syrupi simplicis, ℥ss.

Fiat mistura.—A table-spoonful to be taken every one or two hours during the absence of fever.

* Vide Dunglison, *New Remedies*, p. 408

Mixture with Extract of Bark, &c.

R. Extracti Cinchonæ, ℥j.

Decocti Cinchonæ, ℥vj.

Tincturæ Cardamomi, ℥iv.

Misce.—A table-spoonful for a dose, as above directed.

Tincture of Cinchona, Valerian, &c.

R. Tincturæ Cinchonæ,

———— Valerianæ, aa. ℥j.

———— Cardamomi, ℥ij.

Aquæ menthæ, ℥iv.

Fiat mistura.—Of this mixture a table-spoonful may be given every third hour. This preparation is especially designed as a tonic in nervous temperaments.

Mixture of Fowler's Solution, &c.

R. Liquoris potassæ arsenitis, gtt. lx.

Tincturæ Opii, gtt. xxx.

Spiritus Lavendulæ compositi, ℥j.

Aquæ Cinnamomi, ℥iij.

Fiat mistura.—Signa.—A table-spoonful for an adult, a tea-spoonful for a child; to be administered every two hours, during the apyrexia of intermittents.

Arsenical Mixture.

R. Liquoris arsenicalis, gtt. x.

Aquæ destillatæ, ℥j.

Tincturæ Opii, gtt. x.

Spiritus lavendulæ compositi, ℥ss.

To be taken at a dose.

This mixture is recommended by Dr. Graves, when the simple arsenical solution irritates the stomach.

Acidulated Tincture of Gentian.

R. Tincturæ Gentianæ compositæ, ℥j.

Acidi sulphurici aromatici, ℥j.

Misce.—A small tea-spoonful may be taken in sugar and water, three or four times a-day. In dyspepsia.

Tincture of Bark.

R. Tincturæ Cinchonæ compositæ, ℥j.

Signa.—A tea-spoonful for a dose, every two hours.

In the same doses as the above may be taken.

Tinctura Colombæ—et

Tinctura Gentianæ composita.

Muriated Tincture of Iron.

R. Tincturæ Ferri muriatis, ℥j.

From twenty to sixty or eighty drops may be taken in a glass of cold chamomile tea, twice or thrice a-day.

Infusion of Colomba and Ginger.

R. Radicis Colombæ contusæ, ℥j.

—— Zingiberis, ℥ij.

Aquæ bullientis, Oj.

Fiat infusio.—Give of the strained liquor, (cold,) a wine-glassful every two hours.

This infusion, when freely used, has proved successful in chronic diarrhœa.

Infusion of Colomba, Rhubarb, &c.

R. Seminum Carui contusorum,

Radicis Colombæ,

—— Rhei, āā. ℥j.

Aquæ ferventis, ℥viiij.

Digest for two hours, and strain.

R. Liquoris colati, ℥ijss.

Tincturæ Rhei, ℥j.

Syrupi Zingiberis, ℥ij.

Misce.—Dose.—A tea-spoonful to a table-spoonful for children, according to their age. In Diarrhœa.

Mixture of Sulphate of Iron and Elixir of Vitriol.

R. Ferri sulphatis, gr. iv.

Acidi sulphurici aromatici, gtt. xl.

Aquæ destillatæ, ℥j.

Fiat mistura, et signa.—A tea-spoonful to be taken three times a-day in half a wine-glass of water.

Vinegar Draught.

R. Acidi acetici, ℥j.

Tincturæ cardamomi compositæ, ℥ss.

Syrupi simplicis, ℥ss.

Aquæ, ℥x.

Misce.—To be taken every twenty minutes as a draught. Advised by Dr. Howard, of London, in sick headach.

Acorn Coffee.

R. Pulveris Glandis torrefactæ, ℥j.

Aquæ bullientis, Oj.

Fiat infusio.—Three or four tea-cupsful may be taken during the day, and augmented according to circumstances.

The above is a very favourite remedy with the Germans. "It is one of the most successful in mesenteric atrophy, commencing rachitis, glandular swellings, asthma, and cough. Continued for a long time, it is one of the most powerful means we possess, for destroying the scrofulous disposition."—*Hufeland's Treatise*, p. 215.

Infusion of Quassia.

R. Ligni Quassiæ,

Radicis Serpentariæ,

Corticis Aurantii contusæ, āā. ℥ss.

Aquæ bullientis, Oij.

Fiat infusio et cola.—Of which a tea-cupful may be taken cold, three times a-day.

Decoction of Bark and Valerian.

R. Corticis Cinchonæ contusæ, ℥j.

Boil in a pint of water for ten minutes, and strain.

R. Radicis Valerianæ contusæ, ʒj.

Aquæ bullientis, Oj.

Infuse for one hour and strain.—Add the decoction of bark to this infusion, and give a tea-cupful cold, three or four times a-day.

This was a favourite prescription with Dr. Parrish, in certain cases of headach, which appear to depend upon rheumatism, and which are entirely unmanageable with the lancet, purgatives, &c. &c.

Infusion of Chamomile and Orange Peel.

R. Florum Anthemidis nobilis, ʒj.

Corticis Aurantii, ʒss.

Aquæ frigidæ, ℥iij.

Macerate for twenty-four hours.—Dose.—A tea-cupful three or four times a-day.

The infusion of chamomile made with cold, is in general more grateful to the patient than when made with boiling water; the same remark applies to infusions generally, and they are also less liable to ferment when made with cold water.

Decoction of Angustura Bark.

R. Corticis Cuspariæ febrifugæ contusæ, ʒj.

Aquæ fontanæ, Ojss.

Boil for fifteen or twenty minutes, and strain.—Of this decoction a wine-glassful may be taken every two hours.

Decoction of Wild-Cherry Tree Bark.

R. Corticis Pruni Virginiani contusi, ʒj.

—— Aurantii contusæ, ʒij.

Aquæ fontanæ, Oj.

Boil the bark alone for half an hour, and then add the orange-peel.—A wine-glassful may be taken every hour or two in consumptive cases, asthma, &c.

Infusion of Serpentaria.

R. Serpentariæ Virginianæ, ʒss.

Aquæ bullientis, Oj.

Infuse in a covered vessel for one hour.—A wine-glassful every hour or two hours, *pro re nata*.

Compound Tincture of Aloes, Gentian, &c.

R. Aloes socotorinæ, ℥j.

Pulveris Zedoariæ,

———— Gentianæ,

———— Croci,

———— Rhei,

———— Agarici,

Syrupi, āā. ℥j.

To be mixed into a quart of brandy. This is the celebrated Baume de Vie, or Elixir of Life. It is a powerful tonic, and one of the most effectual febrifuge medicines. In intermittents, especially, it is justly celebrated. Dose.—A table-spoonful three times a-day.

Infusion of Sage and Boneset.

R. Salviæ officinalis,

Eupatoriæ perfoliatæ, āā. ℥ss.

Corticis cascarillæ, ℥j.

Mix, and infuse in three half pints of boiling water until cold. Dose.—A wine-glassful every two or three hours. In the colliquative perspiration of hectic fever.

Decoction of Dog-Wood Bark.

R. Corticis Corni Floridæ contusæ, ℥j.

Aquæ fontanæ, Oj.

Boil for twenty or thirty minutes, and strain.—A wine-glassful may be given every hour, as a substitute for Peruvian Bark in intermittents.

Infusion of Hops.

R. Humuli lupuli, ℥j.

Aquæ ferventis, Oj.

Infuse for two hours, and strain by expression. Of this a wine-glassful may be taken three or four times a-day. In dyspepsia.

Infusion of Boneset.

R. Eupatorii perfoliati, ℥j.

Aquæ bullientis, Oj.

Infuse for two hours, and strain.—Of this a wine-glassful may be taken every one or two hours, cold. In intermittent fever.

Enema of Extract of Bark, &c.

R. Extracti Cinchonæ, ℥ss.

Aquæ tepidæ, ℥iv.

Solve; dein adde—

Olei olivarum, ℥ss.

Tincturæ Opii, gtt. x.

Fiat enema.—Signa.—To be thrown up every fourth hour. In intermittents.

Where it is preferred, the powdered bark may be used—say ℥ij. mixed with some of the decoction instead of common water. The bark thus administered may occasionally be made to arrest an intermittent when the patient cannot swallow it. Bark is also employed externally in the form of a waistcoat, worn next the skin. A poultice made by mixing the powder and decoction into a paste, is occasionally applied to the region of the stomach; and lastly, the decoction may be used as a semicupium.

Enema of Quinine.

Quinine is employed in enemata, and in this form has an immediate and powerful effect on intermittent fevers. From eight to twelve grains of the sulphate of quinine, dissolved in a tea-cupful of flax-seed or other mucilage, may be administered at a time, and repeated in twelve hours.

Infusion of Cinchona with Lime Water.

R. Cinchonæ corticis cordifoliæ, ℥ij.

Liquoris calcis, Oij.

Misce, et fiat infusio.

This infusion was formerly much employed and esteemed as an excellent compound. A wine-glassful may be taken three or four times per diem, or once every hour, in intermittent and other forms of fever.

GENTIANIN.

This substance is obtained by an elaborate chemical process from the root of the *gentiana lutea*. It is intensely bitter, but possesses no advantages over the common preparations of Gentian. Majendie gives two formulæ for its administration :

R. Gentianinæ, gr. v.

Alcoholis, ʒj.

Fiat tinctura.—Dose.—Two tea-spoonsful, which will contain a little more than a grain of the medicine. Commended in scrofulous affections.

Sirup of Gentianin.

R. Gentianinæ, gr. xvj.

Syrupi, Oj.

Misce.—Dose.—A wine-glassful, which will contain a grain.

Compound Infusion of Gentian.

R. Radicis Gentianæ concisæ, ʒss.

Corticis Aurantii contusæ, ʒij.

Seminis Cardamomi, ʒss.

Aquæ ferventis, Oj.

Fiat infusio.

Gentian is an excellent bitter, and the above will prove useful in wine-glassful doses, in debility of the digestive organs, &c.

ALTERATIVES.

Alteratives have been defined—those medicines which are given with a view to re-establish the healthy functions of the animal economy, without producing any active evacuation. This class of remedies, therefore, embraces numerous preparations which produce a slow but decided effect on the various secreting organs, sometimes without any sensible increase of the

secretions themselves, but in other instances obviously augmenting or magnifying them. Such is the action of minute doses of mercury, iodine and other substances upon the glandular apparatus. These medicines are generally classed with stimulants and tonics, and some of them with narcotics, and others have been placed with those preparations to which they seem most allied when given in an over-dose,—Sialagogues, for example. Without attempting to investigate this question farther than regards practical convenience, we proceed in this place to indicate the more active and important Alteratives; merely premising the familiar truth, that medicines of almost every class of the materia medica become alteratives by being administered in very small doses at intervals of a few hours.

Æthereal Tincture of Iodine.

R. Iodini, gr. vj.

Ætheris Sulphurici, 3j.

Misce.—Thirty drops contain one grain of iodine.—Dose.—Ten drops twice or three times a-day. In cutaneous and glandular diseases.

Compound Tincture of Iodine.

R. Iodinæ, 3j.

Potassæ iodidæ, 3ij.

Spiritus rectificati, Oij.

Fiat tinctura.—Dose.—Five to fifteen drops, *pro re nata*.

Mixture of the Hydriodate of Potash, &c.

R. Iodidi Potassæ, gr. ij.

Sulphatis Magnesicæ, 3ss.

Tartratis Antimonii et Potassæ, gr. ss.

Aquæ destillatæ, 3vj.

Misce.—Signa.—A tea-spoonful three or four times a-day. In scrofula.

Iodine with Hydriodate of Potash.

R. Iodinæ, gr. iij.

Potassæ iodidi, gr. vj.

Aquæ destillatæ, 3j.

Solve et signa.—Dose.—Six to ten drops thrice a-day in a wine-glass of cold water.

If it produce dizziness, pain in the bowels, or other unpleasant symptoms, diminish the quantity.

Dr. Morton ("Illustrations of Pulmonary Consumption,") strongly recommends this mixture in catarrh, phthisis, &c. It has a powerful effect in glandular disease, and in all affections of the mucous membranes.

Mixture of Tincture of Iodine, &c.

R. Tincturæ Iodini, f. ℥j.

Mucilaginis Acaciæ, f. ℥ij.

Aquæ destillatæ, f. ℥vj.

Fiat mistura et signa.—A table-spoonful every two hours. Employed in cases of ulceration accompanied by purulent discharge from the meatus auditorius, and where there exists a scrofulous diathesis.

See Astringents.

Solution of Hydriodate of Iron.

R. Iodidi ferri, ℥j.

Aquæ destillatæ, ℥j.

Fiat solutio.—Dose.—Six to ten drops morning, noon and night, in a wine-glass of cold water.

Solution of Iodide of Potassium.

R. Potassii iodidi, gr. xxxvj.

Aquæ destillatæ, ℥j.

Fiat solutio.—Dose.—Five to twenty drops, morning, noon and night, in a wine-glass of sweetened water.

Majendie.

Solution of the Ioduretted Hydriodate of Potassa.

R. Iodinæ, ℥j.

Potassii iodidi, ℥ij.

Aquæ destillatæ, ℥vj.

Fiat solutio.—Dose.—Six drops, morning and evening, in a wine-glass of water. In scrofulous and scirrhus diseases.

Dunghison, New Remedies, p. 396.

Tyrrell's Mixture of Ioduretted Hydriodate of Potassa.

R. Iodinæ, gr. ss.

Potassæ iodidi, ℥ss.

Syrupi papaveris, ℥ss.

Aquæ destillatæ, ℥ss.

Fiat mistura.—Given in dose of a wine-glassful three times a-day. In syphilis combined with scrofula.

Vide Dunglison, ut supra.

Mixture of Iodo-Hydrargyrate of Potassium.

R. Hydrargyri deuto-iodidi, gr. iv.

Potassii iodidi, ℥j.

Aquæ destillatæ, ℥j.

Misce.—Dose.—Five drops three times a-day, much diluted.

Tincture of Iodo-Hydrargyrate of Potassium.

R. Potassæ iodo-hydrargyratis, gr. j.

Spiritus vini tenuioris, ℥j.

Fiat tinctura.—Dose.—Ten drops three times a-day, much diluted.

Ioduretted Sulphuric Ether.

R. Etheris sulphurici, ℥j.

Iodidi puri, gr. vj.

Solve.—Thirty drops of this solution contain a grain of Iodine. Six drops constitute a full dose.

Majendie's Anti-Epileptic Solution of Iodine.

R. Iodureti potassii, ℥iv.

Iodidi, gr. ij.

Aquæ menthæ destillatæ, ℥vj.

Fiat solutio.—Dose.—A spoonful thrice per diem.

Iodine Waters.

The physicians of Europe, and especially of Paris, have made extensive and salutary use of Iodine waters, both internally and externally. In these the Iodine is very much diluted, and generally with the addition of more or less common salt. These waters are prepared

on a large scale and with great precision by the Pharmaceutists of Paris, but they have been but partially introduced into this country. M. Majendie, however, gives formulæ for some extemporaneous Iodine waters, which may answer all the purposes of the more elaborate preparations. The following is an example:—

Ioduretted Water.

R. Iodureti Potassii, gr. vj.

Iodini, gr. j.

Aquæ puræ, Oij.

Fiat solutio.—This solution is directed to be drank at meals, in place of common water.

That Iodine is one of the most powerful alteratives at present known, there can be no question. There are few diseases dependent upon or connected with morbid secretion of the glandular structures, but what are more or less modified by its use, while in others it acts as a direct curative agent. It is chiefly, however, in scrofulous, glandular and cutaneous affections that it acts most beneficially, dispersing indurations of the glands, and restoring the secretions of the cutaneous and mucous apparatus: whence its advantages in eruptive diseases, bronchitis, leucorrhœa, amenorrhœa, &c. &c. It has been found to restore suppressed hæmorrhoidal and catamenial discharges, and when pushed too far, has even produced alarming hemorrhage of the lungs, nose and uterus. These facts should make the practitioner extremely cautious in its administration, beginning with small doses, and watching their effects. Should it produce any of the inconveniences already mentioned, or dizziness, nausea, purging, burning of the skin or swelling of the gums, it should be at once suspended or given less frequently and in smaller doses.

It is well known that Iodine was first discovered in sea-weed and sponges; and the latter having long had the reputation of being a cure for Goitre, it was found that Iodine was the active remedial agent. We

give below two prescriptions for the use of sponge as formerly in vogue, but now nearly superseded by the preparations of pure Iodine.

Burnt Sponge.

R. Spongiæ ustæ, ℥j.

Sacchari albi, ℥j.

Misce et divide in chartulas, vj.—One three times a-day, gradually augmenting the dose. In Bronchocele.

Decoction of Sponge.

R. Spongiæ concisæ et torrefactæ, ℥j.

Aquæ destillatæ, Oj.

Coque et infunde duodecem horas.—Of this, filtered and rendered palatable by the addition of a little mint water, a table-spoonful or more may be taken every three hours.

STRYCHNINE.

We have already (under the head of Narcotics) given several prescriptions for using this extremely poisonous substance; but as it is chiefly given as an alterative, we here add some additional modes of administration. Strychnine is now much in vogue in the whole class of nervous diseases, paralysis, whether general or local, neuralgia, chorea, hysteria, &c. It is also employed in syphilis, amenorrhœa, dyspepsia, bronchitis, and in fact, in all cases where an alterative course is indicated.

Solution of Acetate of Strychnine.

R. Strychniæ acetatis, gr. iij.

Alcohol, ℥j.

Aquæ cinnamomi, ℥vij.

Misce.—Dose.—Five drops twice a-day, gradually increasing the quantity.

Dunghlison, New Remedies, p. 450.

Tincture of Acetate of Strychnine.

R. Strychniæ acetatis, gr. iss.

Alcohol, ℥ss.

Fiat tinctura.—Dose.—Five to twenty drops thrice per diem. In syphilitic pains of the bones. *Dunglison, ut supra.*

A Sulphate of Strichnine is also in use. It is prepared like the Acetate, of course substituting sulphuric acid. It has no advantages over the preparations already given, and is little used.

See Majendie, Formulaire, 9vme. Ed. p. 30.

Brucine.

The alkaline principle called Brucia or Brucine, is contained in the Augustura bark, Nux vomica and St. Ignatius's bean. That obtained from the bark is preferable for medicinal purposes. It possesses the properties of Strychnia, but in less degree. It may be administered in doses of one to three grains without danger; but the minimum dose should be first given, and the quantity gradually augmented.

Pills of Brucia.

R. Bruciæ puræ, gr. xij.

Conservæ rosarum, q. s., ut fiat massa.

Divide in pilulas xxiv.

Signa.—One pill may be taken four times a-day, and gradually increased.

Tincture of Brucia.

R. Alcoholis, ℥j.

Bruciæ puræ, gr. xvij.

Fiat tinctura.—From six to twenty-four drops may be taken in a draught.

Mixture of Brucia.

R. Bruciæ, gr. vj.

Aquæ destillatæ, ℥iv.

Sacchari albi, ℥ij.

Fiat mistura. Dose.—A table-spoonful night and morning. These preparations are chiefly used in paralysis, and the entire series of spasmodic affections.

Bromine.

This substance, which is allied to chlorine and iodine, is obtained from salt springs and other mineral waters. In medicine it is chiefly used in combination with potassa.

Solution of Bromide of Potassium.

R. Potassii bromidi, gr. xvij.
 Aquæ lactucæ destillatæ, ℥ij.
 Syrupi althææ, ℥j.

Fiat mistura.—Majendie directs the preceding quantity to be given during the twenty-four hours, in doses of a table-spoonful. It is recommended in scrofula, glandular diseases, amenorrhœa, &c.

Pills of Bromide of Iron.

R. Bromidii Furri pulverisate, gr. xij.
 Conservæ rosarum, gr. xvij.
 Gummi Arabici, gr. xij.

Misce, and fiant pilulæ xx secundum artem. Two pills to be taken in the morning, and two in the evening, or, one three times a-day. In Hypertrophy of the Heart, and scrofulous affections.

Majendie.

Mercury.

The mercureal preparations constitute the most decided and efficient medicines of this class. In addition to several formulæ inserted under the head of Sialagogues, the following will be appropriately noticed in this place.

Calomel Pills.

R. Calomelanos, gr. ij.
 Conservæ rosarum, q. s., ut fiant pilulæ xij.

Of which one may be taken every one or two hours. In cases of marasmus, &c. &c.

The use of calomel in minute doses has become very general, and certainly, if we may be allowed to give our experience, very deservedly so. In the hepatic and intestinal complaints of children, it is productive of the happiest results when carefully managed.

In infants, for example, the sixteenth or eighth of a grain, repeated every hour or two hours, according to age and circumstances, will often check diarrhœa, and even dysentery, which have resisted every treatment. If the discharges are attended by pain and tenesmus, small doses of powdered opium or laudanum may be added; but, as a general rule in these cases, the anodyne had better be given by injection.

The preparation called blue pill (*massa ex hydrargyro*) is indicated in similar conditions, and is especially beneficial wherever an alterative is called for. In these cases a grain given at bed time, or even a grain in divided portions through the day, will often produce the desired effect. Hence their extensive use in chronic and obscure derangements of the digestive organs, especially of the stomach and liver;—in the whole class of syphilitic affections, glandular enlargements, &c.

In like manner all the mercurial preparations mentioned under the head of *SIALAGOGUES*, become alteratives by cautious administration in doses more or less minute, and especially when they are given in conjunction with the preparations of sarsaparilla.

Pills of Quinia, Opium, &c.

R. Calomelanos, gr. vj.

Pulveris Opii, gr. iij.

Sulphatis Quiniæ, gr. xij.

Syrupi simplicis, q. s., ut fiat massa et in pilulæ xij dividenda.

Signa —One night and morning.

Dr. Meigs.

Pills of Blue Mass, Quinia, &c.

- R. Massæ ex Hydrargyro,
 Sulphatis Quiniæ,
 Pulveris Aloes, āā. gr. xij.
 Syrupi Rhæi aromatici, q. s. ut fiant pilulæ xij.
 Signa.—One three or four times a-day.

Either of the above prescriptions will be found useful in that condition of the system which supervenes an attack of bilious intermittent or remittent fever. They may be denominated tonic alteratives.

Arsenic.

Arsenic is a powerful alterative, and one which should be administered with the utmost circumspection. In addition to the formulæ already given, the following will be found useful.

Pills of Arsenic and Black Pepper.

- R. Protoxidi arsenici, gr. iv.
 Piperis nigri, ʒix.

Pound these substances in a mortar for a few days at intervals, after which water enough is to be gradually added to form a pill mass, which should be divided into one hundred pills: the French physicians direct double that number, and restrict the patient to one or two pills daily. This preparation of arsenic is regarded in India as an infallible cure for Tuberculous Lepa.

Pills of the Arseniate of Iron.

- R. Proto-arseniatæ Ferri, gr. iij.
 Extracti humuli lupuli, ʒij.
 Pulveris althææ, ʒss.
 Syrupi, q. s. ut fiat massa.

To be divided into forty-eight pills, of which one is given daily. Recommended by Dr. Biett in scrofulous, cancerous, and herpetic diseases.

Hydriodate of Arsenic and Mercury.

- R. Liquoris hydriodatis arsenici et hydrargyri, ʒij.
 Aquæ destillatæ, ʒiiss.
 Syrupi zinziberis, ʒss.

Misce.—To be taken in four draughts, night and morning. Chiefly given in scrofula and cutaneous eruptions.—*Donovan, in Dunglison, New Remedies, p. 288.*

MINERAL ACIDS.

Nitro-Muriatic Acid.

R. Acidi nitrici, ℥j.
 — Muriatici, ℥ss.
 Aquæ destillatæ, ℥vijss.

Misce, et adde—

Spiritus ætheris nitrici, ℥ijss.

Fiat mistura.—Signa.—A tea-spoonful to be given every two hours, in a wine-glass of sweetened water.

These acids, in the same proportions, are also employed as a bath in hepatic derangements, and by some are esteemed highly beneficial. To accomplish this end, put a tea-spoonful of the compound acid into a quart of warm water, and apply it to the region of the liver night and morning, by means of a soft sponge.

Diluted Nitric Acid.

R. Acidi nitrici, ℥j.
 Aquæ destillatæ, Oij.
 Sacchari albi, ℥j.

Fiat mistura.—Signa.—This quantity to be taken daily by the patient.
 In Lues Venerea. *Ferriar, Med. Hist.*

Mixture of Nitric Acid.

R. Acidi nitrici, ℥iss.
 Pulveris Acaciæ,
 Sacchari albi, āā. ℥iij.
 Aquæ destillatæ, ℥vj.

Fiat mistura.—Signa.—A table-spoonful *pro re nata*.

Nitric acid may be prescribed as a tonic in the above form, or simply diluted with water. It is best

taken through a quill, as the teeth suffer when it comes in contact with them. We have seen it arrest the colliquative sweats of hectic fever when other remedies had failed.

Mixture of Nitrous Acid, Laudanum, &c.

R. Acidi nitrici, ʒj.

Misturæ Camphoræ, ʒviiij.

Tincturæ Opii, guttas xl.

Misce.—Signa.—One fourth part to be taken every three or four hours.

This compound is highly recommended by Mr. Thomas Hope, Surgeon, Chatham, in dysentery and cholera.—*N. A. Medical and Surgical Journal*, Oct. 1826.

CLASS XV.

ASTRINGENTS.

DOCTOR CULLEN describes astringents to be "such substances as applied to the human body, produce contraction and condensation in the soft solids, and thereby increase their density and force of cohesion." "And by the operation of this corrugating power," adds Dr. Murray, "either directly exerted on a part, or extended by sympathetic action, the morbid affections arising from a state of relaxation are supposed to be removed."

In the use of this class of medicines there is one caution suggests itself; namely, that they should not be given to arrest evacuations that are critical, and designed by nature to relieve a plethoric state of the system; unless, however, such evacuation exist to an alarming extent.

PILLS, POWDERS, &c.

Compound Powder of Alum.

R. *Aluminis*, ʒj.

Pulveris Kino, gr. v.

Misce.—Fiant pulveres, v.—Signa.—One every two hours. In obstinate diarrhœa.

This compound, though much employed, and praised by some practitioners, contains substances incompati-

ble with each other. It is, nevertheless, very useful in menorrhagia, &c.

Powder of Sugar of Lead and Calomel.

R. Hydrargyri chloridi mitis, gr. ij vel iv.

Acetatis Plumbi, gr. ss ad gr. j.

Fiant pulveres iv.—One every three hours. *In Cholera Infantum.*

The above dose is adapted to any period between ten and twenty months of age.

Dr. T. D. Mitchell.

Powder with Alum and Opium.

R. Aluminis, ʒss.

Pulveris Opii, gr. iij.

Fiant pulveris vj.—Signa.—One to be taken every three or four hours. In diarrhœa.

Powder of Blue Vitriol, &c.

R. Sulphatis Cupri,

Boli Armeniæ, āā. ʒj.

Camphoræ, ʒij.

Fiat pulvis.—Project ʒj. of this powder into a pound of boiling water—filter the solution, and mix ʒj. of it with ʒj. of clear water, and apply it to the purulent eyes of infants by means of a syringe, *pro re nata*.—See *Cooper's Surgery*.

Pills of Acetate of Lead and Calomel.

R. Acetatis plumbi, ʒss.

Calomel, gr. v.

Conservæ rosarum, q. s.

Ut fiant pilulæ x. One to be given every two, three, or four hours, according to circumstances. In Hemorrhage, and especially in Hematemesis.

Sugar of Lead and Opium.

R. Acetatis plumbi, ʒj.

Gum Opii, gr. j.

Misce, et divide in pilulas xij.—A pill to be taken every hour at first, and then every two hours. This is Dr. Graves's prescription for the treatment of Cholera.

Pills of Acetate of Lead and Opium.

R. Acetatis Plumbi, gr. xij.

Pulveris Opii, gr. vj.

Conservæ rosarum, q. s., ut fiat massa, in pilulas vi dividenda.

One to be taken for a dose, according to circumstances. In hæmoptysis and other hemorrhages.

Bolus with Alum and Extract of Bark.

R. Aluminis,

Extracti Cinchonæ,

Nucis myristicæ, aa. ʒss.

Syrupi simplicis, q. s., ut fiat bolus.

To be given in uterine and other hemorrhages, and repeated *pro re nata*.

Pills of Alum, Catechu, &c.

R. Aluminis, gr. vj.

Extracti Opii, gr. j.

Catechu, gr. j.

Misce.—To be divided into six pills, one of which is to be given every two, three or four hours. In passive hemorrhage, and in atonic mucous discharges.

Pills of Rhatany.

R. Extracti Rhatanix, ʒj.

Pulveris kino, q. s.

Ut fiant pilulæ viij.—Signa.—A pill to be given every hour or two hours. In the Profluvia.

Pills of Creosote.

R. Creosoti, gtt. x.

Pulveris glycyrrhizæ, ʒj.

Mucilaginis, q. s.

Ut fiant xx.—Dose—a pill thrice in the day. In Bronchitis, Neuralgia, &c.

Creosote Mixture.

R. Creosoti, gtt. xij.

Mucilaginis Arabici, ℥iv.

Sacchari albi, ℥j.

Fiat mistura.—Dose—a table-spoonful every two or three hours. In Hemorrhages.

Alcoholic Solution of Creosote.

R. Creosoti, gtt. j.

Alcoholis rectificati, gtt. xvj.

Misce.—A drop or two of this solution dropped into the cavity of an aching tooth, is said to afford immediate relief. *Marinus.*

Tannin Pills.

R. Tanninæ puræ, gr. ij.

Gummi vel Syrupi, q. s.

Ut fiant pilulæ viij. One may be given every two or three hours. In Gonorrhœa, &c-

Chalk Mixture.

R. Cretæ præparatæ: ℥iss.

Sacchari albi,

Gummi Arabici, āā. q. s.

Aquæ destillatæ, ℥iv.

Olei cinnamomi, gtt. ij.

Tincture Opii, gtt. xl vel lx.

Misce.—Dose—a table-spoonful every two hours. In Diarrhœa and Dysentery.

Another formula for chalk mixture will be given under the head of Absorbents and Antacids.

When a strongly astringent chalk mixture is desirable, we may add to the above formula Tinct. Kino, ℥ij., or the same quantity of Tinctura Rhataniæ, or Catechu, or Nut-galls.

Camphor, Laudanum and Nitric Acid.

R. Aquæ camphoræ, ℥iv.

Acidi nitrici, gtt. iv.

Tincturæ Opii, gtt. xl vel lx.

Fiat mistura. Dose—a table-spoonful every two hours. In Diarrhœa and Dysentery.

This is the well known and very efficacious preparation called *Hope's Mixture*. If the nitric acid is added in larger proportion, it is very liable to defeat the object of the prescription.

Camphor Water, Lavender, &c.

R. Aquæ camphoræ, ℥iv.

Tincturæ lavendulæ compositæ, ℥ij.

Tincturæ Opii, gtt. xl vel lx.

Fiat mistura. Dose—a table-spoonful every two hours.

This prescription is employed in the same cases as the preceding one, and was strongly commended by the late Dr. Parrish.

Electuary of Catechu and Opium.

R. Electuarii Catechu, ℥j.

Suited to cases of diarrhœa, in tea-spoonful doses. Ten scruples contain one grain of opium.

Pills of Kino, Opium, &c.

R. Pulveris Kino, gr. xx.

——— Opii, gr. ij.

Mucilaginis Acaciæ, q. s. ut fiant pilulæ iv.

Of which take one every four hours. In diarrhœa, pyrosis, &c.

Monesia.

This substance is the product of a tree which grows in Chili, but its botanical affinities are yet unknown. It is given in the form of Extract, twelve or twenty grains daily, divided into three grain pills. In Leucorrhœa, Diarrhœa, Menorrhagia, &c. &c.

COLLYRIA.

These are preparations applied to the eyes; they are sometimes dry, but generally liquid—consisting of infusions, decoctions, or distilled waters, with the addition of various substances.

Dry Collyrium of Sugar, &c.

R. Sacchari albi,
Oxidi Zinci, āā. partes equales.
Tere in pulverem.

M. Recamier.

Dry Collyrium of Mercury, &c.

R. Sacchari albi, ʒij.
Oxidi Hydrargyri rubri, gr. x.
Oxidi Zinci impuri præparati, ʒj.
Fiat pulvis.

M. Dupuytren.

Dry Collyrium of Opium, &c.

R. Pulveris Opii, gr. iv.
Hydrargyri chloridi mitis,
Sacchari purificati, āā. ʒj.
Tere bene.

These compounds are blown into the eyes for the removal of specks on the cornea resulting from inflammation.

Alum Curd.

R. Pulveris Aluminis, ʒss.
Albuminem Ovi, j.

Agitate them well until a coagulum be formed.—To be applied on a linen rag to inflamed eyes. It should be removed in three or four hours.

Alum Collyrium.

R. Pulveris Aluminis, gr. xij vel xx.
Aquæ rosarum, ʒiv.

Fiat solutio.—To be applied to the eye in chronic inflammation

Collyrium of Ioduret of Potassium.

R. Aquæ rosarum, ℥vj.
 Iodureti potassii, gr. xxiv.
 Iodini, gr. j vel ij.

Fiat collyrium. To be applied to the eyes three or four times in the day. In scrofulous ophthalmia, and ulcerations of the cornea and conjunctiva.

Collyrium of Acetate of Zinc.

R. Sulphatis Zinci,
 Acetatis plumbi, āā. gr. vj.
 Aquæ rosarum, ℥iv.
 Fiat solutio.

In this prescription a double decomposition takes place. Acetate of zinc is held in solution, while sulphate of lead falls to the bottom. It should be filtered, or used without shaking.

Collyrium of Acetate of Lead, &c.

R. Liquoris plumbi acetatis, gtt. xij.
 Vini Opii, gtt. xl.
 Aquæ rosarum, ℥iv.
 Fiat collyrium.

Collyrium of Vinegar, &c.

R. Aceti destillati, ℥j.
 Spiritus Vini diluti, ℥ss.
 Aquæ rosarum, ℥viij.
 Fiat mistura.—Signa.—To be applied to weak eyes, and in ophthalmia after depletion.

Collyrium of Acetate of Ammonia und Camphor.

R. Liquoris Ammoniaæ acetatis, ℥ij.
 Misturæ Camphoræ, ℥vj.
 Misce.—Signa.—To be applied to eyes requiring a mild astringent and stimulant remedy.

Collyrium of Acetate of Ammonia with Opium.

R. Liquoris Ammoniaë acetatis, ℥ij.

Aquæ ferventis, ℥vj.

Extracti Opii mollis, gr. x.

Dissolve the extract of Opium in the hot water—strain, and add the liquor of acetate of ammonia. Useful in acute ophthalmia when considerable pain remains after depletion.

Collyrium of Sulphate of Copper.

R. Sulphatis Cupri, gr. vj.

Camphoræ, ℥j.

Aquæ ferventis, ℥viiij.

Rub the camphor with the water—then strain, and add the sulphate of copper.

This is nearly the same as the formula recommended by Mr. Ware in the treatment of purulent ophthalmia.

Collyrium of Corrosive Sublimate.

R. Hydrargyri chloridi corrosivi, gr. ij.

Aquæ destillatæ, ℥viiij.

Fiat solutio.

The above may be advantageously used in gonorrhœal ophthalmia, as also in that of a scrofulous character.

Collyrium of Opium and Camphor.

R. Extracti Opii mollis, gr. x.

Camphoræ, gr. vj.

Aquæ ferventis, ℥xij.

Rub the camphor and opium well together in a mortar and add the water. Then strain before using.

Collyrium of Poppies, &c.

R. Fomenti papaveris capsularum, ℥iv.

Aquæ rosarum,

Misturæ Camphoræ, aa. ℥ij.

Misce.—Either of the last two collyria may be employed in the early stages of ophthalmia.

All of these remedies can be most successfully applied to the eyes by introducing the point of a small syringe into the outer angle, and gradually injecting the collyria.

Collyrium of Nitrate of Silver.

R. Nitratis Argenti, gr. ij.

Aquæ destillatæ, ℥ij.

Fiat solutio.—To be applied at the close of acute ophthalmia, and to promote the absorption of nebulæ of the cornea.

Emollient Collyrium.

R. Radicis Althææ officinalis, ℥ij.

Aquæ destillatæ, ℥j.

Infuse for three hours near the fire, and strain.—Employed in inflammation of the conjunctiva accompanied with violent irritation.

Anodyne Collyrium.

R. Colchici autumnalis, ℥j.

Aquæ Lini bullientis, ℥iv.

Tincturæ Opii, ℥j.

Fiat mistura.—Employed in ophthalmia accompanied with great sensibility.

M. Jadelot.

Strichnine Collyrium.

R. Strichniæ puræ, gr. ij vel iv.

Acidi acetici diluti,

Aquæ destillatæ, āā, ℥j.

Fiat solutio.—A drop or two, diluted or not according to circumstances, to be applied to the eye several times a-day. Commended by Dr. Ryan in Amaurosis.

INJECTIONS

Injectiōns are liquid compositions, containing emollient, astringent, or stimulating substances. The parts into which they are thrown are, the external auditory passages, the canal of the urethra, the bladder, and also fistulous cavities, the tunica vaginalis in the case of hydrocele, and the vagina. When employed in the cure of gonorrhœa, two circumstances merit especial attention, viz. the manner of throwing them up, and the frequency of using them. When the common syringe is used, the pipe should be perfectly smooth, short, and conical, so that it cannot pass more than a quarter of an inch into the urethra. The patient should oil the pipe before using it, seat himself on the edge of a chair so as to avoid pressing the perinæum, draw the penis forward on the syringe with the left hand, and throw up the injection to the seat of disease, (previously ascertained,) with the other. As respects the frequency of injecting, it may be observed, that six or eight times a-day are not too often, and that the cure will be retarded by using these remedies at longer intervals. We cannot forbear remarking, that gonorrhœal inflammation, situated near the extremity of the penis, can be often cured by a *routine plan*; but when seated higher up the canal, and affecting Cowper's gland or the prostate, it becomes a serious disease, the cure of which ought not to be undertaken by those ignorant of the structure of the parts and the nature of morbid action.

Injection of Pyroligneous Acid for the Ear.

R. Acidi pyrolignei, ℥ij.

Aquæ destillatæ, ℥vj.

Fiat injectio.—Signa.—To be injected twice a-day with a silver syringe, in purulent discharge from the meatus auditorius.

Buchanan on the Ear.

Injection for Catechu for the Ear.

R. Extracti Catechu, gr. xij.

Aquæ bullientis, ℥vj.

Fiat injectio.—To be used four or five times a-day. *Idem.*

Pyroligneous Acid, Ether, &c., for the Ear.

R. Acidi pyrolignei,

Spiritus Ætheris sulphurici,

Olei Terebinthinæ, aa. partes equales.

Misce.—Signa.—Introduce two drops into the meatus auditorius every night, when the cerumen is deficient in quantity.—*Idem.*

Injection of Sulphate of Zinc, &c.

R. Sulphatis Zinci, gr. x.

Pulveris Acaciæ, ℥ij.

Tincturæ Opii, ℥j.

Aquæ destillatæ, ℥viij.

Fiat injectio.—Used in gonorrhœa virulenta.

Injection of Corrosive Sublimate, &c.

R. Sulphatis Zinci, gr. x.

Chloridi Hydrargyri corrosivi, gr. ij.

Aquæ rosarum, ℥viij.

Fiat injectio.—To be used in gleet.

This powerful remedy should be employed with great caution.

Peruvian Balsam, &c., for the Ear.

R. Fellis Bovini, ℥iij.

Balsami peruviani, ℥j.

Misce.—Signa.—To be dropped occasionally into the ear to correct a fetid discharge, syringing the ear every day with a weak solution of soap and water.

Dr. Hugh Smith.

Injection of Muriate of Ammonia, &c.

R. Muriatis Ammoniaë, gr. j vel ij.
 Chloridi Hydrargyri corrosivi, gr. ij.
 Aquæ destillatæ, ℥iv.
 Fiat injectio.—To be used in chronic gonorrhœa.

Injection of Acetate of Zinc.

R. Acetatis Zinci, gr. viij.
 Aquæ rosarum, ℥iv.
 Fiat injectio.—Used in gonorrhœa virulenta.

For all these preparations, rose water is an admirable menstruum ; it possesses the advantages of being slightly astringent and rendered pure by distillation, and seems therefore particularly applicable to the formation of injections and collyria.

Injection of Armenian Bole.

R. Sulphatis Zinci, ʒj.
 Boli Armeniaë, ʒij.
 Pulveris Acaciæ, ʒij.
 Aquæ rosarum, ℥viij.
 Fiat injectio.—To be used in gonorrhœa.

Injection of Calomel, &c.

R. Hydrargyri chloridi mitis, ʒj.
 Gummi Arabici,
 Tincturæ Opii, āā. ʒss.
 Aquæ destillatæ, ℥viij.
 Fiat injectio.—Signa.—For gonorrhœa or gleet.

Injection of Sulphate of Copper.

R. Cupri sulphatis, gr. vj.
 Aquæ destillatæ, ℥vj.
 Tincturæ Opii, ʒj.
 Fiat solutio.—Signa.—To be thrown up the urethra in chronic gonorrhœa.

Injection of Tea.

R. Theæ viridis foliorum, ʒj.

Aquæ ferventis, ʒvj.

Fiat infusio et cola.—To be used as an injection in gonorrhœa virulenta.

Injection of Ammoniuret of Copper.

R. Cupri Ammoniureti, gr. v.

Aquæ rosarum, ʒviij.

Fiat solutio.—To be used in chronic gonorrhœa.

GARGLES, MOUTH WASHES, &c.

Gargles, like injections, are astringent and stimulating; to be of any service they must be frequently applied and persevered in for some time.

Gargle of Brandy.

R. Spiritus vini gallici,

Aquæ fontanæ, āā. partes equales.

Brandy and water form an excellent gargle for some cases of inflamed fauces.

Gargle of Galls and Alcohol.

R. Infusionis gallarum, ʒvij.

Spiritus vini rectificati, ʒj.

Fiat gargarysma.—To be used frequently, in relaxation of the uvula and soft palate.

Gargle of Infusion of Oak Bark.

R. Corticis Querci falcata, ʒj.

Aquæ fontanæ, ℥jss.

Boil to a pound, and strain; then add—

Sulphatis aluminis, ʒss.

Spiritus vini gallici, ʒij.

Misce.—To be applied to the uvula when swelled or relaxed, and to be used as an injection in fluor albus.

Gargle of Sulphate of Quinia, &c.

R. Sulphatis Quiniæ, gr. xij.

———— cupri, gr. xvj.

Acidi sulphurici aromatici, ℥j.

Aquæ fontanæ, ℥viij.

Fiat mistura.—Signa.—A table-spoonful to be used three or four times a-day, in obstinate sore throat.

Dr. Hartshorne.

Gargle of Borax, &c.

R. Boratis sodæ, ℥j.

Tincturæ myrrhæ, ℥ss.

Mellis despumati, ℥j.

Aquæ rosarum, ℥iv.

Fiat gargarysma.—Signa.—To be used as a gargle or mouth wash, in ptyalism or scorbutic affections of the gums.

Gargle of Sulphate of Zinc, &c.

R. Sulphatis Zinci, ℥j.

Mellis, ℥ss.

Tincturæ Myrrhæ, ℥j.

Spiritus vini gallici, ℥j.

Aquæ rosarum, ℥iv.

Fiat gargarysma.—To be applied to ulcerated gums, occasioned by excessive salivation. Water can be added if it be too strong.

In many cases of mercurial sore mouth, common lead water forms a most soothing and useful application. It, makes the teeth black—an effect which is only temporary.

Dr. Blake's Toothach Solution.

R. Aluminis, ℥ij. (in pulverem subtilem tritandum.)

Spiritus ætheris nitrici, ℥vij.

Fiat solutio.—Said by Dr. Blake, of London, to be an infallible cure for toothach, where this is not connected with rheumatism.

Gargle of Alum.

R. Sulphatis aluminæ et potassæ, ℥ij.

Aquæ, ℥iv.

Solve.—This is Dr. Cavarra's gargle for fœtor oris, or offensive breath.

Gargle of Pomegranate.

R. Corticis punicæ granati, ℥ss.

Petalarum rosæ rubræ, ℥ij.

Aquæ bullientis, ℥vj.

Fiat infusio.—Cola et adde—

Mellis despumati, ℥j.

Aluminis, ℥ij.

Fiat gargarysma.—Signa.—To be used in inflammations of the throat and fauces. This gargle is highly praised by Dr. Meigs.

Muriatic Acid Gargle.

R. Infusionis Cinchonæ, ℥iv.

Mellis despumati, ℥j.

Acidi muriatici, gtt. xvij.

Fiat gargarysma.—To be used in chronic inflammation, when pain and swelling have been reduced by depletion—also in cynanche maligna.

Ratier's Formulary.

Wash of Sulphate of Copper, &c.

R. Sulphatis Cupri, ℥ij.

Pulveris Cinchonæ, ℥ss.

Aquæ destillatæ, ℥iv.

Misce.—Signa.—To be used in the gangrenous sore mouth of children; and to be applied to the full extent of the ulceration and excoriation.

Dr. B. H. Coates.

Linctus with Muriatic Acid, &c.

R. Acidi Muriatici, ℥j.

Mellis,

Aquæ rosarum, āā. ℥j.

Misce.—Signa.—To be applied to scorbutic gums three or four times a-day.

Detergent Gargle—Sulphuric Acid, &c.

R. Decocti Hordei, ℥iv.

Mellis despumati, ℥ss.

Acidi sulphurici, gtt. xx.

Fiat gargarysma.

Ratier's Formulary.

Mercurial Gargle.

R. Chloridi Hydrargyri corrosivi, gr. ij.

Mellis despumati, ℥ss.

Aquæ destillatæ, ℥iv.

Fiat solutio.—To be used in venereal sore throat.

Idem.

Acetous Gargle.

R. Acidi acetici, ℥ij.

Muriatis Ammoniacæ, ℥j.

Mellis, ℥iss.

Aquæ fontanæ, ℥xij.

Fiat gargarysma.—To be used in inflamed fauces.

Idem.

Astringent Gargle of Acetate of Lead, &c.

R. Acetatis Plumbi, ℥ss.

Syrupi simplicis, ℥j.

Decocti Hordei, Oj.

Fiat solutio.—In mercurial sore mouth, &c.—*Ratier's Formulary.*

MIXTURES, INFUSIONS, &c.

Infusion of Galls.

R. Gallæ contusæ, ℥ij.

Aquæ bullientis, ℥x.

Infuse for four hours, and strain.—Signa.—To be taken in doses of a table-spoonful every hour or two hours. In diarrhœa. It may be rendered more agreeable by the addition of some aromatic.

Infusion of Galls and Chalk.

R. Infusionis Gallarum, ℥iv.

Cretæ præparatæ, ℥ss.

Tincturæ Opii, ℥j.

Gummi Acaciæ, ℥j.

Fiat mistura.—Dose.—A table-spoonful every two hours. In diarrhœa.

Infusion of Logwood.

R. Hæmatoxyli ligni, ℥ss.

Aquæ bullientis, Oj.

Infuse for two or three hours, and strain.—Of this a table-spoonful to be given to children every two or three hours. In cholera and diarrhœa.

Mixture of Extract of Logwood, &c.

R. Extracti Hæmatoxyli, ℥ij.

Tincturæ catechu, ℥ij.

Aquæ fontanæ, ℥vij.

Misce.—Signa.—Two spoonfuls to be given once in three or four hours. In the advanced stages of diarrhœa and dysentery.

Mixture with Extract of Bark and Alum.

R. Extracti Cinchonæ, ℥ss.

Aluminis, ℥ij.

Aquæ Cinnamomi, ℥v.

Syrupi Limonis, ℥j.

Misce.—Signa.—Two spoonfuls may be taken every two hours. In low diseases accompanied with hemorrhage.

Mixture with Tincture of Catechu, &c.

R. Tincturæ Catechu, ℥j.

——— Opii, gtt. lx.

Pulveris Acaciæ, ℥ij.

Aquæ Cinnamomi, ℥vj.

Misce.—Signa.—A table-spoonful every two hours. In diarrhœa.

Solution of Vitriolated Zinc and Alum.

As under Emetics.

Alum Whey.

R. Lactis Vaccinæ bullientis, Oj.

Pulveris Aluminis, ℥ij.

Boil them until the coagulum separates, and strain. A cupful to be taken occasionally. In diabetes, &c.

Peruvian Bark with Lime Water.

R. Pulveris corticis Cinchonæ, ℥ss.

Aquæ Calcis, ℥viij.

Misce.—Signa.—A table-spoonful to be taken every two hours, or oftener if necessary. In diarrhœa and menorrhagia.

Sirup of Galls, Brandy, &c.

R. Gallarum contusarum, ℥ij.

Spiritus gallici, ℥viij.

Misce.—Set the brandy on fire and burn out the spirit;—then add—

Corticis Cinnamomi,

Macis, āā. ℥ij.

While hot, add sugar by burning it over the inflamed brandy until a sirup is formed.—Decant the pure sirup.—Dose.—A table-spoonful every two hours. In chronic and obstinate diarrhœa.

Dr. Parrish.

Clyster of Lime Water and Catechu.

R. Electuarii Catechu, ℥ss.

Aquæ Calcis, ℥x.

Fiat solutio.—Signa.—One-half may be injected at a time; in the diarrhœa of continued fevers.

Solution of Sulphate of Copper, &c.

R. Sulphatis Cupri, gr. iij.

Acidi sulphurici, gtt. xx.

Aquæ destillatæ, ℥ij.

Fiat solutio.—Signa.—From twenty to forty drops to be given every one or two hours in sirup, *pro re nata*. In menorrhagia.

Thatcher's Practice.

Enema of Acetate of Lead, &c.

R. Acetatis Plumbi, ℥j.

Tincturæ Opii, gtt. lx.

Aquæ tepidæ, ℥ij.

Fiat solutio.—Signa.—To be thrown up the rectum in uterine hemorrhage.

Dewees's Syst. of Mid., p. 420.

Mixture of Extract of Rhatany Root, &c.

R. Extracti Rhatanix, ℥j.

Syrupi papaveris,

Aquæ rosarum, āā. ℥j.

Fiat solutio.—Signa.—A tea-spoonful for a dose, *pro re nata*. In passive hemorrhage and chronic dysentery.

M. Fouquier.

LINIMENTS, LOTIONS, &c.

Lotion of Lime Water, Zinc, &c.

- R. Oxidi Zinci, gr. xij.
 Cupri sulphatis, gr. iij vel iv.
 Mellis rosacei, ʒj.
 Aquæ calcis, ʒij.

Fiat lotio.—Recommended as very useful in phagedænic ulcers of the genitals.
Vide Swediaur on Syphilis.

Liniment of Lime Water and Linseed Oil.

- R. Olei Lini, ʒiij.
 Liquoris Calcis, ʒvj.

Fiat linimentum.—To be applied to burns and scalds by means of a feather or soft rag.

Lotion of Lime Water and Alcohol.

- R. Spiritus vini rectificati, ʒij.
 Liquoris Calcis, Oss.

Fiat mistura.—To be applied in the same cases as the above.

Lotion of Goulard's Extract of Lead, &c.

- R. Liquoris acetatis plumbi, ʒj.
 Spiritus Camphoræ, ʒiij.
 Aquæ fontanæ, Oj.

Fiat lotio.—To be used as the former.

Chloride of Soda.

- R. Chloridi Sodæ, quantum placet.

This compound is much recommended by some of the French surgeons, applied by means of lint in the treatment of old ulcers.

Epithem of Acetate of Lead.

- R. Confectionis rosæ, ʒj.
 Mellis,
 Liquoris plumbi acetatis,
 Tincturæ Opii, āā. ʒij.

Misce.—Signa.—To be applied to painful venereal ulcers, and to chancres after the use of lunar caustic.

Compound Lotion of Sulphuret of Potash.

R. Sulphureti Potassæ, ℥ss.
Saponis venetæ, ℥j.
Spiritus rectificati, ℥iv.
Tincturæ Myrrhæ, ℥ss.
Liquoris Calcis, Oj.

Fiat mistura secundum artem.—To be applied to the head night and morning, in cases of tinea capitis. The hair must be shaved previous to its employment.

Astringent Fomentation.

R. Radicis Bistortæ,
Corticis Punicæ granati, āā. ℥ij.
Muriatis Ammoniacæ, ℥ij.
Vini rubri, Oj.

Misce.—Signa.—To be applied to indolent inflammatory swellings.
Ratier's Formulary.

Lotion of Bark.

R. Cinchonæ rubræ contusæ, ℥j.
Aquæ fontanæ, Oj.
Boil for ten minutes, strain, then add—
Alcoholis camphorati, ℥viij.

This preparation may be used whenever a tonic and stimulant lotion is required.

Lotion of Sulphuret of Potassa, &c.

1. R. Sulphureti Potassæ, ℥j vel ℥ij.
Aquæ, Oj.

Misce.—

2. R. Acidi hydrochlorici, ℥j vel ℥ij.
Aquæ destillatæ, Oij.

Mix one ounce of each lotion with four ounces of warm water, and apply by means of a sponge in cases of psora.—*M. Alibert.*

OINTMENTS.

Ointment of Alum, Calomel, &c.

R. Hydrargyri chloridi mitis, ℥ij.

Sulphatis Aluminis exsiccati,

Oxidi plumbi albi, āā. ℥ss.

Olei Terebinthinæ, ℥ij.

Cerati simplicis, ℥iss.

Fiat unguentum.

This ointment is particularly efficacious in *Tinea Capitis*. It should be applied every night going to bed, and washed off in the morning; as soon as the child's head is dry, it should be dusted over with powdered starch, which in the evening is again to give place to the ointment. The hair should be cut off *previous to the application of the ointment*.

Ointment of Gall-nuts.

R. Pulveris Gallarum, ℥j vel ij.

Adipis suillæ, ℥j.

Fiat unguentum.—Used in hæmorrhoidal affections.

Ointment of Oxide of Zinc.

R. Oxidi vel florum, ℥j.

Butyri recentis, ℥vj.

Fiat unguentum.—To be applied to the eyes in cases of defluxions arising from relaxation or slight inflammation of the lids.

Ointment of Rhubarb, Opium, &c.

R. Pulveris Rhei, gr. x.

——— Opii, gr. v.

Cerati simplicis, ℥iss.

Fiat unguentum.—This ointment has been found very efficacious in indolent ulcers.

Ointment of Tar, &c.

R. Unguenti picis liquidæ, ℥j.

——— hydrargyri nitratis vel citrini, ℥ss.

Fiat mistura.—In tinea capitis.

Ointment of Tur and Opium.

R. Pulveris Opii, ℥ij.

Unguenti picis, ℥j.

Fiat unguentum.—Signa.—Hæmorrhoidal ointment.

This compound has been found very efficacious in hæmorrhoids.

Kirkland's Neutral Cerate.

R. Emplastri plumbi, ℥iv.

Olei olivarum,

Cretæ præparatæ,

Acidi acetici, aa. ℥ij.

Super-acetatis plumbi, ℥jss.

The plaster and oil are first to be melted over a gentle fire. The prepared chalk is then to be briskly stirred in, and when the mixture is sufficiently cooled, the vinegar and superacetate of lead are to be added, and the whole stirred until nearly cold.

This cerate, which is not to be met with in the pharmacopœia, is strongly recommended in abraded irritable surfaces, and the latter stages of burns.

Aromatic Sulphur Ointment.

R. Potassæ carbonatis, ℥ss.

Aquæ rosæ, ℥j.

Hydrargyri sulphureti rubri, ℥j

Olei essentialis Bergamotæ ℥ss.

Sulphuris sublimati et loti,

Adipis suilli, aa. ℥x.

Misce secundum artem.—Employed in scabies.

Bateman's Synopsis.

CLASS XVI.

ABSORBENTS OR ANTACIDS.

ABSORBENTS or Antacids form a class of remedies for correcting acidity in the stomach and bowels. As a general rule they should be preceded by an emetic or cathartic, or both. When a permanent effect is desired, they are usually combined with tonics, aromatics, and sometimes narcotics. When given alone, they too frequently afford but transient relief.

POWDERS.

Compound Cretaceous Powder with Opium.

R. Cretæ præparatæ, ℥iss.

Pulveris Cinnamomi, ℥j.

——— Piperis longi, gr. viij.

——— Opii, gr. vj.

Fiat pulvis.—Divide in chartulas xij.—One three or four times a-day.
In diarrhœa.

The above formula, with a slight alteration, is from the London Pharmacopœia.

Powder with Crabs' Claws and Rhubarb.

R. Chelæ cancerorum, ℥j.

Pulveris Rhei, ℥ij.

Misce, et divide in pulveres iv.—Of which one may be taken once or twice a-day, according to circumstances. In acidity of the *Primæ viæ*.

This preparation is called by the several names of crabs' claws, eyes, and stones. It is a carbonate of lime united with some animal gelatine, and which, like prepared oyster shells, has little or no superiority over the prepared chalk.

Powder with Magnesia, Fennel Seeds, &c.

R. *Magnesia*, ℥ss.

Pulveris seminum Fœniculi,

—— *Cinnamomi*, āā. ʒj.

Fiat pulvis.—Dose.—A small tea-spoonful three or four times a-day.

Powder with Soda, Magnesia, &c.

R. *Magnesia calcinata*, ℥ss.

Bi-carbonatis Sodæ,

Pulveris Zingiberis, āā. ʒj.

Misce.—Signa.—A small tea-spoonful occasionally. In cardialgia.

Powder with Magnesia and Colombo.

R. *Magnesia ustæ*, ʒjss.

Pulveris Colombæ, ʒj.

Misce.—Divide in chartulas vj.—Of which one may be taken three times a-day, in sugar and water. In Cardialgia.

PILLS.

Pills of Soda, Rhubarb, &c.

R. *Pulveris Rhei*,

Carbonatis Sodæ exsiccata,

Extracti Gentianæ, āā. ʒj.

Hydrargyri chloridi mitis, gr. iij.

Misce.—Fiant pilulæ xx.—Signa.—Two occasionally. In dyspepsia.

MIXTURES, SOLUTIONS, &c.

Chalk Mixture.

R. Cretæ præparatæ, ℥iij.

Sacchari albi,

Pulveris gummi Acaciæ, āā. ℥iss.

Aquæ Menthæ, ℥vj.

Fiat mistura.—Signa.—A table-spoonful for a dose every two hours. When administered in Diarrhœa, Astringents are added.

Solution of Salt of Tartar.

R. Carbonatis Potassæ, ℥j.

Sacchari albi, ℥j.

Tincturæ Lavandulæ compositæ, ℥ij.

——— Opii, gtt. xl.

Aquæ Menthæ, ℥iv.

Misce.—Signa.—A table-spoonful every hour or two hours. In sickness of the stomach occasioned by acidity.

The opium in the above prescriptions may be omitted at the discretion of the practitioner.

Mixture of Salt of Tartar, Ammonia, &c.

R. Spiritus Ammoniæ,

Carbonatis Potassæ, āā. ℥ij.

Aquæ Cinnamomi, ℥iv.

Misce.—Signa.—A large tea-spoonful for a dose *pro re nata*. In Pyrosis.

Lime Water and Milk.

R. Aquæ calcis,

Lactis recentis, āā. ℥iij.

Misce.—Signa.—Two table-spoonsful for a dose. In sickness of the stomach few remedies are more prompt in their effect. An antacid remedy admirably adapted to children, is prepared by pouring four ounces of boiling water on a heaped tea-spoonful of calcined magnesia, and giving a dessert-spoonful of the clear solution every hour or two. It is slightly aperient.

Mixture with Ammonia, Magnesia, &c.

R. Magnesiae ustæ, ʒj.
 Aquæ Ammoniae, ʒj.
 Spiritus Cinnamomi, ʒiij.
 Aquæ destillatæ, ʒvss.

Misce.—Signa.—A table-spoonful every two hours. Usually given in the sickness accompanying pregnancy.

Mixture of Magnesia, Camphor, &c.

R. Magnesiae albæ, ʒj.
 Camphoræ, ʒss.
 Sacchari albi,
 Gummi Acaciæ, āā. ʒij.
 Ætheris sulphurici, ʒss.
 Aquæ destillatæ, ʒiv.

Misce.—Signa.—A table-spoonful for a dose four or five times a-day. In flatulency and irritable stomach.—*Dr. E. Griffiths.*

Dyspeptic Lie.

Take of

Hickory Ashes	-	-	-	-	-	1 quart,
Soot	-	-	-	-	-	6 ounces,
Boiling Water	-	-	-	-	-	1 gallon.

Mix and let them stand for twenty-four hours, frequently stirring the ingredients. A tea-cupful may be given of this liquor, (decanted,) three times a-day. It should be decanted always at the end of twenty-four hours, for by standing on the materials a long time the water acquires almost caustic properties, and may do serious injury.

This domestic remedy we have thought best to couch in the English language. It deserves an insertion from the circumstance that it was employed by Dr. Physick in his own case, and we are informed was of decided advantage when the usual alkaline remedies had failed.

Solution of Salt of Tartar, &c.

R. Carbonatis Potassæ, ℥j.
 Pulveris gummi Acaciæ, ℥ss.
 Olei Menthæ sativæ, gtt. ij.
 Tincturæ Opii, gtt. x.
 Aquæ æratæ, ℥iij.

Fiat solutio.—Signa.—A table-spoonful for a dose, *pro re nata*. In sickness of the stomach.—*Dr. Meigs*.

Mixture of Colombo, &c.

R. Radicis Colombæ, ℥ss.
 Coque in aqua, ℥iij ad ℥v.
 Cola et adde—
 Carbonatis Potassæ, ℥ss.
 Succi limonis, ℥iij.
 Tincturæ Opii, gtt. xij.

Misce.—Signa.—From a tea to a table-spoonful may be given every hour.

This mixture is said to be powerfully *anti-emetic*; and is much praised by *Dr. Meigs*.

Anti-Emetic Mixture.

R. Tincturæ corticis Aurantii, ℥ss.
 ——— Aloes socotrinæ,
 ——— Castorei, āā. ℥j.

Misce.—Signa.—The dose may be from forty to fifty drops, three or four times a-day.

In chronic vomiting, occasioned by atony or spasm of the stomach, the above is strongly recommended in *Hufeland's Treatise on Scrofula*.

Mixture with Carbonate of Soda, &c.

R. Sodæ carbonatis, ℥iv.

Infusionis Gentianæ compositæ,

Aquæ cinnamomi, āā. ℥iij.

Tincturæ Cardamomi, ℥ss.

Fiat mistura.—**Signa.**—A table-spoonful every two or three hours.

In acidity and flatulency of the stomach, &c.

APPENDIX, NO. I.

DIETETIC PREPARATIONS.

THE importance of an acquaintance with those preparations of diet peculiarly designed for the patient during disease and convalescence, is so obvious that it requires little or no comment. It was remarked by Dr. Rush, in his lectures, that a physician should spend six months in a kitchen before entering upon his practical career. As the medical attendant, therefore, is called upon frequently to give directions and rules for preparing the diet for the sick, we have inserted a few formulæ for those articles which are in daily use. However apparently trifling such a class may appear to some persons, the physician will find that an attention to it will place the patient more completely under his own control, and inspire confidence in the relations and attendants. The rules to be observed in giving nourishment to the sick, will generally, present themselves; but there is one in particular which should not be overlooked. In all low diseases, the nurse should be directed never to give too much at once, but frequently to repeat a small proportion. The lapse of a *very* short time without food, will frequently in cases of great prostration, reduce the patient very low, *and may endanger* his life.

This rule is not intended to be applied to those

chronic derangements of the digestive functions, which are to be remedied principally by a regulation of diet, and in which, the old adage, "a little and often," has too frequently, we are convinced, been perniciously advised.

Decoction of Barley; or, Barley Water.

Take of

Pearl Barley,	-	-	-	-	-	2 ounces,
Boiling Water,	-	-	-	-	-	2 quarts.

Before adding the boiling water, let the barley be well washed. Then boil it to one-half, and strain the liquor. A little lemon juice and sugar may be added, if desirable. To be taken *ad libitum* in inflammatory diseases.

Rice Water.

Take of

Rice,	-	-	-	-	-	2 ounces,
-------	---	---	---	---	---	-----------

Let it be well washed, and add to it—

Water,	-	-	-	-	-	2 quarts.
--------	---	---	---	---	---	-----------

Boil it for an hour and a half, and then add sugar and nutmeg as much as may be required. To be taken *ad libitum*.

Rice, when boiled for a considerable time, assumes a gelatinous form, and is a very excellent diet, mixed with milk, for children. It is believed by many persons to have a constipating property. This opinion, however, appears to be destitute of foundation.

Infusion of Flaxseed.

As under Expectorants.

Mucilage of Gum Arabic.

As under Expectorants.

Decoction of Bran.

Take of

Wheat Bran, recent,	-	-	-	-	-	1 pint,
Water,	-	-	-	-	-	3 quarts.

Boil down one-third, strain off the liquor, and add sugar, honey or molasses, according to the taste of the patient.

Bran tea may be made by using boiling water, and suffering the mixture to stand in a covered vessel for three or four hours.

Sage Tea.

Take of

The dried leaves of sage,	-	-	half an ounce,
Boiling Water,	-	-	1 quart.

Infuse for half an hour, and then strain. Sugar and lemon juice may be added in the proportion required by the patient. In the same manner may be made balm and other teas.

These infusions form very agreeable and useful drinks in fever, and their diaphoretic powers may be increased by the addition of sweet spirits of nitre; or, what is better, by antimonial wine.

Infusion of Malt.

Take of

Ground Malt,	-	-	-	-	1 pint,
Scalding Water,*	-	-	-	-	3 pints.

Infuse for two hours, and strain off the liquor, to which may be added sugar or lemon juice, if required.

This is a favourite preparation with Dr. Parrish, and employed by him in inflammatory fevers.

Lemonade.

Take of

Fresh Lemon Juice,	-	-	-	-	4 ounces,
Fresh and very thin peel of Lemon,	-	-	-	-	half an ounce,
White Sugar,	-	-	-	-	4 ounces,
Boiling Water,	-	-	-	-	3 pints.

* Water not quite brought to the boiling point.

Let them stand until cold, then strain off for use. When employed in fevers, a little nitrate of potash or sweet spirits of nitre may be added.

This preparation, when freely taken by the patient, sometimes produces pain in the bowels. It must therefore be given with some reserve as a daily drink.

Water Gruel.

Take of

Oatmeal,	-	-	-	-	-	2 ounces,
Water,	-	-	-	-	-	1 quart.

Boil for ten or fifteen minutes, and strain, adding sugar and salt to make it agreeable.

Milk for Infants.

Take of

Cow's Milk,	-	-	-	-	-	1 part,
Water,	-	-	-	-	-	2 parts.

Loaf Sugar as much as may be agreeable.

It is necessary, when children are to be raised by the hand, to dilute the milk. These proportions may be altered as the child advances in age. The object is to make a diet as near the qualities of the maternal milk as possible.

Rice Gruel.

Take of

Ground Rice,	-	-	-	-	-	1 ounce,
Cinnamon,	-	-	-	-	-	1 drachm,
Water,	-	-	-	-	-	1 quart.

Boil for forty minutes, adding the aromatic near the conclusion. Strain and sweeten it. Wine may be added if necessary.

Panado.

Take of

Wheat Bread,	-	-	-	-	-	1 ounce,
Cinnamon,	-	-	-	-	-	1 drachm,
Water,	-	-	-	-	-	1 pint.

Boil them until well mixed, then add a little grated nutmeg and sugar.

Wine or butter may also be added if desirable.

Compound Salep Powders.

R. Salep,

Gummi Tragacanthi,

Sago, āā. ℥iv.

Cochineal, ℥ss.

Testæ ostrearum preparatæ, ℥j.

To be carefully mixed and divided into powders of ℥j each. Stir one of these powders into a pint of milk, and boil them together for ten or fifteen minutes. To be drank freely in dysentery and diarrhœa.

These are the dysenteric powders of Dr. Castanello. A somewhat analogous preparation which bears the same name, is prepared as follows:

R. Pulveris gummi arabici,

Gummi Tragacanthi,

Marantæ,

Sago,

Tapiocæ, āā. ℥ij.

Mix them well together, and boil in a pint of milk, flavoured with nutmeg or cinnamon. To be used as a diet in dysentery, diarrhœa, &c. &c.

Sago.

Take of

Sago,	-	-	-	-	-	2 spoonsful,
-------	---	---	---	---	---	--------------

Water,	-	-	-	-	-	1 pint.
--------	---	---	---	---	---	---------

Boil gently until it thickens, frequently stirring. Wine, sugar, and nutmeg, may be added according to circumstances.

Tapioca Jelly.

Take of

Tapioca,	-	-	-	-	-	-	2 spoonsful,
----------	---	---	---	---	---	---	--------------

Water,	-	-	-	-	-	-	1 pint.
--------	---	---	---	---	---	---	---------

Boil it gently for an hour, or until it assume a jelly-like appearance. Add sugar, wine, and nutmeg, with lemon juice, to suit the taste of the patient and the nature of the case.

Arrow Root.

Take of

Arrow Root,	-	-	-	-	-	1 table-spoonful,
Sweet Milk,	-	-	-	-	-	half a pint,
Boiling Water,	-	-	-	-	-	half a pint.

Boil them a few moments over the fire.

Sweetened with loaf sugar it is one of the best aliments for children, and more especially when their bowels are irritated.

Boiled Flour.

Take of

Fine Flour, - - - - - a pound.

Tie it up in a linen cloth as tight as possible, and after frequently dipping it in cold water, dredge the outside with flour till a crust is formed round it, which will prevent the water soaking into it while boiling. It is then boiled until it becomes a hard, dry mass.

Two or three spoonful of this may be grated and prepared in the same manner as arrow root, for which it forms an excellent substitute, and can be obtained in the country, when perhaps the other cannot.

Vegetable Soup.

Take one turnip, one potato, one onion. Let them be sliced, and boiled in one quart of water for an hour. Add as much salt as is agreeable, and pour the whole upon a piece of dry toast.

This preparation forms an agreeable substitute for animal food, and may be given when the former is inadmissible.

Chocolate.

When this is desired by the patient, it should be suffered to stand until cold, after having been made in the usual way. The oily parts thus collect on the surface and should be taken off. The liquid may then be boiled again, and sugar, &c. added as much as may be agreeable.

Beef Tea.

Take of

Lean Beef, cut into shreds,	-	-	-	-	1 pound,
Water,	-	-	-	-	1 quart.

Boil it for twenty minutes, taking off the scum as it rises. After it grows cold, strain the liquor.

This preparation is more nourishing than ordinary broths, and is very palatable.

Hartshorn Jelly.

Take of

Hartshorn shavings,	-	-	-	-	4 ounces,
Water,	-	-	-	-	1 quart.

Boil over a gentle fire until one pint of the water be dissipated. Strain and add lemon juice, sugar and wine.

This forms, either with or without the last ingredients, a very nourishing diet for the convalescent; and when mixed with an equal portion of cow's milk, is very useful in the bowel complaints of children.

Isinglass, or fish glue, may be used instead of the hartshorn shavings, if preferred.

Calves' Feet Jelly.

Take two calves' feet, and add to them one gallon of water, which reduce by boiling to one quart. Strain it, and when cold skim the fat entirely off. Add to this, the white of six or eight eggs well beaten, a pint of wine, half a pound of loaf sugar, and the juice of four lemons, and let them be well mixed. Boil the whole for a few minutes, stirring it constantly, and then pass it through a flannel strainer.

This forms a very nutritious article of diet for the sick and convalescent. When desired the wine can be omitted.

Essence of Beef.

Take of

Lean beef, sliced.

Put a sufficient quantity into a porter bottle to fill up its body, cork it loosely and place it in a pot of cold water, attaching the neck by means of a string to the handle of the vessel. Boil this for three-quarters of an hour, then decant the liquor and skim it.

To this preparation may be added spices, salt, wine, brandy, &c., according to the taste of the patient and nature of the disease. It is greatly employed in low fevers, and therefore every one should be able to direct its *modus præparandi*.

Chicken Water.

Take half a chicken divested of all fat, and break the bones; add to this half a gallon of water, and boil for fifteen or twenty minutes. Season with salt.

This water has been freely employed by some physicians in the early stages of cholera. Taken warm it promotes vomiting and washes out the stomach.

Suet Plisan.

Take of

Sheep's Suet,	-	-	-	-	-	-	2 ounces,
Milk,	-	-	-	-	-	-	1 pint,
Starch,	-	-	-	-	-	-	half an ounce.

Boil slowly for ten minutes.

This may be used as a common drink in dysentery.

Mucilage of Starch, &c.

Take of

Starch,	-	-	-	-	-	-	1 ounce,
Powdered Cinnamon,	-	-	-	-	-	-	1 drachm,
———— Gum Arabic,	-	-	-	-	-	-	1 ounce,
Boiling Water,	-	-	-	-	-	-	3 pints.

Boil until reduced one-third, and strain.

The above may be used as a common drink in dysentery.

Wine and Mustard Wheys.

As under Stimulants.

Rennet Whey.

Take of

New Milk,	-	-	-	-	-	-	-	1 quart,
Rennet,	-	-	-	-	-	-	-	a large spoonful.

Warm the milk and then add the rennet. Boil until the curd separates and is all taken off.

To many persons this forms a very agreeable nutriment.

Vinegar Whey.

Take of

Milk,	-	-	-	-	-	-	-	1 pint,
Vinegar,	-	-	-	-	-	-	-	half an ounce.

Boil for a few minutes, and separate the curd.

Tartar Whey.

Take of

Milk,	-	-	-	-	-	-	-	1 quart,
Cream of Tartar,	-	-	-	-	-	-	-	half an ounce.

Boil until the curd separates.

Sarsaparilla Beer.

R. Sarsaparillæ contusæ, ℥ij.

Pulveris corticis Guaiaci, ℥viiij.

Ligni Guaiaci concisi,

Seminum Anisi,

Radicis Glycyrrhizæ contusæ, āā. ℥iv.

Corticis radiceis Mezereii, ℥ij.

Sacchari liquidi, ℥ij.

Caryophilli contusi, ℥ss.

Aquæ bullientis, congios iv.

Mix these ingredients, and agitate the vessel thrice a-day. When a fermentation has well begun it is fit for use, and may be taken in the dose of a small tumblerful twice or thrice a-day.

“This is the Spanish recipe for the Jarave or diet drink, somewhat modified. If intended for old and obstinate complaints, as leprous affections, elephantiasis, various anomalous ulcerations and foul disorders of the skin, there may be added to the jug, *after fermentation has well begun*, tartrate of antimony twelve grains, corrosive sublimate eight or ten grains, muriate of ammonia one drachm, dissolved previously in a little water. These remedies greatly increase the alterative powers of the vegetable infusion, and so effectually prevent its decomposition that it may be kept for a long time unaltered, *even in a hot climate*.”—*See Hancock on Sarsaparilla, Journal of the Philadelphia College of Pharmacy, Vol. I. No. 4.*

APPENDIX, NO. II.

POISONS.

CASES of poisoning are so sudden and so awful that no time is to be lost in administering relief. To do this with judgment and effect, it is requisite to be familiar with all the usual antidotes: of these we have accordingly selected some which possess an acknowledged efficacy, omitting at the same time many others that are unworthy of confidence. For the purpose of convenience we have arranged the *Poisons* alphabetically, with the appropriate *antidotes* attached to each. But, as has been remarked when speaking of Emetics, our chief reliance in these cases is upon free vomiting, induced by those articles which are most speedy in their effect. Dr. Beck prefers sulphate of zinc, whenever it can be procured, and adds, "but if vomiting be present, we may aid it by diluents or a vegetable emetic. Tartrate of antimony should never be given." Yet even from this treatment little can be expected where the deleterious substance has remained long in the stomach; and the stomach tube and syringe will be indispensable.

To protect the stomach and intestines from the direct operation of poisons, it is generally recommended to give milk, lime water, soap, or drinks sweetened with sugar or honey. Oils and other fatty substances are rather injurious than otherwise.

When the vomiting has ceased, and retching and pain continue, Mr. Hume advises the free administration of carbonate of magnesia, with tincture of opium, suspended in water.

A.

ACIDS, NITRIC, MURIATIC, AND SULPHURIC.

The proper antidotes to these are the alkalies, alkaline earths and soaps; common pearl ash, or carbonate of potash; calcined, or carbonate of magnesia; chalk; soap in solution; accompanied with copious draughts of warm water or warm flaxseed tea. While the antidotes are in preparation, use freely any mild fluid, milk or oleaginous matters being preferred.

ALKALIES.

The best antidote is common vinegar; it is always at hand, and its affinity for these substances is sufficiently strong to neutralize them. Also, a *fixed oil*, as olive oil should be given in *large quantity*—several pounds being required where a large portion of the *alkali* has been taken; it acts by rendering the vomiting easy and free, and partly by converting the alkali into a soap.

ANTIMONY.

The preparation of this mineral by which poisoning is commonly produced, is *Emetic Tartar*. This of itself produces violent vomiting, and therefore our first efforts should be directed to keeping up the emesis by means of mild diluents and mucilaginous preparations. Its antidotes are the alkalies, bitter and astringent infusions, such as common tea, Peruvian bark, the Yellow Bark and its tincture—the administration

of the bark in *powder*, before the decoction is ready, &c. &c. These, and especially the bark, decompose the antimonial salt. Where the vomiting is excessive, opium will often assist in allaying it.

ARSENIC.

Until lately there has been no known antidote to this deadly poison, and our sole dependence has been upon mild diluents, albumen, (such as the whites of eggs suspended in milk) or warm water and sugar. Various substances have been proposed as antidotes, but without success. It is best first (if vomiting has been delayed) to give an emetic of the Sulphate of Lime, and milk should be drunk liberally both before and after the vomiting has begun. The patient should never be allowed to exhaust his strength in retching, without some milk or other fluid in his stomach to act on.

THE HYDRATED PEROXIDE OF IRON, *freshly* prepared, was announced in the year 1834, by Dr. Bunsen of Gottingen, as an antidote for poisoning by Arsenic; and its efficacy has been confirmed by M. M. Orfila, Lesueur, Soubiaïn and Miguel, and also in this city by various chemists, and by experiments upon the inferior animals. The antidote is thus prepared.

Hydrated Peroxide of Iron.

R. Sulphuric acid (67° Baumé)	8 oz.	-	-	16 parts.
Iron wire	-	8 oz.	-	16 “
Nitric acid - (49° Baumé)	5½ oz.	-	-	11 “
Water of Ammonia	-	q. s.	-	-
Water	-	1½ gal.	-	384 “

Mix the Sulphuric Acid with the water in a glass vessel; then add the iron, and after the effervescence has ceased, filter. Add the Nitric Acid in divided portions.

and apply heat so long as orange-coloured fumes are given off. To the heated solution, pour in the Water of Ammonia until a decided excess has been added, then wash the precipitate by decantation, until the washings give no precipitate with Nitrate of Baryta. The water is then to be drawn off until just enough remains to give the consistence of thick cream.

Bottles of half a pint are the most convenient size for containing it; to each bottle add two ounces of un-crystallizable sugar or honey to prevent the oxide from separating from the water.

The following direction will enable the most ignorant to use this antidote until medical advice can be obtained. "This antidote must be administered AS SOON AS POSSIBLE after the discovery that arsenic has been taken, and as it causes no bad effects itself, it should be given EVERY FIVE OR TEN MINUTES, UNTIL ENTIRE RELIEF IS OBTAINED. The dose for a grown person is a table-spoonful; for children a tea-spoonful. The bottle must be well shaken before each dose.

Should the patient survive (and he will, if the antidote prepared as above be faithfully given) the inflammatory symptoms must be combated on the general principles for treating inflammations. The diet should consist exclusively of milk, gruel, cream, rice and similar bland articles.

In a case of poisoning in this city, where *half an ounce* of the *powdered arsenic* had been taken and which caused extreme thirst and vomiting, all the symptoms were allayed immediately after freely taking the *Hydrated peroxide of Iron*. Only a slight attack of Gastritis followed, which was subdued by the usual treatment, cupping, &c. &c.

B.

BARYTES.

The carbonate of this earth is poisonous, and might possibly be taken by mistake. Its true antidote is sul-

phate of magnesia or soda; with these a sulphate of barytes is formed, which is perfectly insoluble, and may be brought off by emetics and copious diluents, oleaginous or albuminous.

C.

COPPER.

The sulphate of this metal is, perhaps, the only chemical preparation that is likely to be taken into the stomach. A more ordinary manner is the careless use of culinary vessels, or mineral water fountains. All saccharine preparations, and the prussiates of potash and iron are considered as antidotes. But our safest reliance is upon *albumen*, with copious draughts of warm milk, and *brown* sugar or molasses. Vinegar in all cases should be avoided.

G.

GOLD.

The nitro-muriate of gold is a powerful poison. Its antidotes are sulphate of iron, (which throws down the gold in a metallic state,) and the free use of mucilaginous drinks.

L.

LAUDANUM.

Laudanum, when taken in too large quantity, must be withdrawn from the stomach by active emetics, or by tickling the fauces. The stomach-tube may be used to remove whatever remains of the poison. Diluted vegetable acids, and vinegar in particular, are then necessary, and active stimulants are also often

required. Iodine, bromine and chlorine are stated by M. Donné, of Paris, to be antidotes for poisoning by nux vomica and opium. M. Orfila thinks that the decoction of galls may with propriety be used as an imperfect antidote, until the narcotic can be evacuated from the stomach.

LEAD.

The acetate is the only preparation which is resorted to as a poison. The sulphate of magnesia or soda is its true antidote, as it converts the poison into a sulphate, which is inert;—the alkaline carbonates and particularly the bi-carbonates. Phosphate of soda is also a good antidote. If the patient does not vomit, it is best to give an emetic of the sulphate of zinc.

M.

MERCURY.

All the preparations of this metal are poisonous, and corrosive sublimate in particular. Our chief and only dependence, according to M. Orfila, is upon the albumen of eggs, copiously swallowed, or introduced through the *stomach tube*, which should be resorted to on all occasions where the patient cannot swallow or vomit. Enemata of the same should also be employed.

If the albumen be taken in sufficient quantities, it decomposes the sublimate, forming a triple compound of albumen, muriatic acid and calomel. If the remedy excite fresh vomiting, so much the better, and if symptoms of gastritis are present, blood-letting should not be overlooked.

The gluten of wheat flour is also strongly recommended, on the same principle, as an antidote to corrosive sublimate. To administer it, it is only requisite to mix the flour with water. The quantity taken must be

very considerable. This remedy reduces the muriate to the state of calomel.

Gluten of wheat flour is best given in the form of an emulsion with soft soap—thus: by mixing partly in a mortar and partly with the hand, 5 or 6 parts of fresh gluten with 50 parts of soft soap.

When neither albuinen or gluten is at hand, milk is a good substitute.

N.

NARCOTICS.

The remedies employed in all cases of poisoning from this class of substances, are nearly the same, viz: the most active emetics, as sulphate of zinc in large doses, sulphate of copper and ipecacuanha. In all cases where it is possible, the stomach tube should be employed to wash out that organ. After these evacuations, coffee and the vegetable acids are proper.

NITRE.

This salt is often taken in mistake for Epsom or Glauber's salts. When this happens, vomiting should be excited as speedily as possible, and be followed by the free use of mucilaginous drinks.

O.

OPIUM.

We cannot do better in this place than transcribe what Dr. T. R. Beck, of Albany, says of the antidotes to opium.

“The experiments of Orfila have demonstrated that

the administration of vinegar, or other vegetable acids, previous to the evacuation of the poison by vomiting, will accelerate and aggravate the action of the opium; but that when the opium has previously been expelled, water acidulated with vinegar, or any other vegetable acid, will tend to diminish and correct the effects. Coffee, when prepared in strong decoction or infusion, rapidly lessens the symptoms, but cannot be considered as an antidote. The result of his experiments led him to give the following directions for the treatment of a person poisoned by opium. Induce vomiting, if possible, with sulphate of zinc, or sulphate of copper: but the administration of watery liquors with the view of promoting the vomiting, should be forbidden, as they dissolve the opium and promote its absorption. In order to ensure the action of the emetic, the patient must be kept *roused* either by discharging cold water over the head and breast, yet cold affusion must be avoided in the advanced stage, when the body is cold, and the breathing scarcely perceptible. To ensure success in every case, the patient must be kept awake, either by dragging him up and down between two men, or by flagellation on the back and soles of the feet. Artificial respiration may be used in desperate cases with propriety. Bleed the patient immediately after the rejection of the poison, and repeat this if necessary. Afterwards administer alternately water acidulated with any vegetable acid, and a strong infusion of coffee warmed—and in ten or twelve hours exhibit an enema, and let the arms and legs of the patient be well rubbed. Purgative clysters should be continued, if it be suspected that any of the opium still remains in the large intestines.”—*Dr. Beck's Med. Jurisprudence, Vol. II. p. 403.*

OXALIC ACID.

Death is generally sudden when this poison has been taken. A mixture of chalk and water is recommended

by Dr. Thompson, for the purpose of converting the acid into oxalate of lime. The alkalies must be avoided, as they act with an energy very little inferior to the acid itself.

It is particularly necessary to avoid giving *warm water* with a view to accelerate vomiting; for dilution will promote the entrance of the poison into the blood, if it has not the effect of *immediately* expelling it.

S.

SILVER.

The nitrate of this metal is sometimes taken by mistake: its antidote is common salt, *largely given*, with copious draughts of mucilaginous drinks to promote vomiting. The salt produces an insoluble muriate which has no power on the system. *Beck.*

T.

TIN.

When any of the salts of tin are taken, albumen of eggs, or milk given freely, is the appropriate remedy. Milk forms a coagulum with the muriate of tin, and renders it innoxious.

Z.

ZINC.

The sulphate of this metal is occasionally taken in an over-dose. Warm water and emollient drinks are proper in such cases; but milk is particularly useful, from its power of decomposing the sulphate. Also albumen.

The crystallized sulphate of zinc very closely resembles the sulphate of magnesia, and *has been mistaken* for it. This fact should be borne in mind by apothecaries and physicians.

The following works have been referred to in forming this Chapter on Poisons. A Treatise on Poisons, by Robert Christison, M. D. Turner's Chemistry. Manuel de Médecine Legale, par A. Brierre de Boismont. Orfila, Toxicol. Generale. Henry, Thompson, &c. &c.

APPENDIX, NO. III.

ON THE ENDERMIC USE OF
MEDICINES.

"THE science of medicine," says Dr. Gerhard,* "has recently received an important auxiliary to its resources, by the discovery that remedial substances when applied to the cutis, after the removal of the epidermis, produce their peculiar effects with nearly the same energy as if introduced into the alimentary canal. This mode of exhibition, which has received the name of the *endermic method*, or *endermic medication*, is always found much more efficacious than the iatatreptic method of Christien, or the administration of medicines by means of frictions on the cuticle."

The endermic method requires the previous separation of the cuticle. This is best effected by a blister about four inches square. A small portion of the cuticle should be removed, and the medicine inserted into the opening so that it may come in contact with the vesicated surface. Substances used in this way, must be applied in two or three times the quantity requisite when given by the mouth. "The remains of the substance applied are much less than the original quantity, and sometimes no trace of it can be perceived; the whole

* North Amer. Med. and Surg. Journ. April, 1830.

appearing to be absorbed or mingled with the secretions."

We avail ourselves of Dr. Gerhard's valuable paper for the following facts in relation to those medicines which act most effectually in the endermic method.

Sulphate of Quinia.—Blister to be applied to the epigastrium during the apyrexia; and if the paroxysm returns before the quinia can be used, defer it until the next apyrexia. Apply four grains of powdered quinia, (either alone or with an equal quantity of starch,) four times a-day, for two successive days. A less irritating plan, however, is to dissolve sixteen grains of quinine in two ounces of water, with which the blistered surface is to be repeatedly wet during the day, by means of a soft linen cloth. There is no occasion to *remove* the loose epidermis; it is all-sufficient to *puncture* it in several places to allow the solution to touch the denuded surface.

Morphia.—The acetate and sulphate act readily and effectually. The sulphate is preferable because more soluble and less irritating. It may be applied in powder, in solution, or in the form of cerate.—The dose varies from half a grain to two grains. Its effects are speedily perceptible.

Digitalis.—Dose, two grains three times a-day.

Aloes.—This medicine acts on the bowels with nearly the same certainty as if taken internally. Dose, ten grains, to be repeated, if necessary, in a few hours.

Gamboge, owing to its solubility, purges freely in the endermic method. Dose from five to eight grains.

Rhubarb, *Jalap*, *Elaterium* and *Croton Oil*, contrary to what might be expected, do not produce catharsis, although they excite severe local irritation.

Squills.—This medicine in powder, sprinkled on a blistered surface three times a-day in the quantity of four grains at each application, causes copious diuresis. Its power is increased by combination with *Digitalis*, in the proportion of four grains of the former to one of the latter.

Emetics from the *vegetable* kingdom all produce their effect in the endermic method. The *metallic* medicines of this class produce merely local irritation.

Mercury.—Every practitioner is well aware of the endermic action of mercury.

Iodine.—This substance acts rapidly: it is best applied in the form of ointment, made by rubbing a drachm with an ounce of cerate.

INDEX.

- A.
 Absorbents, 222.
 Acetate of ammonia, 49, 162.
 of ammonia, embrocation, 107.
 of ammonia and camphor, 206.
 of ammonia and opium, 206.
 of lead with calomel, 201.
 of lead with opium, 201, 202.
 217.
 of lead, epithem of, 218.
 of mercury, 152.
 of morphia, 142, 148.
 of zinc, injection of, 211.
 Acorn coffee, 184.
 Acid, prussic, 80.
 sulphuric, 166.
 nitrous, 198.
 nitro-muriatic, 198.
 nitric, 37, 198.
 pyroligneous, 209.
 linctus of muriatic, 187.
 Acids considered as poisons, 239.
 Aloes with canella, 26, 88.
 with rhubarb and soap, 31.
 with gentian, 31, 186.
 with ipecacuanha, 32.
 with asafætida and soap, 32.
 with blue pill, 33.
 with guaiacum and rhubarb, 34.
 Alcohol with lime water, (lotion,) 191.
 Alkalies considered as poisons, 239.
 Alkaline bath, 124.
 Alteratives, 188.
 Althea collyrium, 208.
 Aloetic enema, 100.
 Almond emulsion with squills, 75.
 Alum, burnt, 124.
 with kino, 200.
 with opium, 201.
 with extract of bark, 202, 216.
 whey, 216.
 curd, 205.
 collyrium of, 205.
 Alum and sulphate of zinc, 21.
 with oak bark, (gargle,) 212, 213.
 with spirits of nitre, 186.
 with calomel, 220.
 Ammoniated counter-irritants, 106.
 Ammonia, nitrate of, 67.
 carbonate of, with valerian, 134.
 acetate of, 48, 64.
 muriate of, with arsenic, &c. 175.
 muriate of, injection, 211.
 Amber oil, 116.
 salt with castor, 156.
 Ammoniac mixture, 84.
 and squills, 77.
 gum, with squills, 72.
 with conium, 72.
 and nitric acid, 80.
 Ammoniuret of copper, 174, 178.
 of copper, injection of, 212.
 Anodynes, 141.
 Anodyne draught, 147.
 pills, 64.
 Angustura bark, 171, 185.
 Antacids, 222.
 Anti-emetic mixture, 226, 18.
 Antilithics and lithontriptics, 56.
 Anthelmintics, 95.
 Antidynous lotions, 106.
 Antispasmodics, 156.
 Antimonial wine, 20.
 wine with nitre, &c. 64.
 wine with water of ammonia, 64.
 wine with tincture of tolu, 64.
 powder, 62.
 powder with calomel, &c. 63.
 Antimony, antidotes of, 239.

Aromatic Pills, 133.
 sulphur ointment, 221.
Arsenic, white oxide of, 124. 197.
 ointment of, 124.
Arsenic, pills of, 175.
 antidotes to, 240.
Arsenical solution, 182.
Arsenious acid, 124.
Arsenical powder, 125.
Arsenate of iron, 197.
Arrow root, 233.
Asafœtida with magnesia, 35.
 with aloe, 32.
 with tolu, &c. 77.
 milk of, with aq. ammoniæ,
 78.
 with soap, 156.
 with castor, 158.
 mixture, 158.
 with valerian, &c. 158.
 enema, 159.
Astringents, 200.

B.

Baths, 119.
 Barèges, 120.
 sulphureo-gelatinous, 120.
 nitro-muriatic acid, 120.
 emollient, 120.
 sublimate, 121.
 alkaline, 121.
 artificial Harrogate, 121.
Barytes, muriate of, 69.
 antidotes of, 241.
Balsam copaiva, mixture of, 51. 54.
 copaiva with cubebs, 53.
 copaiva with calomel, 54.
 copaiva with Bolus, 54.
 copaiva with tolu, 78.
 copaiva in pills, 46.
 Peruvian, 210. 79.
Barley water, 229.
 with muriatic acid, 56.
 with nitre, 69.
Belladonna, 80. 112. 152. 153.
Beef tea, 234.
 essence of, 235.
Bismuth with cubebs, 134.
 pills, 175.
Blake's toothach solution, 213.
Blister issue, 129.
Blisters, 102.
Blue pill, 164.
Blue pill with jalap and aloe, 29.
 with rhubarb, 29.
 with colocynth, 32.
 with aloe, 33.

Black wash, 127.
 drop, 151.
Blood root, 78.
Boneset, 68. 179.
Boiled flour, 233.
Boracic acid, 159.
Borage flowers with sarsaparilla, 66.
Bougies, 130.
 Sharp's, 130.
 Dr. Swediaur's, 130.
Bole with sulph. copper, 211.
 with sulph. zinc. (injection,) 211.
Borax, (gargle,) 213.
Brown mixture, 74.
Bran, decoction of, 229.
Brucia, pills of, 194.
 tincture of, 194.
Bromine, 195.
Bromide of potassium, 195.
 of iron, 195.
Butternut, extract of, 32.
Burdock in decoction, 66.
Burnt sponge, 193.
Burnt alum, 124.

C.

Cathartics, 24.
Calomel with jalap, 25.
 with jalap and rhubarb, 26.
 with rhubarb, 26.
 with rhubarb and opium, 30.
 with extract of colocynth, 33.
 with squills, 47. 72. 73.
 with squills and digitalis, 47.
 with nitre and tartar emetic,
 60.
 with ipecacuanha, nitre, &c.
 60.
 with opium, 61.
 with opium and antim. powder,
 61. 62.
 with gamboge, 95.
 with pink root, 95.
 with lime water, 157.
 with iodine, 108.
 pills, 151.
 with acetate of lead, 201.
 with laudanum, (injection,) 184.
 with alum, ointment of, 220.
 with quinia and opium, 196.
 with quinia and tart. antimon., 220.
Calves' feet jelly, 234.
Camphor with emetic tartar, &c. 63.
 with guaiacum, 138.
 mixture, 92. 136.

- Camphor** julep, 123.
 enema, 100, 135.
 and mercury, 104.
 liniment, 113.
 with cantharides, 132.
 in milk, 136.
 with magnesia, 136.
 with myrrh, 137.
 with musk, 157.
 with acetate of ammonia, 65.
 with opium, 204.
 with Hoffman's anodyne, 150.
 cinchona, 172.
 with nitric acid, &c. 203.
 water with laudanum, &c. 204.
 with magnesia, 225.
Canella with aloe, 26.
Carbonate of soda with ipecacuanha, 76.
 of soda with rhubarb, 223.
 of soda with magnesia, 223.
 of soda with gentian, 227.
 of potash with cochineal, 76.
 of potash, 87, 224.
 of potash with opium, &c. 224, 226.
 of iron, 96, 157, 172.
Castor oil, 34.
 with spirits of turpentine, 35.
 enema of, 42.
Catechu and alum, 202.
 injection of, 210.
 electuary, 204.
 mixture, 216.
 with logwood, 216.
Castor with amber, 156.
 with asafetida, 158.
Cantharides, Dewees's tincture of, 92.
 with spirits of turpentine, 111.
 liniment of, 113.
 with camphor, 133.
Cataplasma, antiseptic, 119.
 of pepper, &c. 116.
 of mustard, 117.
 of garlic, 117.
 of yeast, 118.
 of carbon, 118.
 of iodine, 117.
 of salt, 118.
 of yeast, 118.
 antiseptic, 119.
 resolvent, 119.
Caustics, 123.
Caustic potash, 123.
 lunar, 123.
Cayenne pepper and alcohol, 116.
Cayenne pepper infusion, 130.
 in pills, 132.
Charcoal, 29.
Chicken water, 235.
Chamomile infusion, 185.
Chiococca anguifuga, 57.
Chocolate, 233.
Chalk with opium, 222.
 with rhubarb, 222.
 mixture, 203, 224.
Cherry tree bark, 173.
Chloride of soda, lotion of, 218.
Chlorine fumigation, 87, 122.
Cider with parsley, &c., 48.
Cinchona, 170.
 with uva ursi, 55.
 with serpentaria, 171.
 with sulph. magnesia, 171.
 with cloves, 171.
 with camphor, 172.
 with Epsom salts, 159.
 extract of, with gentian, 173.
 electuary, 177.
 tincture of, 183.
 decoction of, 179, 181.
 in Port wine, 181.
 with valerian, 184.
 extract of, enema of, 187.
 with lime water, 216.
 extract of, 182.
 lotion, 219.
Cinchonia, 173.
Cinchonina, 181.
Citrate of ammonia, 67.
 of potash, 63.
Cicuta. See Conium.
Cloves, 118.
Copaiva. See Balsam.
Colchicum wine 39, 52.
 with Epsom salts, 40.
 with magnesia, 39.
 with elaterium, &c., 41.
 with squills and tobacco, 50.
Cochineal with common salt, 96.
 with carb. potash, 76.
Confection of sulphur, 32.
Conium with tolu, paregoric, &c., 75.
 with ipecacuanha, 71.
 with hyoscyamus, 143.
 extract of, 144.
 with calomel, 145.
 with squills, 72.
 infusion of, 152.
Collyrium, opiate, 205.
 of alum, &c., 205.
 of vinegar, 206.
 of acetate of ammonia and camphor, 206.

- Collyrium, of acetate of ammonia and opium, 206.
 of sulphate of copper, 207.
 of corrosive sublimate, 207.
 of opium and camphor, 207.
 of poppies, 207.
 of nitrate of silver, 208.
 of sugar, 205.
 of mercury, 205.
 of dry opium, 205.
 of acetate of zinc, 206.
 of acetate of lead, 206.
 of ioduret of potassium, 206.
 of strychnine, 208.
 emollient, 208.
 anodyne, 208.
 of colchicum, 208.
- Colocynth with calomel, 33.
 with antimony, &c., 33, 34.
- Confection of sulphur, 33.
 of opium, 160.
- Corrosive sublimate and lime water, 126.
 pills, 161.
 with hemlock, 162.
 mixture of, 165.
 solution of, 165, 166.
 collyrium of, 207.
 injection of, 210.
- Cowhage electuary, 97.
- Coxe's hive sirup, 23, 75.
- Copper, sulphate of, 175.
 ammoniuret of 174, 178.
 sulph. of, with bole, 211.
 sulphate, collyrium of, 207.
 antidotes of, 242.
- Cough lozenges, 73.
 mixture, 74.
- Columbo with ginger, 172, 183.
 with rhubarb, 172, 183.
 with magnesia, 223.
 mixture of, 226.
- Croton oil, pills of, 31.
 mixture of, 36.
 with potash, 36.
- Cream of tartar in solution, 48.
 with juniper berries, 49.
 and sulphur, 27.
- Creosote liniment, 115.
 ointment, 112.
 pills, 202.
 mixture, 203.
 alcoholic solution of, 203.
- Crabs' claws with rhubarb, 195.
- Cretaceous powder and opium, 222.
- Cubebs with copaiva, 53.
 with ergot, 47.
- Cubebs lozenges, 75.
 with bismuth, 134.
 with carb. of soda, 134.
- Cyanide of potassium, 83.
 of mercury, 163.
- D.
- Dandelion, infusion of, 55.
 pills, 47.
 with blue mass, 47.
- Denarcotized laudanum, 150.
- Decoction of burdock, 67.
 of cainca, 51.
 of pipsisseway, 55.
 of madder, 91.
 of cantharides, 111.
 of mezereon, 67.
 of lobelia, 67.
 of sarsaparilla, 68.
 of barley, 229.
 of barley, with nitre, 69.
 of Iceland moss, 81.
 of seneka root, 84, 90.
 of aloes, 39.
 of Geoffrea bark, 98.
 of pomegranate, 100.
 of turpentine and cantharides, 111.
 of sponge, 193.
 of bitter sweet, 152.
 of stramonium, in milk, 153.
 of cinchona 179.
 of cinchona with valerian, 184.
 of Augustura bark, 185.
 of wild cherry bark, 185.
 of dogwood, 186.
 of bran, 229.
- Dewees's tincture of guaiacum, 92.
 tincture of cantharides, 94.
 carminative, 35.
- Deuto-iodide of mercury, 132, 168.
- Dietetic preparations, 228.
- Digitalis with opium and calomel, 46.
 with squills and colomel, 47.
 and acetate of potash, 51.
 tincture of, with opium, &c., 77.
 with opium, 144.
 tincture of, 151.
 infusion and tincture of, 50.
- Diaphoretics, 59.
- Diuretics, 45.
- Diuretic drops of Ferriar, 49.
 liniment 51.
- Doses of laudanum, 147.
- Dover's powders, 61.

Dogwood bark, infusion of, 186.
 Dulcamara, decoction of, 152,
 Dupuytren's antispasmodic pills, 162.
 Dyspeptic lie, 225.

E.

Effervescing draught, 64.
 Elaterium with nitre and squills, 41.
 Electuary of senna and figs, 34.
 of Jalap, nitre, &c., 41.
 of sulphur and senna, 42.
 of tin, 97.
 of cowhage, 97.
 of cinchona, iron, &c. 177.
 of catechu, 204.
 Elixir paregoric, 151.
 vitriol with sulph. of iron, 171.
 Embrocations, 114.
 Emetics, 18.
 Emetina lozenges, 21, 74.
 solution of, 21.
 Emmenagogues, 88.
 Emulsion of phosphorus, 137.
 Endermic use of medicines, 248.
 Enema of castor oil, salts, &c., 43.
 of Epsom salts and oil, 43.
 of colocynth and manna, 43.
 of spirit of turpentine, 43.
 of Epsom salts and senna, 44.
 of tartar emetic, 44.
 of tobacco, 44.
 of aloes, 100.
 of camphor, 100.
 of opium, 153.
 of laudanum, 154.
 of asafoetida, 159.
 of musk, 160.
 of extract of bark, &c., 187.
 of quinine, 187.
 of bark and lime water, 216.
 of catechu and lime water, 217.
 of sugar of lead, 217.
 Epsom salts with tartar emetic, 28, 38.
 with senna and manna,
 38, 39.
 with nitric acid, 37.
 with colchicum, 40.
 Epithem of tincture of cantharides, 115.
 of camphor, and carb.
 ammonia, 116.
 of acetate of lead, 218.
 Epispastic ointment, 104.
 Epispastics, 102.
 Ergot, 89, 91.
 and cubebs, 47.
 powder of, 89.
 infusion, 91.

Ergot decoction, 91.
 sirup of, 91.
 Essence of beef, 235.
 Escharotics, 123.
 Ether with iodine, 128.
 with laudanum, 159.
 Epectorants, 70.
 Extract of hyoscyamus, 152.
 of stramonium, 143.
 of conium, 144.
 of belladonna, 152.
 of bark with gentian, 173.
 of quassia and gentian, 173.
 of cinchona, &c., 187.
 of cinchona with alum, 216.
 of logwood with catechu, 188.
 Euphorbia, oil of, 36.

F.

Febrifuge bolus, 174.
 Ferriar's hydragogue cathartic, 55.
 Ferro-cyanate of quinia, 180.
 Fig electuary, 34.
 Flaxseed infusion, 81.
 Fothergill's pills, 33.
 Fowler's solution in mixture, 182.
 Fomentation, 119, 121.
 Fumigation of chlorine, 87, 122.
 of tar, 85.

G.

Gamboge with nitre, &c., 52.
 with jalap, &c., 26.
 with calomel, 95.
 Garlic cataplasm, 117.
 Gargle of galls and alcohol, 212.
 of oak bark, &c., 212.
 of sulphate of zinc, 213.
 of borax, &c., 213.
 of pomegranate, 214.
 of brandy, 212.
 of quinia, 213.
 of muriatic acid, 214.
 detergent, 214.
 mercurial, 215.
 acetous, 215.
 of acetate of lead, 215.
 Galls, infusion of, 215, 217.
 sirup of, 217.
 ointment of, 220.
 Gelatinous bath, 120.
 Gentianin, 188.
 Gentian with aloes, 31, 186.
 with magnesia, 56.

Gentian with quassia, 176.
 with sulphuric acid, 182.
 infusion of, 188.
 Geoffrea bark, 98.
 Gold, nitro-muriate of, 177.
 antidotes of, 242.
 Ginger draught, 136.
 with columbo, 172.
 Goulard's extract, 218.
 Granville's ammoniated lotions, 106.
 Griffith's myrrh mixture, 178.
 Griffiths' pills, 28.
 Gruel, water, 231.
 rice, 231.
 Guaiacum with aloes and rhubarb, 34.
 volatile tincture of, 92.
 with tartar emetic, &c., 65.
 Dewees's tincture of, 92.
 with camphor, 138.
 Gum ammoniac mixture, 83.
 with squills, 77.

II.

Hartshorn jelly, 234.
 Hellebore, tincture of, 92.
 with myrrh, &c., 92.
 with savin, 92.
 Hemlock mixture, 146.
 See Conium.
 Hive sirup, 23, 75.
 Hiera picra, 27, 88, 94.
 tincture of, 94.
 Honey with borax, 213.
 Hoffman's anodyne, 158.
 Hooper's pills, 50.
 Hop, tincture of, 149.
 infusion of, 186.
 Hope's mixture, 204.
 Huxham's tincture of bark, &c., 179.
 Hydrogogue draught, 55.
 Hydrocyanic acid, 81, 82.
 Hydrocyanate of potash, 83.
 Hyoscyamus with ipecac., 143.
 with turpentine, 98.
 with conium, 143.
 pills of, 143.
 infusion of, 142.
 and myrrh, 73.
 Hydriodate of potash, 91.
 ointment of, 109.
 Hydragogue draught, 52.
 Hydrated peroxide of iron, 240.

I & J.

Jalap with cream of tartar, 26, 41.
 with gamboge, 26.

Jalap with extr. of butternut, 32.
 with squills, 49.
 Iceland moss, 80.
 Inhalation, 84.
 Infusion of parsley, 49.
 of juniper berries, &c., 49.
 of digitalis, 50.
 of digitalis with tinc. digit, 50.
 of uva ursi, 52.
 of senega, 74.
 of flaxseed, 81.
 of ergot, 90.
 of pinkroot, 98.
 of wild cherry bark, 185.
 of gentian with magnesia, 56.
 of boneset, 68, 179, 186.
 of pleurisy root, 68.
 of pipsisseway, 69.
 of malt, 230.
 of flaxseed 80.
 of dandelion, 53.
 of pink root, 96.
 of galls, &c., 215.
 of sassafras, 126.
 of logwood, 215.
 of Cayenne pepper, 139.
 of cloves, 139.
 of ginger, 140.
 of conium, 152.
 of hyoscyamus, 152.
 of columbo and ginger, 172.
 of quassia, 184.
 of chamomile, 185.
 of serpentaria, 185.
 of hops, 186.
 of cinchona with lime water,
 187.
 of gentian, 188.
 of seneka with ipecac., &c. 74.
 Injections per anum. See Enemata.
 for gonorrhœa, 209.
 Injection of ammoniuret of copper, 212.
 of sulphate of zinc, 210.
 of corrosive sublimate, 210.
 of same with mur. ammonia,
 211.
 of acetate of zinc, 211.
 of sulphate of zinc and bole,
 211.
 of sulph. of copper, 211.
 of tea, 212.
 for the ear, viz.
 of pyroligneous acid, &c.,
 209, 210.
 of infusion of catechu, 210.
 of Peruvian balsam and Ox-
 gall, 210.
 Iodate of zinc, ointment of, 109.

Iodate of zinc, 109.
Iodo-hydrargyrate of potassium, 191.
Iodine, tincture of, 93. 190.
 ointment, 110.
 liniment, 110.
 ethereal tincture of, 189.
 compound tincture of, 189.
 endermic use of, 250.
 and calomel, 108.
 waters, 191.
 inhalation, 87.
 cataplasm, 117.
Iodide of potassium, 93. 189. 190.
 ointment of, 110.
 of arsenic, 111.
 of sulphur, 111.
 of mercury, 163.
 of iron, 190.
Ioduretted iodide of potassium, 190, 191.
 ether, 191.
 waters, 192.
Ipecacuanha, 18. 22.
 with tartar emetic, 18. 19.
 with rhubarb, 19.
 with squills, 21.
 with aloes, 32.
 with calomel and opium,
 60, 61.
 with guaiacum, nitre, &c.,
 60.
 with myrrh, 70.
 pills, 73.
 with hyoscyamus, 143.
 wine with antimony, 19.
Iron, sulphate of. See Sulphate of
 Iron.
 muriate of, with aloes, 94.
 carbonate of, 96.
 prussiate of, with guaiacum, 173.
 iodide, 190.
 arsenate of, 197.

Issues and setons, 128.

Juniper berries, &c., in cider, 49.
 with cream of tartar, 49.
 oil, 50.

K.

Kentish ointment, 112.
Kino with alum, 200.
 with opium, 204.
Kirkland's neutral cerate, 221.

L.

Lac sulphuris, 28.
 ammoniac, 80.
Lactucarium, 142.

Laudanum, graduated doses of, 147.
 with spirits of turpentine,
 150.
 denarcotized, 150.
 enema, 154.
 with ether, 159.
 antidotes of, 242.

Laffecteur, rob antisiphilitic of, 66.
Lemonade, 230.

Lead, acetate of, with calomel, 177.
 acetate, collyrium of, 206.
 acetate, antidotes of, 243.

Lime water with alcohol, (lotion,) 218.
 with carbonate of potash,
 58.
 with corrosive sublimate,
 126.
 with calomel, 127.
 with cinchona, 216.
 with chatechu, (enema,) 217.
 and zinc, (lotion,) 218.
 and linseed oil, 218.
 with milk, 224.

Liquorice with guaiacum, &c., 59.

Lithontriptics, 56.

Liniment, of ammonia and turpentine,
 105.
 with camphor, 113.
 with soap, 113.
 of oil of amber, 116.
 with ammonia and oil, 113.
 of creosote, 115.
 of oil of amber, 116.
 with ammonia and tartar
 emetic, 106.
 of cantharides, 113.
 of opium, 114.
 of tar, 115.

Linctus, white, 76.
 of spt. of turpentine, 137.

Lozenges, of emetina, 21.
 of liquorice, opium, &c., 73.

Logwood, infusion of, 215, 216.
Lotion of lime water, zinc, &c., 218.
 of same and alcohol, 218.
 of myrrh, 127.
 antipsoric, 127.

Lobelia, 67. 22.
 inflata, 22.
Lugol's iodine ointment, 110.

Lupulin pills, 142.
 tincture of, 149.

Lunar caustic, 123.

M.

Magnesia with rhubarb, 25. 38.

- Magnesia** opium and asafetida, 35.
 with sulphur, 27.
 with Epsom salts, 37.
 with colchicum wine, 52.
 with infusion of gentian, 58.
 with fennel seed, 223.
 with carbonate of soda, 223.
 with columbo, 223.
 with aqua ammonia, 225.
Madder, 89. 91.
Marsh-reed grass, 66.
Male fern, 96.
Malt, infusion of, 230.
Mezercon, 67.
Mercury, acetate of, 162.
 sulphuret of, with serpentina, 62.
 with chalk and ipecacuanha, 164.
 sulphate of, 164.
 red oxide of, 165.
 protoiodide of, 109. 163.
 cyanide of, 112. 162.
 deuto-iodide of, 132.
 cyanuret of, 169.
 antidotes of, 243.
Mercurial plaster, 104.
 ointment with camphor, 166.
 cerate, 112.
 solution, 165.
 gargle, 215.
 pills, 164.
 ointment, 166.
 lotion, 167.
 collyrium, 178.
Mineral acids, 124. 198.
Morphia with tolu, 84.
 sulphate of, 142.
 acetate of, 142.
Mosely's solution, 21.
Monesia, 204.
Moxa, 125.
Mucilage of starch, 235.
Mustard draught, 21.
 cataplasm, 117.
 fomentation, 122.
 seed, 138.
 whey, 139.
Muriatic acid and barley water, 58.
 with nitric acid, 189.
 linctus of, 214.
Muriate of barytes, 69.
 of iron with aloes, &c., 94.
 of iron, tincture of, 183.
 of soda with cochineal, 93.
 of ammonia with arsenic, &c., 175.
 of ammonia with iron, &c. 164.
Muriate of ammonia, injection of, 211.
Muriated tincture of iron, 183.
Mugwort, 125.
Musk and camphor pills, 156.
 with sub-carbonate of ammonia, 157.
 enema, 160.
 mixture, 158.
Myrrh with henbane, 73.
 with sulphate of iron, 89.
 with hellebore, 89.
 with ipecacuanha, 70.
 lotion, 127.
 with camphor, 137.
 mixture, Griffith's, 178.
 with zinc, 71.
- N.
- Narcotics**, 141.
 antidotes of, 244.
Neutral mixture, 62. 63.
 certate, Kirkland's, 221.
Nitric acid with Epsom salts, 37.
 acid diluted, 198.
 lac ammoniac, 80.
 with camphor, 203.
Nitrate of silver, 123.
 of silver, pills of, 177.
 of silver, collyrium of, 208.
 of potash with tartar emetic, 59.
 of potash with guaiacum, &c., 60. 65.
Nitre with barley water, 69.
 antidotes of, 244.
 with calomel and ipecacuanha, 71
Nitro-muriatic acid, 198.
Nitro-muriate of gold, 177.
Nitrous powders, 59. 62.
 acid with laudanum, 167.
Nut galls, infusion of, 215.
Nux vomica powders, 145.
 resin of, 145.
 tincture of, 146.
- O.
- Oak bark gargle**, 212.
Oil of amber with tolu, 76.
 of juniper, 50.
 of turpentine, 58. 137. 150.
 of wormseed, 99.
 camphorated, 115.
 liniment, 116.
Ointment of tartar emetic, 105.
 of arsenic, 113.

Ointment of red precipitate, 126.
 of white precipitate, 126.
 of iodine and calomel, 108.
 of iodide of zinc, 109.
 of hydriodate of potash, 109.
 of iodine and potassium, 109, 110.
 of ioduretted iodide of potassium, 110.
 of iodide of arsenic, 111.
 of cyanide and mercury, 110, 111.
 of iodide of sulphur, 111.
 of bromium, 111.
 of creosote, 112.
 of belladonna, 112.
 of sulphur, 112.
 Kentish, 111.
 of gall nuts, 220.
 of oxide of zinc, 220.
 of rhubarb and opium, 220.
 of tar and opium, 221.
 comp. and sulph. 221.
 of protoiodide of mercury, 109.
 of iodide of zinc, 109.
 of belladonna, 152.
 of alum and calomel, 220.

Oleaginous mixture, 34.

Olive oil with ammonia, 99.

Opium with nitre, &c., 60.
 with guaiacum, nitre, &c., 60.
 with ipecacuanha, &c., 61.
 with calomel and ipecacuanha, 61.
 with camphor, &c., 204.
 liniment, 114.
 with soap in pills, 141.
 with hyoscyamus, 142.
 with digitalis, 46. 144.
 enema, 153.
 with lime water, 154.
 suppository, 154.
 plaster, 154.
 with acetate of lead, 201, 202.
 antidotes of, 244.
 with quinia and calomel, 196.

Opiate confection, 144. 160.
 mixture, 146.
 collyrium, 147.
 enema, 153.

Oxide of bismuth, 175.
 of zinc, 176. 218.
 of arsenic, 124.
 of zinc, ointment of, 220.

Oxalic acid, antidotes of, 245.

P.

Paregoric with cicuta, &c. 75.
 with squills, 74.
 elixir, 151.

Parsley infusion, 49.

Panado, 231.

Pea issue, 129.

Peruvian bark. See Cinchona.
 balsam for the ear, 210.

Pellitory, 166.

Pectoral mixture, 82.

Phosphoric acid, 126.

Phosphorous emulsion, 137.

Phosphate of soda, 28.

Pipsisseway, 55.

Piperina with gentian, 175.

Pink root with calomel, 95.
 with savin, 96.
 infusion, 98.

Plummer's pills, 64.

Pleurisy root, 68.

Plaster, warm, 119.
 of mercury, 104.
 of opium, 154.
 of mercurial ointment, &c. 104.
 of tartar emetic, 105.

Poppies, collyrium of, 207.

Pomegranate bark, 100.
 gargle, 214.

Potash, sub-carb. of, with lime water, 58.
 caustic, 114.

Potassium, cyanide of, 83. 113.
 iodide of, 190.

Pommade ammoniacale, 107.

Poultices. See Cataplasm.

Poisons, 238.

Protoiodide of mercury with morphia, 129.

Protoiodide of mercury, 109. 163.

Prunes, decoction of, 38.

Prussic acid solution, 81, 82.
 mixture, 82.

Prussiate of iron and guaiacum, 173.

Pyroligneous acid, 209.
 mixture of, 182.

Q.

Quassia, infusion of, &c. 184.
 pills of extract of, 176.
 with ammoniuret of copper, 165.
 with serpentaria, 172.

Quinia pills, 173.

with carbonate of ammonia, 174.

in sirup, 179.

in mixture, 180.

with calomel, &c. 196.

with blue mass, &c. 197.

enema, 187.

gargle, 213.

R.

Red precipitate ointment, 168.

Rhatany root, 202. 217.

Rhubarb with calomel, 25, 26.

with soap, 29, 30.

with aloes, 31.

with sulphate of iron, 30.

with magnesia, 26.

with ipecacuanha, 128.

with columbo, 170.

with opium, (ointment,) 220.

with chalk, 222.

with carbonate of soda, 223.

Rennet whey, 236.

Rice water, 229.

gruel, 231.

Rob antisypilitique of Laffeteur, 66.

Rubefacients, 104.

S.

Sage tea, 230.

Saline mixture, 62.

Sarsaparilla decoction, 68.

with senna, &c. 66.

beer, 236.

Sago, 232.

Sal æratus with Seltzer water, 57.

Salep powders, 232.

Salicine, 173.

Savin with ginger, &c. 89.

with cantharides, 89.

with hellbore, 92.

with pink root, 93.

with acetate of copper, 128.

Sassafras infusion, 126.

Scuddamore's gout mixture, 40.

Seltzer water, 55.

Serpentaria with sulphuret of mercury, 63.

infusion of, 185.

Secale cornutum, 89. 91.

Setons, 128, 129.

Senna with figs, 34.

with manna and salts, 37.

Senna, infusion of, 38.

infusion of, with rhubarb, 38.

with pink root, 95.

Serpentaria, 185.

Seneka with squills, 23.

infusion, 74.

in decoction, 84. 90.

Seidlitz powders, 28.

Sialagogues, 161.

Sirup of galls and brandy, 189.

Snake root, Virginia. See *Serpentaria*.

Soap pills, 57.

liniment, 113.

Soda, carbonate of, with gentian, 199.

chloride of, (lotion,) 191.

solutio vitriolica, 21.

phosphate of, 28.

sub-carb. of, with copaiva, 57.

Sponge, burnt, 193.

Spurge, 36.

Spermaceti mixture, 80.

Spirits of turpentine, 58.

with castor oil, 36.

enema of, 43.

mixture of, 50.

124.

with hartshorn, 100.

with ether, 100.

with cantharides, 105.

with laudanum, &c. 137. 150.

linctus of, 137.

Spice plaster, 117.

Spurred rye, 89. 91.

Starch, mucilage of, 235.

Strychnia pills, 145.

tincture of, 145.

draught of, 146.

acetate of, 193.

collyrium, 208.

Stramonium, extract of, 143. 153.

Stimulants, 132.

Squills with potash, 45.

with morphia, 83.

with calomel, 47.

with calomel and digitalis, 47.

with paregoric, &c. 74.

with colchicum and tobacco, 49.

with carbonate of potash, &c. 51.

with cicuta, &c. 72.

with calomel and ammoniac, 72.

with calomel, opium, &c. 73.

with seneka, 22.

Steel filings, 176.

- Sub-carbonate of potash. See Carbo-
nate of Potash.
- Sugar of lead. See Acetate of Lead.
- Sulphate of copper, 19. 175. 201. 217.
of copper, pills of, 175.
of copper with bole, 211.
of copper, collyrium of, 207.
of copper, injection of, 211.
of zinc, 19. 21.
of zinc with alum, 20.
of zinc with myrrh, 71.
of zinc in pills, 72.
of zinc, (injection,) 210.
of zinc, (gargle,) 214.
of zinc with bole, (injection,) 211.
of potash with rhubarb, 27.
of potash with opium, &c. 60.
of magnesia. See Epsom
Salts.
of alumine with kino, 200.
of alumine with opium, 201.
of morphia, 142.
of iron with senna, jalap, &c. 90.
of iron with aloes, &c. 90.
of iron with extract of bark, &c. 172.
of iron with sulphuric acid, 184.
of quinia. See Quinia.
- Sulphur and magnesia, 27.
confection of, 32.
electuary, 42.
ointment, (aromatic,) 221.
- Sulphuric acid mixture, 178.
- Suet ptisan, 235.
- Supertartrate of potash. See Cream of Tartar.
- Sulphuret of antimony, 61.
of mercury, 63.
of potash, lotion of, 219.
- Sulphurous water, 120.
- Sulphureo-gelatinous bath, 120.
- Sublimate bath, 121.
- Suppository of opium, 144.
- Sweet spirits of nitre, &c. 64.
- Swaim's panacea, 66.
- T.
- Tapioca jelly, 232.
- Tartar emetic solution, 20.
and quinine, 20.
enema, 43.
with nitre and calomel, 59.
- Tartar emetic with camphor and opium, 63.
plaster, 105.
pills, 72.
ointment, 105.
- Tartar whey, 236.
salt of. See Carbonate of Potash.
- Tar liniment, 115.
ointment, 220, 221.
pills, 72.
water, 79.
fumigation, 85.
- Tannin pills, 203.
- Tea injection, 212.
- Tin electuary, 97.
antidotes of, 246.
- Tincture of tolu with antimonial wine, 64. 78.
of iodine, 93.
of blood-root, 77.
of guaiacum, 92.
of digitalis, 77. 141.
of bark with citrate of potash, 166.
of bark, 170.
of muriate of iron, 170.
of catechu, (mixture,) 189.
- Tolu with digitalis, 79.
with ammoniac, 84.
with asafœtida, 77.
with amber oil, 76.
inhalation of, 84.
with morphia, 84.
inhalation, 85.
with belladonne, 80.
- Tobacco fomentation, 122.
poultice, 23.
enema, 44.
with squills and colchicum, 49.
- Tonics, 170.
- Turpentine pills, 48. 98. 132.
and cantharides, 111. 1.
mixture, 34, 35. 137.
oil of. See Spirits of Turpentine.
- U.
- Uva ursi with soda, 46.
with cinchona, &c. 55.
infusion, 52.
- V.
- Valerian with cinchona, 184.
with carb. of iron, 157.

- Valerian bolus, 157.
 with musk and camphor, 158.
 with Hoffman's anodyne, 158, 159.
 Veratrina, tincture pills, and ointment of, 40, 41.
 Vegetable soup, 233.
 Vermifuges, 95.
 Vermifuge bolus, 97.
 Venice turpentine pills, 96.
 Vitriol, blue. See Sulphate of Copper.
 Verdigris, powder of, 127.
 Vinegar collyrium, 206.
 whey, 236.
 draught, 184.
 Volatile tinct. of guaiac. 92.
- W.
- Warm plaster, 119.
 Water gruel, 231.
 Warner's cordial, 39.
 White's precipitate ointment, 115.
 Whey, alum, 216.
 wine, 139.
 rennet, 236.
- Whey, tartar, 236.
 vinegar, 236.
 Wine of ipecacuanha with antimony, 19.
 of colchicum, 39. 52.
 whey, 139.
 mixture, 140.
 of opium, 151.
 Wistar's cough lozenges, 73.
 Wild cherry bark, 185.
 Wormseed oil, 99.
 Worm tea, 98.
- Y.
- Yest poultice, 118.
 Yellow wash, 126.
- Z.
- Zinc, pills of oxide of, 176. 218.
 sulphate of, 210.
 antidotes of, 246.
 acetate of, (collyrium,) 206.
 iodide of, 109.
 ointment of, 220.

Lea & Blanchard have lately published.

N E W R E M E D I E S,
The Method of Preparing and Administering them;
THEIR EFFECTS

UPON THE
HEALTHY AND DISEASED ECONOMY, &c. &c.

BY ROBLEY DUNGLISON, M. D.

*Professor of the Institutes of Medicine and Materia Medica in
Jefferson Medical College of Philadelphia; Attending Physi-
cian to the Philadelphia Hospital, &c.*

THIRD EDITION BROUGHT UP TO 1841.
IN ONE VOLUME.

A N E W E D I T I O N

Completely Revised, with Numerous Additions and Improvements,

OF

DUNGLISON'S DICTIONARY

OF

MEDICAL SCIENCE AND LITERATURE:

CONTAINING

A concise account of the various Subjects and Terms, with a vocabulary of
Synonymes in different languages, and formulæ for various officinal and empiri-
cal preparations, &c.

IN ONE ROYAL 8vo. VOLUME.

A Fourth Edition Improved and Modified, of

D U N G L I S O N ' S
H U M A N P H Y S I O L O G Y :

ILLUSTRATED WITH NUMEROUS ENGRAVINGS.

IN TWO VOLUMES, OCTAVO.

Brought up to the present day.

A PRACTICAL TREATISE

ON THE

HUMAN TEETH:

Showing the causes of their destruction and the means of their preservation. By Wm Robertson: with plates. First American, from the second London edition. In one volume

OUTLINES

OF A

COURSE OF LECTURES, ON MEDICAL JURISPRUDENCE.

BY THOMAS STEWART TRAILL, M. D.

From the Second Edinburgh Edition,

WITH AMERICAN NOTES AND ADDITIONS.

ARNOTT'S ELEMENTS OF PHYSICS.

Complete in One Volume.

A new edition of Elements of Physics, or Natural Philosophy, general and medical, written for universal use, in plain or non-technical language, and containing New Disquisitions and Practical Suggestions, comprised in five parts: 1st. Somatology, Statics and Dynamics. 2d. Mechanics. 3d. Pneumatics, Hydraulics, and Acoustics. 4th. Heat and Light. 5th. Animal and Medical Physics. Complete in one volume. By Neil Arnott, M. D., of the Royal College of Physicians. A new edition, revised and corrected from the last English edition, with additions, by Isaac Hays, M. D.

THE NINTH BRIDGEWATER TREATISE.

A F R A G M E N T,

BY

C H A R L E S B A B B A G E, E S Q.

From the Second London Edition.

IN ONE VOLUME, 8vo.

A New Edition with Supplementary Notes, and Additional Plates; of BUCKLAND'S GEOLOGY AND MINERALOGY, considered with reference to Natural Theology; from the last London Edition with nearly one hundred Maps and Plates.

PROFESSOR GIBSON'S RAMBLES IN EUROPE, in 1839:—Containing Sketches of Prominent Surgeons, Physicians, Medical Schools, Hospitals, &c. &c. In One Volume.

AN ATLAS OF PLATES, illustrative of the Principles and Practice of Obstetric Medicine and Surgery, with descriptive Letter Press, by Francis H. Ramsbotham. This will form a large super royal volume, with over One Hundred lithographic plates—to be ready in November.

THE PRINCIPLES AND PRACTICE of MEDICINE. By Professor Dunglison in 2 vols. 8vo.

This work will be ready the approaching fall.

THE LIBRARY OF PRACTICAL MEDICINE. Edited by Tweedie, is now complete in five volumes, royal octavo, handsomely bound in leather, to match. The different volumes may be had separate, bound in extra cloth.

CATALOGUE

OF

BOOKS,

PUBLISHED

BY

LEA & BLANCHARD,

PHILADELPHIA,

AND

FOR SALE

BY

1840

BOOKS,

PUBLISHED

BY

LEA & BLANCHARD,

PHILADELPHIA.

A

NAVAL HISTORY

OF

THE UNITED STATES,

BY J. FENIMORE COOPER, ESQ.

IN TWO HANDSOME VOLUMES, BOUND IN EMBOSSED CLOTH.

A new edition revised and corrected with an index to the volumes.

We have pursued this history with no little curiosity and with great interest. Considering the brief existence of the American marine, its annals are more eventful, more romantic, and more various, than any in existence. Nothing can surpass the energy which enabled the United States to form an effective navy, at a time when they could hardly be said to have had a political existence, and when they were beset by greater difficulties than any which an infant nation had ever yet to encounter. This consideration has animated the present historian, whose enthusiasm seems to be kindled by his office of chronicler, even more than when he formerly sought inspirations from the same source in constructing his famous stories of the sea. His national pride has, however, not tempted him to be, after the manner of his countrymen, vain-glorious as regards his own nation, and abusive towards others. His work, accordingly, is more fair and candid than could have been expected on such a theme from an American pen. Altogether this history is a valuable one, and cannot fail to pass into universal circulation. The incidents which took place in the naval war with Tripoli, are grander and more heroic than any thing in the circle of romance, and are detailed with all the vigor and animation of Mr. Cooper's genius.—*British Naval and Military Magazine*.

The History of the Navy of the United States from the earliest period of its existence, in the dawn of the Revolution, through

all its discouragements, reverses, trials and glory, was a task worthy of the author who had established a reputation as a describer of nautical events, superior to that of any other living writer. The task has been so performed as to leave nothing to desire. No work of higher interest has been published in the United States for many years. The theme is one which Mr. Cooper seems to treat *con amore* and for which his early life and education fitted him, above all other men. If we are not mistaken, the publication of this book is calculated to heighten the already exalted estimate in which the Navy is held, and to render it still more, if possible, a favourite with the nation. Whilst Mr. Cooper has, at all times, given full credit to the officers and crews of the vessels whose victories, during the late war, shed so much renown upon our arms, he has not been guilty of the bad taste, which a writer of less discrimination would scarcely have avoided, of indulging a vainglorious spirit and a disparaging tone in reference to our great rival upon the ocean. The glories of American victories are fully portrayed, whilst, at the same time, care is taken, in every case to exhibit a fair and impartial estimate of the strength and appointments of antagonists' vessels or fleets. In this way the work acquires the credit due to a grave and impartial history. The book is splendidly published by Lea and Blanchard.—*Baltimore Chronicle*.

SCOTT'S POETICAL WORKS.

A fine edition in Six Volumes, Royal 12mo., printed on a beautiful paper, and handsomely bound in embossed cloth, or in full coloured calf, of

THE

POETICAL WORKS

OF

SIR WALTER SCOTT,

COMPLETE.

LIFE OF SIR WALTER SCOTT,

To match the above, in Seven Royal 12mo. Volumes.

MEMOIRS OF THE LIFE OF

SIR WALTER SCOTT.

BY J. G. LOCKHART, Esq.

A cheaper edition of this work may also be had in two volumes octavo, done up in embossed cloth, or bound up in leather.

We have here a recurrence of that pleasing excitement, through all the reading world, which was wont to attend the appearance of each new "Waverly novel." The voice of the mighty minstrel again rises, as it were, from the grave, and we cease for a time in the midst of the anxious and feverish contentions of faction, to listen to the tones which delighted and astonished our early years.—*Dundee Chronicle.*

To match the above, a new and complete edition of

THE PROSE WORKS

OF

SIR WALTER SCOTT.

(In preparation.)

A new and complete edition of
THE POETICAL WORKS

OF

THOMAS MOORE,

TO MATCH

SCOTT'S POETICAL WORKS.

(In preparation.)

MRS. HEMANS POETRY,

A COMPLETE AND UNIFORM EDITION
OF

MRS. HEMANS WORKS;

TO WHICH IS PREFIXED

A MEMOIR BY HER SISTER,
MRS. HUGHES.

Together with an Essay on her Genius, by

MRS. SIGOURNEY,

In Seven Volumes Royal 12mo., to Match the Works of Scott, &c. &c. With a fine portrait.

CRABBE'S LIFE AND POEMS.

The Posthumous Poems of the Rev. George Crabbe, with his Letters and Journals, and a Memoir of his Life. By his Son and Executor. In two handsome vols.

MATHEWS'S MEMOIRS.

MEMOIRS OF CHARLES MATHEWS, COMEDIAN, by Mrs. Mathews.

A man so various, that he seemed to be, Not one, but all mankind's epitome.—*Dryden.* "Proteus for shape, and mocking bird for tongue."

In two volumes, 12mo.

ALSO

A CONTINUATION OF THE MEMOIRS OF CHARLES MATHEWS, COMEDIAN, by Mrs. Mathews, including his correspondence and an account of his residence in the United States. In two vols. 12mo.

"Every page of this work teems with entertainment."—*Morning Chronicle.*

"These volumes are 'heaped up and running over' with anecdotes of the most singular and diverting nature. They furnish us with racy personal sketches of numerous eminent literary, political, and theatrical characters."—*Sun.*

"Mr. Mathews was an honour to his country and his profession. His memoirs, which have now been furnished to the world by Mrs. Mathews, are replete with all that can interest the admirers of this singular man. His letters are very amusing and display all the different phases of his character with singular fidelity."—*Corsair.*

MOORE'S LALLA ROOKH:

BEAUTIFULLY ILLUSTRATED WITH THIRTEEN ENGRAVINGS,

Finished in the highest style of art, under the superintendence of

MR. CHARLES HEATH.

In one super royal 8vo. volume, handsomely bound in extra embossed cloth or in superb white calf and morocco, gilt edges.

LIST OF ILLUSTRATIONS.

LALLA ROOKH, <i>by Meadows.</i>	THE DEPARTURE OF HAFFET, <i>by Corbould.</i>
THE PERI AT THE GATE OF EDEN, <i>by Meadows.</i>	DEATH OF HINDA, <i>by do.</i>
THE PERI'S FIRST PILGRIMAGE, <i>by Corbould.</i>	ZELICA, <i>by do.</i>
THE PERI'S SECOND PILGRIMAGE, <i>by Corbould.</i>	AZIM AND ZELICA, <i>by do.</i>
THE PARTING OF HINDA AND HAFFET, <i>by Stephanoff.</i>	ZELICA DISCOVERING THE VEILED PROPHET, <i>by do.</i>
	HINDA, <i>by Stephanoff.</i>
	NOURMAHAL ASLEEP, <i>by do.</i>
	NAMOUNA, <i>by Meadows.</i>

This is a beautiful and splendid volume. The nineteenth edition is so fittingly adorned, that it will be very apt to cause its eighteen predecessors to be neglected, if not forgotten. —*Literary Gazette.*

This exquisite poem, long the admiration of readers of all classes, has received in the costly volume before us a form worthy of its

internal beauty and universal popularity. The noble dimensions of the book, the clearness of the type, and the general splendour of the style in which it is prepared, in all its departments of art, will recommend it afresh to the world of readers and lovers of poetry. —*Atlas.*

JAMES' BOOK OF THE PASSIONS, TO MATCH LALLA ROOKH.

In one handsome royal octavo volume, printed on fine paper, with 16 beautiful engravings done on steel, from designs by the best artists; and bound in extra embossed cloth or in superb white calf and morocco extra with gilt edges.

WALPOLES LETTERS.

NOW PREPARING A COMPLETE EDITION
OF
THE LETTERS OF HORACE WALPOLE,
(EARL OF ORFORD.)

INCLUDING UPWARDS OF THREE HUNDRED NEW LETTERS.

The whole now first chronologically arranged, the Names inserted at full length, and explanatory Notes added.

FORMING AN ANECDOTAL HISTORY
OF THE CHIEF PART OF THE LAST CENTURY—AND TO WHICH
WILL BE ADDED, HIS REMINISCENCES.

To be handsomely printed in Octavo.

JACOB'S ON METALS.

AN HISTORICAL INQUIRY INTO THE PRODUCTION AND CONSUMPTION OF THE PRECIOUS METALS, from the Earliest Ages, and into the Influence of their Increase or Diminution on the Prices of Commodities. By William Jacob, Esq. F. R. S. In 8vo.

MUDIE'S HEAVENS.

THE HEAVENS. With cuts. By Robert Mudie, 1 vol. 12mo.

MUDIE'S EARTH.

THE EARTH. By Robert Mudie, 1 vol. 12mo.

GRANT'S LIFE IN LONDON.

EVERY DAY LIFE IN LONDON, By James Grant, Author of "Random Recollections of the Lords and Commons," "The Great Metropolis," &c. &c. In two vols. 12mo.

DUMAS'S FRENCH REVOLUTION.

Memoirs of his Own Time, including the Revolution, the Empire, the Restoration, and his Residence in the United States. By Lieut. General Count Mathieu Dumas, Private Secretary to Napoleon. In two royal 12mo vols.

"As a history his book is replete with the most instructive information upon the most extraordinary events that have ever agitated the world—events in which he performed no obscure or insignificant part; and as a personal memoir, it is one of the most deeply interesting we have ever read.

"Few men have in a long life passed through so many strange scenes, and been actors in so many great and surprising events as Count Dumas, and the narrative of these is given with a modesty and impartiality that enhances their value, and with a clearness and distinctness that throws much light upon all we have before known of the history of those eventful times."—*Evening Gazette and Galaxy*.

RUSCHENBERGER'S VOYAGE.

A VOYAGE ROUND THE WORLD.

INCLUDING AN EMBASSY TO MUSCAT AND SIAM IN 1835, 1836, and 1837. By W. S. W. Ruschenberger, M. D., *Surgeon U. S. Navy; Honorary Member of the Philadelphia Medical Society; Member of the Academy of Natural Sciences of Philadelphia, &c.*; Author of "Three Years in the Pacific." In one vol. 8vo.

LIFE OF JEFFERSON.

THE LIFE OF THOMAS JEFFERSON, third President of the United States, with parts of his correspondence never before published, and notices of his opinions on questions of Civil Government, National Policy, and Constitutional Law. By George Tucker, Professor of Moral Philosophy, in the University of Virginia. With an engraved bust. In two volumes.

LADY MONTAGUE'S LIFE AND WORKS.

THE CORRESPONDENCE OF LADY MARY WORTLEY MONTAGUE, edited by Lord Wharnccliffe, including upwards of 150 letters, never before published. A memoir of the Court of George I. by Lady Montague, and a Life of the Authoress, illustrative anecdotes, and notes added, and the suppressed passages restored. Handsomely bound in two volumes, embossed cloth.

LA MARTINE'S PILGRIMAGE.

A PILGRIMAGE TO THE HOLY LAND, comprising Recollections, sketches, and Reflections made during a tour in the East, by Alphonse de la Martine. Fourth Edition.

KEBLE'S CHRISTIAN YEAR.

THOUGHTS IN VERSE, FOR SUNDAYS, AND HOLY DAYS THROUGHOUT THE YEAR. By the Rev. John Keble, Professor of poetry, in the University of Oxford.

"In quietness and confidence shall be your strength."

—ISAIAH XLV. 15.

A new Edition, with a farther revision; and an Introduction by the Right Rev. George W. Doane, Bishop of New Jersey, now preparing in one neat volume.

"These verses were singularly beautiful in conception and composition, and breathe the purest poetic taste, and the most sincere and fervent spirit of piety."—*Gazette*.

"The Christian Year, a collection of little poems." "By the Rev. John Keble, a name well known, and justly dear to Oxford."—"We scarcely ever remember to have read so exquisite an invocation of religion in poetry."—"We have so high a sense of its genuine worth, that we hope this notice of it may be the means of introducing it into many families."—*British Critic*.

SMOLLETT'S WORKS.

The Select works of TOBIAS SMOLLETT, with a Memoir of the Life and Writings of the Author. By Sir Walter Scott. In two handsome Royal Volumes. Various bindings.

FIELDING'S WORKS.

The Select Works of HENRY FIELDING, with a Memoir of the Life of the Author. By Sir Walter Scott. And an Essay on his Life and Genius, By Arthur Murphy, Esq. In two handsome Library Volumes. Various bindings.

THE PEOPLE'S LIBRARY.

"The editors and publishers should receive the thanks of the present generation, and the gratitude of posterity, for being the first to prepare in this language what deserves to be called, not the *Encyclopædia Americana*, but *THE PEOPLE'S LIBRARY*."—*New York Courier and Inquirer*.

THE ENCYCLOPÆDIA AMERICANA.

A Popular Dictionary of Arts, Sciences, Literature, History, and Politics, brought down to the Present Time, and including a copious Collection of Original Articles in American Biography; on the basis of the Seventh Edition of the German *Conversations-Lexicon*. Edited by Francis Lieber, assisted by Edward Wigglesworth and T. G. Bradford, Esqs. In Thirteen Large Volumes, Octavo.

HISTORY OF CONGRESS.

Exhibiting a Classification of the Proceedings of the Senate and House of Representatives, from March, 1789, to March, 1793, embracing the First Term of the Administration of General Washington. In one volume, octavo.

THE ENCYCLOPÆDIA OF GEOGRAPHY.

By Hugh Murray. Revised, with Additions, by Thomas G. Bradford. In Three Beautiful Volumes. With over Eleven Hundred Cuts. (Published by subscription.)

FRENCH REVOLUTION.

SKETCHES OF THE MEN AND TIMES OF THE FRENCH REVOLUTION, By Madame Tussaud. In two volumes, 12mo.

BROUGHAM'S STATESMEN.

HISTORICAL SKETCHES OF STATESMEN WHO FLOURISHED IN THE TIMES OF GEORGE III. BY LORD BROUGHAM. First and second series. In 4 vols. 12mo.

"The readers of the *Edinburgh Review* are already familiar with these powerful sketches of the great English Statesmen and Orators of the past generation; written by one who is himself, perhaps, the greatest statesman and orator of this. They who have read them, will need no urging to read them again, and those who have not, will require no other inducement than the author's name. These sketches deserve to be studied alike for the strength and purity of the style, and the vigorous thought and deep interest of the matter."—*Ev. Gazette*.

BROUGHAM'S OPINIONS.

Opinions of Lord Brougham on Politics, Theology, Law, Science, Education, Literature, &c. &c. as exhibited in his Parliamentary and Legal Speeches, and Miscellaneous Writings, together with a Memoir of the author, in 2 volumes, 12mo.

The object of this publication was not only to embody the most brilliant passages from the most celebrated speeches and writings of Lord Brougham, but also to develop, in a gradual manner, the particular mind and genius of the man. This design is well carried out, and we arise from the earnest perusal of the book with our opinion strengthened, if

possible, in regard to the extraordinary character and exceeding vigour of the intellect discussed. Perhaps, however, the best portion of the work is embodied in the Prefatory Memoir, which contains more complete, accurate, and satisfactory information about his Lordship, both in his public and private life, than any thing of the kind hitherto published.—*Gentleman's Magazine*.

BROUGHAM'S SPEECHES.

A Selection of the Speeches of Lord Brougham. In 1 vol. 8vo. In preparation.

TIMES OF GEORGE IV.

A CONTINUATION OF THE DIARY ILLUSTRATIVE OF THE TIMES OF GEORGE IV., interspersed with Original Letters from Queen Caroline, the Princess Charlotte, and other distinguished persons. Edited by John Galt, Esq. In 2 vols. 12mo.

"A work replete with deep interest to all who would know those whose deeds make the sum of history. The times of George III., and the times of those who having lived years with him still survive to give consequence to succeeding reigns, and to influence indeed the affairs of the world, are full of interest and instruction."—*U. S. Gazette*.

ELLIOTT'S TRAVELS.

"Travels in the Three Great Empires of Austria, Russia, and Turkey, by C. B. Elliott, F. R. S., Vicar of Godalmin, late of the Bengal Civil Service, and author of 'Letters from the North of Europe.' In two vols. royal 12mo.

These highly entertaining and instructive volumes will be sought after with avidity, by those who are desirous of becoming acquainted with the manners and habits of the people of Austria, Russia, and Turkey. The work is one of intense interest, and is written in a style at once free, easy and familiar. The social, domestic, and political relations of each nation are dwelt upon, while the accounts of cities, scenery, and antiquities are equally interesting. Religion and Literature are not forgotten, but introduced throughout the volumes in a highly philosophical spirit.—*Saturday Chronicle*.

CITY OF THE CZAR; OR

A Visit to St. Petersburg, in the Winter of 1829—30. By Thomas Raikes, Esq. In two vols. 12mo.

THE STRANGER IN CHINA;

Or, the Fan-qui's Visit to the Celestial Empire. By C. T. Downing. Two volumes, 12mo.

THE POETIC WREATH.

Consisting of Select Passages from the Works of English Poets, from Chaucer to Wordsworth, Alphabetically arranged, with Twenty-six Beautiful Wood Cuts. In one neat Royal 18mo volume, handsomely bound in embossed cloth, with gilt edges.

The selections are made with much taste, presenting characteristic and striking passages from the best of the English poets, in various ages, forming, as it were, a history of style in this species of composition; and the mechanical execution of the Wreath is such as to render it one of the prettiest volumes of the season, the embellishments, binding, and typography, being all elegant and appropriate.—*Pennsylvanian*.

The object of the editor has been to present a collection of beautiful or striking extracts, and to furnish at the same time specimens of the authors from whose works they are selected. A more delightful little volume, or one better calculated to grace the centre table or the private library, has not been examined by us for many a day.—*Pennsylvanian Inquirer*.

THE LANGUAGE OF FLOWERS.

With Illustrative Poetry. To which is now first added, The Calendar of Flowers. Revised by the Editor of "Forget Me Not." A New Edition, with New Plates.

Handsomely done up in embossed leather and gilt edges.

THE SENTIMENT OF FLOWERS;

Or, Language of Flora. Embracing an account of nearly 300 Different Flowers, with their powers in language. With Coloured Plates. A small volume in embossed cloth.

POCKET LAÇON.

In two handsome volumes.

The Pocket Lacon, comprising near one thousand extracts from the best authors, selected by John Taylor.

A work that, dip where the reader may, he will find a fund of knowledge: and which he may continue to peruse, lay down and take up at pleasure, without breaking the thread, or interrupting the chain of reasoning.

These volumes illustrate the advantage of a practice which we have never had the patience to adopt, viz: that of keeping a common-place book. The compiler has read largely, and to good purpose; and has read much that we like to read—that is, among the old authors. Adopting the practice of noting the best thoughts of the best writers of all ages; in their own words, he has collected a casket of literary gems which he spreads out before us with lavish profusion, and at small price.—*American*.

MRS. HEMANS MEMOIRS.

Memoir of the Life and Writings of Mrs. Hemans. By her sister, Mrs. Hughes.

The publishers deserve great praise for the very handsome manner in which they have got up this volume. Its appearance is as creditable as its contents are agreeable.

The life of Mrs. Hemans by one who had access to her at all times—one who was familiar with her from her infancy upwards—one who knew of all her sorrows and her triumphs, and who fully understood the peculiar organization of her mind, is, of course, a book full of interest. We have found it eminently so, and we cordially recommend it to all who cherish the memory of this gifted woman.—*Lady's Book*.

THE YOUNG WIFE'S BOOK.

A MANUAL OF RELIGIOUS, MORAL AND DOMESTIC DUTIES.

A small volume bound in extra, with plates engraved on steel.

THE YOUNG HUSBAND'S BOOK.

A Manual of the Duties, Moral, Religious, and Domestic, imposed by the Relations of Married Life.

A small volume, bound in extra, with plates engraved on steel.

LAWS OF ETIQUETTE;

Or, Rules and Reflections for Conduct in Society. By a Gentleman.

A Small Pocket Volume. A work of which many thousands have been sold.

ETIQUETTE FOR THE LADIES.

With Hints on the Preservation, Improvement, and Display of Female Beauty.

A small volume to match the Laws of Etiquette.

THE CANNONS OF GOOD BREEDING;

Or the Hand-Book of the Man of Fashion, by the author of "The Laws of Etiquette;" a small pocket volume in embossed cloth.

It is original; written with remarkable ease and elegance, most agreeably seasoned with wit and anecdote; and while it lays down with precision the laws of behaviour in good society, it affords a rich treat of pleasant reading.

"The truth is, that the volume abounds in good things. We may safely say that, in a compass so small, we never before met with an equal radiancy of fine wit, so well commingled with scholar-like observation and profound thought—thought sometimes luminously logically, and always elegantly expressed."—*Gentleman's Magazine*.

ADVICE TO A YOUNG GENTLEMAN.

Upon Entering Society. By the Author of the "Laws of Etiquette" A small volume, in embossed cloth.

This is the work of an original, independent and vigorous writer. It is in a much more serious and earnest style than the "Laws of Etiquette." The same acuteness of perception, knowledge of human character and fine classical taste, characterise this work, of which were so conspicuous in the former. As a monitor to the young gentleman, on his entrance into society, it will be variously estimated according to the preconceived notions of its readers; but one thing is certain, that no young man can read it without learning much that will be useful to him in his intercourse with mankind.—*Weekly Messenger*.

The book is replete with a worldly wisdom even profound; it is the product of a vigorous and cultivated mind imbued with a thorough knowledge of its subject, and discussing it *con amore*.

Among the detailed precepts which form its body, there is, perhaps, little to be found which the letters of Lord Chesterfield have not already given. But without the offensive heartlessness of these very objectionable writings, the American work equals them, at least, in all their reputable points—in vigour of thought and diction, in acumen, in practicability and in evidences of worldly knowledge.—*Gentleman's Magazine*.

MACKINTOSH'S REVOLUTION.

HISTORY OF THE REVOLUTION IN ENGLAND, IN 1688: By the late Right Hon. Sir JAMES MACKINTOSH. And completed to the settlement of the crown, by the editor. In 1 vol. 8vo.

SISMONDI'S ROME.

SISMONDI'S, History of the Fall of the Roman Empire. Comprising a view of the Invasion of the Barbarians. 1 volume, Octavo.

A NEW HISTORY OF ROME.

Principally from the German of Schlosser, Wachsmuth, Heeren, Niebuhr, &c. In 1 volume, Octavo.

This work is well calculated for the higher classes in Seminaries, as well as for the Library.

"The book is an indispensable one, from the complete *coup-d'œil* which it presents of Roman History illustrated by modern lights; it is a very readable one, from the closeness and rapidity of its march; and it is a very extraordinary one, from the immense quantity of matter comprised in so small a compass."—*Spectator*.

"We recommend this work very strongly to the English reader, as being the first compendium of Roman history yet published in England, in which the early ages of the Roman states have been exhibited in the new and striking light thrown upon them by the late researches and speculations of Niebuhr, Wachsmuth, and other recent German writers of eminence. Such a work has long been much wanted, and we trust it will be as extensively read as it deserves to be."—*Examiner*.

RAUMER'S ENGLAND.

ENGLAND IN 1835.

Being a Series of letters written to Friends in Germany, during a Residence in London and Excursions into the Provinces.

By FREDERICK VON RAUMER, Professor of the History at University of Berlin. In 1 Volume, Octavo.

MOORE'S IRELAND.

A HISTORY OF IRELAND: by Thomas Moore. Complete in 1 volume 8vo. (In preparation.)

LIFE OF R. H. LEE.

MEMOIRS OF THE LIFE OF RICHARD HENRY LEE. Together with his Correspondence. By his Grandson Richard H. Lee. In 2 vols. Octavo.

LIFE OF BAINBRIDGE.

LIFE AND SERVICES OF COMMODORE WILLIAM BAINBRIDGE, UNITED STATES NAVY.

By Thomas Harris, M. D., Surgeon United States Navy, and Member of the American Philosophical Society.

With a Portrait. 1 volume Octavo.

MEMOIRS OF SIR WILLIAM KNIGHTON, BART., G. C. H.

Keeper of the Privy Purse during the Reign of His Majesty King George the Fourth. Including his Correspondence with many distinguished Personages. By Lady Knighton. In 1 volume Octavo.

MEMOIRS OF CABOT.

A MEMOIR OF SEBASTIAN CABOT, with a Review of the History of Maritime Discovery. Illustrated by Documents from the Rolls, now first published.

"Put forth in the most unpretending manner, and without a name: this work is of paramount importance to the subjects of which it treats."—*Literary Gazette*.

IRVING'S FLORIDA.

THE CONQUEST OF FLORIDA BY HERNANDO DE SOTO.

By Theodore Irving, Esq. In 2 volumes, 12mo.

IRVINGS INDIAN SKETCH BOOK.

Indian Sketches taken during an Expedition to the Pawnee Tribes. By John T. Irving, Jun. 2 vols. 12mo.

CAREY ON CREDIT.

THE CREDIT SYSTEM OF FRANCE, GREAT BRITAIN, &c. By H. C. Carey, Author of Principles of Political Economy, &c.

CONTENTS.

Chapter I. Rise and Progress of Credit —2. Of Credit in France—3. Of Credit in England—4. Of Credit in the United States —5. Insecurity of Property and its Effects: England, France, United States—6. Fric-

tion: England, France, Scotland, United States—7. The United States in 1836-37—8. England in 1835 and 1836—9. Conclusion.

CAREY ON CURRENCY.

ANSWERS TO THE QUESTIONS:—What constitutes Currency? What are the causes of Unsteadiness of the Currency? And what is the Remedy? By H. C. Carey.

CAREY ON POLITICAL ECONOMY.

PRINCIPLES OF POLITICAL ECONOMY. Of the Laws of the Production and Distribution of Wealth.

By H. C. Carey, Author of an Essay on the Rate of Wages. In 3 volumes Octavo.

CAREY ON WAGES.

ESSAY ON THE RATE OF WAGES, With an Examination of the Causes of the Differences in the Condition of the Labouring Population throughout the World.

By H. C. Carey. 1 Volume.

GALLATIN ON BANKING.

CONSIDERATIONS ON THE CURRENCY AND BANKING SYSTEM OF THE UNITED STATES. By Albert Gallatin.

RAGUET ON FREE TRADE.

THE PRINCIPLES OF FREE TRADE, illustrated in a series of short and familiar Essays. By Condé Raguet, Esq. 1 vol. 8vo.

LIFE OF JESUS, &c.

JESUS AND HIS BIOGRAPHERS; OR THE REMARKS ON THE FOUR GOSPELS, revised with copious additions, by W. B. Furness. In 1 volume 8vo.

To all who are acquainted with the clearness, beauty and accuracy of style; the diligent research, the logical strictness of argument, and the profound truthfulness of Mr. Furness as a writer, and especially as an expositor of the Scriptures, nothing need be said to induce them to read with interest and attention this work on the Four Gospels. But we would say to all who are seeking for moral truth—for the light of

pure Christianity—thirsting for a real personal knowledge of the Lord Jesus Christ—and for a higher and truer estimation of those matchless records, his gospel, turn ye to this book—pursue this most remarkable exposition, the work of a pure minded, able, truth loving man, with the same pure spirit and sincere desire after the truth, that animates his pages, and ye will not find your labour in vain.—*Evening Gazette*.

GEISLER'S TEXT BOOK.

TEXT BOOK OF ECCLESIASTICAL HISTORY. By J. C. I. Geisler, Doctor of Philosophy and Theology, and Professor of Theology in Gottingen.

Translated from the Third German Edition, by Francis Cunningham. In 3 vols. 8vo.

Dr. Geisler's History is very highly recommended by gentlemen whose testimony to its value will have great weight, viz: Professors Stuart and Emerson, of Andover; Hodge, of Princeton; Sears, of Newton; and Ware, of Cambridge.

THE DISTRICT SCHOOL, OR NATIONAL EDUCATION. By J. Orville Taylor. The third edition.

"The style of the 'District School' is simple, intelligible, appropriate, correct and forcible; and the author displays much acquaintance with the condition of common education. We regard the general circulation of this work as highly desirable, for it will carry to every reader a large amount of important truths, many sound views of education, and an incitement to exertion in its promotion, which must produce some good effects."—*New-York Daily Advertiser*.

A CABINET ATLAS.

THE FAMILY CABINET ATLAS, constructed upon an original plan: Being a Companion to the Encyclopædia Americana, Cabinet Cyclopædia, Family Library, Cabinet Library, &c.

This Atlas comprises, in a volume of the Family Library size, nearly 100 Maps and Tables, which present equal to fifty Thousand Names of Places; a body of information three times as extensive as that supplied by the generality of Quarto Atlases.

MISS MARTINEAU.

How to Observe—Morals and Manners. By Miss Martineau. 1 volume, 12 mo.

WALSH'S DIDACTICS.

Didactic's Social, Literary, and Political. By Robert Walsh, Esq.

In 2 volumes, Royal 12mo. Embossed cloth.

COATE'S FAMILY ADVISER.

POPULAR MEDICINE;

OR, FAMILY ADVISER.

Consisting of Outlines of Anatomy, Physiology, and Hygiene, with such Hints on the Practice of Physic, Surgery, and the Diseases of Women and Children, as may prove useful in families when regular Physicians cannot be procured: Being a Companion and Guide for intelligent Principals of Manufactories, Plantations, and Boarding Schools, Heads of Families, Masters of Vessels, Missionaries, or Travellers; and a useful Sketch for Young Men about commencing the Study of Medicine.

BY REYNELL COATES, M. D.

Fellow to the College of Physicians of Philadelphia—Honorary Member of the Philadelphia Medical Society—Correspondent of the Lyceum of Natural History of New York—Member of the Academy of Natural Sciences of Philadelphia—Formerly Resident Surgeon of the Pennsylvania Hospital, &c.

ASSISTED BY SEVERAL MEDICAL FRIENDS.

In one handsome volume.

"It is with great satisfaction that we announce this truly valuable compilation, as the most complete and interesting treatise on Popular Medicine ever presented to the public. Simple and unambitious in its language, free from technicalities, and embracing the most important facts on Anatomy, Physiology and Hygiene, or the art of preserving health, and the treatment of those affections which require immediate attention, or are of an acute character, this should be in the hands of every one, more particularly of those who, by their situations are prevented from resorting to the advice of a physician; nor would the careful perusal of its pages fail to profit the inhabitants of our cities, by giving them a more accurate knowledge of the structure of the human frame, and of the laws that govern its various functions, whose perfect integrity is absolutely essential to health, and even to existence; the various systems of medical charlatanism, daily imagined to take advantage of the credulity and ignorance of mankind, would be rendered far less prejudicial to the community than they now are. We would particularly direct attention to the Chapter on Hygiene, a science in itself of the utmost importance, and ably treated in the small space allowed to it in this volume."—New York American.

A SERIES OF MEMOIRS of the STATESMEN of the ENGLISH COMMON-WEALTH, commencing with the Origin of the Popular Struggle, in the reign of James I., and closing at the Restoration. With original Portraits of Elliot, Pym, Hampden, Cromwell; and an Historical scene, after a Picture, by Cattermole. By John Forster Esq., of the Inner Temple.

(In preparation.)

TRAVELS IN EGYPT, by Prince Puckler Muskau,

(In preparation.)

CONVERSATIONS ON NATURE AND ART.

For the Use of Schools and Seminaries by a Lady.

With numerous wood cuts in one neat 12mo. volume, bound in embossed cloth, and lettered in gold.

This little work embodies a vast amount of useful, agreeable, and elegant information, and is full of truth and beauty. It is illustrated with engravings.—Saturday Chronicle.

"The rich stores, which are freighted in the discourse of Mrs. Fotescue with Henrietta and Frederick, are remarkable for the interest and importance, as well as the lucid and fascinating mode in which they are imparted. It is an invaluable assistant to the mother, who wishes to have useful and elevating converse with her little ones."

"This is a valuable little volume from the press of Lea & Blanchard, full of useful information, and instructive to all who wish to acquire a knowledge of Nature and Art. It is embellished with nearly

forty cuts, and the whole is well got up. It comes from a talented family, the authoress being the sister of Captain Marryat."

CHITTY'S JURISPRUDENCE.

CHITTY'S MEDICAL JURISPRUDENCE. A valuable work for Lawyers or Physicians. In royal 8vo.

ADVENTURES OF AN ATTORNEY IN SEARCH OF PRACTICE, by the Author of Adventures of a Gentleman in Search of a Horse, 2 vols. 12mo.

HERSCHEL'S PHILOSOPHY.

Preliminary Discourse on the Objects, Advantages, and Pleasures of the Study of Natural Philosophy. By J. T. W. Herschel, A. M. Late Fellow of St. John's College, Cambridge.

"Without disparaging any other of the many interesting and instructive volumes issued in the form of cabinet and family libraries, it is, perhaps, not too much to place at the head of the list, for extent and variety of condensed information, Mr. Herschel's discourse of Natural Philosophy in Dr. Lardner's Cyclopædia."—Christian Observer.

"The finest work of philosophical genius which this age has seen."—Mackintosh's England.

"Mr. Herschel's delightful volume. *** We find scattered through the work instances of vivid and happy illustration, where the fancy is usefully called into action, so as sometimes to remind us of the splendid pictures which crowd upon us in the style of Bacon."—Quarterly Review.

WASHINGTON IRVING'S WORKS.

THE WORKS

OF

WASHINGTON IRVING,

EMBRACING

THE SKETCH BOOK,
KNICKERBOCKER'S NEW
YORK,
BRACEBRIDGE HALL,

TALES OF A TRAVELLER,
THE CONQUEST OF GRANADA,
THE ALHAMBRA,

A New and Beautiful Edition, in Two Royal Octavo Volumes, with a Portrait of the
the Author.

THE ROCKY MOUNTAINS; or
Scenes, Incidents and Adventures in the
Far West.

Digested from the Journal of Capt. B. L.
BONNEVILLE, of the Army of the U. States,
and illustrated from various other sources.
By WASHINGTON IRVING.

With Two Large Maps.
In Two Volumes.

ASTORIA; or, Anecdotes of an Enterprise
Beyond the Rocky Mountains. By WASH-
INGTON IRVING.

Handsomely bound in Two Volumes,
embossed cloth or in leather.

A History of the Life and Voyages of
Christopher Columbus By WASHINGTON
IRVING.

Revised and corrected by the Author.
In Two Volumes, Octavo.

THE CRAYON MISCELLANY.
By WASHINGTON IRVING. In 3 vols.

Containing,

A TOUR ON THE PRAIRIES.
ABBOTSFORD AND NEWSTEAD
ABBEY.

LEGENDS OF THE CONQUEST
OF SPAIN.

NEW EDITIONS OF THE FOLLOW-
ING WORKS MAY BE HAD
SEPARATE.

THE SKETCH BOOK, 2 vols. 12mo.

KNICKERBOCKER'S HISTORY OF
NEW YORK, revised and corrected.
2 vols. 12mo.

BRACEBRIDGE HALL, OR THE HU-
MORISTS, 2 vols. 12mo.

TALES OF A TRAVELLER, 2 vols.
12mo.

A CHRONICLE OF THE CONQUEST
OF GRENADA. In 2 vols. 12mo.

THE ALHAMBRA; a series of Tales
and Sketches of the Moors and Spa-
niards. In 2 vols. 12mo.

THE BEAUTIES OF WASHINGTON
IRVING.

A Small Volume for the Pocket, neatly
done up in extra cloth.

COOPER'S WORKS.

A

NAVAL HISTORY.

OF

THE UNITED STATES.

A new Edition, Revised and Corrected with an Index to the Volumes.

In 2 volumes, 8vo.

GLEANINGS IN EUROPE, CONTAINING:

SKETCHES OF SWITZERLAND.

In 2 vols. 12mo.

A RESIDENCE IN FRANCE, with an Excursion up the Rhine, and a Second Visit to Switzerland. In 2 volumes 12mo.

FRANCE, with Sketches of Parisian Society. In 2 volumes, 12mo.

ENGLAND, with Sketches of Society in the Metropolis. In 2 volumes 12mo.

ITALY with Sketches of Society. In 2 volumes 12mo.

Now ready,

THE PATHFINDER,

OR

THE INLAND SEA. In 2 vols. 12mo.

THE MONIKINS. In 2 volumes 12mo.

A new Edition of Homeward Bound; or, the Chase. In 2 volumes 12mo.

HOME AS FOUND, being a Sequel to Homeward Bound. In 2 vols. 12mo.

PRECAUTION,

A NOVEL.

A new edition, revised by the Author. In 2 volumes 12mo.

This excellent novel, the first that issued from the prolific pen of our admired countryman Cooper, had become extremely

scarce, if not entirely out of print. It is one of the very best productions of its author, and from the mountains of trash under the titles of novels, that have accumulated of late years, it may be selected by the youth of our country as worthy of their perusal for its amusing story and pure English style.—*The Lousianian*.

The great care which the author has bestowed upon this edition, is correcting the errors and amending the faults, of the original, will be appreciated by every reader with whom "Precaution" is an old acquaintance. It is now much more worthy of his time and money.—*Evening Gazette and Galaxy*.

NOVELS AND TALES, IN TWENTY-SIX VOLUMES.

CONTAINING:

The Spy.	In 2 vols.
Pioneers.	"
Pilot.	"
Prairie.	"
Lionel Lincoln.	"
Last of the Mohicans.	"
Red Rover.	"

The Wept of the Wish-Ton-Wish.	In 2 vols.
Water Witch.	"
Bravo.	"
Travelling Bachelor	"
Heidenmauer.	"
Headsmen.	"

Or any work may be had separate in boards.

WORKS BY BOZ.

Illustrated editions of the following works by (BOZ) Charles Dickens, Esq. printed on fine paper, and handsomely bound in embossed cloth to match. The plates are engraved by Yeager, and neatly printed on cream coloured paper.

THE POSTHUMOUS PAPERS OF THE PICKWICK CLUB;

WITH NUMEROUS DESIGNS,

ILLUSTRATED BY

SAM WELLER JR., AND ALFRED CROWQUILL, ESQ.

OLIVER TWIST; OR, THE PARISH BOY'S PROGRESS, WITH TWENTY-FOUR ILLUSTRATIONS;

DESIGNED BY CRUIKSHANK.

SKETCHES BY BOZ;

Illustrative of Every-Day Life, and Every-Day People.

A NEW EDITION,

COMPRISING BOTH THE SERIES;

AND EMBELLISHED WITH NUMEROUS ILLUSTRATIONS,

BY GEORGE CRUIKSHANK.

THE LIFE AND ADVENTURES

OF

NICHOLAS NICKLEBY, WITH THIRTY NINE ILLUSTRATIONS BY PHIZ;

AND

A SPLENDID PORTRAIT OF THE AUTHOR.

Cheap editions of the last three mentioned works may also be had, each work containing two plates.

IN PREPARATION.

A new work by Boz, to be issued in numbers, to match the above.

JACK SHEPPARD,

A NEW ROMANCE

BY THE AUTHOR OF

"ROOKWOOD," "CRITCHTON," &c.

In 2 vols. 12mo. With Plates.

"We are glad to welcome the whole of Jack Sheppard at last. It is one of the richest books of the day. W. H. Ainsworth, the author of Jack Sheppard, has in this work, established his reputation as a fine delineator of low life. His novel is second only to the works of the unrivalled 'Boz,' for the deep interest, graphic sketches of character, and striking views of crime and its inevitable consequences. The hero of the tale is the original of 'Captain Macheath,' of Beggar's Opera celebrity."—*Evening Gazette & Galaxy*.

"Jack Sheppard is so worthy of the hand which gave us the famous Dick Turpin and his Ride to York, that we gladly step out of our way to bid him welcome."—*Literary Gazette*.

"In every respect 'Jack Sheppard' promises well. He is full of interest, life, and spirit. The story is at once startling, impressive, and pathetic."—*Courier*.

"The scenes of 'Jack Sheppard' are full of spirit, incident and variety."—*Spectator*.

"Full of that interest which Mr. Ainsworth imparts to all the productions. It bids fare to share that undisputed sway which 'Boz' has exercised in these pages."—*Weekly Chronicle*.

"The Author of 'Rookwood' has here struck out a new vein in fiction. 'Jack Sheppard' will greatly advance even the present high reputation of Mr. Ainsworth."—*Naval and Military Gazette*.

(In preparation,)

THE TOWER OF LONDON, with Illustrations.

GUY FAWKES, with Illustrations.

By Ainsworth.

"SAM SLICK THE CLOCKMAKER."

THE CLOCK-MAKER OR SAYINGS AND DOINGS OF SAMUEL SLICK OF SLICKVILLE, First and second series in 2 vols. 12mo.

** Either Series may be had separately.

Sam Slick's shrewd observations and odd quaint humour are admirable.—*Athenæum*.

Full of spirit, originality and humour.—*Morning Chronicle*.

The humour of Mr. Slick is by far the best thing of its kind we have ever seen. It is indeed truly delightful. Such a book as this stands in need of neither our praise nor recommendation. It will make its own way; for it is of the right stuff.—*Scotsman*.

Sam Slick is one of the most sagacious as well as amusing authors of his time and country.—*Atlas*.

The author of this work, which has furnished the newspapers with so many *bonne bouches* in the way of extracts, is a man of original genius and shrewd observation, with an inexhaustible vein of wit and humour.—*Dublin Mail*.

One of those genuinely original works which carry their own letter of recommendation, and are every where relished.—*Globe*.

A very clever and characteristic work. The humour is rich and dry, the satire broad and sharp, without ill-nature.—*Spectator*.

THE BUBBLES IN CANADA, by Sam Slick, in 1 vol. 12mo.

SAM SLICK'S LETTER BAG OF THE GREAT WESTERN, OR LIFE IN A STEAMER, by the Author of "The Clockmaker," &c. &c. In 1 vol. 12mo.

(In preparation,)

SAM SLICK IN ENGLAND, By the Author of "The Clockmaker," &c. &c.

BRIDGEWATER TREATISES.

This series of Treatises is published under the following circumstances.—

The Right Honourable and Rev. FRANCIS HENRY, Earl of Bridgewater, died in the month of February, 1835; he directed certain trustees therein named, to invest in the public funds, the sum of eight thousand pounds sterling; this sum, with the accruing dividends thereon, to be held at the disposal of the President, for the time being, of the Royal Society of London, to be paid to the person or persons nominated by him. The Testator farther directed, that the person or persons selected by the said President, should be appointed to write, print and publish one thousand copies of a work, on the Power, Wisdom, and Goodness of God, as manifested in the Creation; illustrating such work, by all reasonable arguments, as, for instance, the variety and formation of God's creatures in the Animal, Vegetable, and Mineral Kingdoms; the effect of digestion, and, thereby, of conversion; the construction of the hand of man, and an infinite variety of other arguments; as also by discoveries, ancient and modern, in arts, sciences, and the whole extent of literature.

He desired, moreover, that the profits arising from the sale of the works so published, should be paid to the authors of the works.

The late President of the Royal Society, DAVIES GILBERT, Esq., requested the assistance of his Grace, the Archbishop of Canterbury, and of the Bishop of London, in determining upon the best mode of carrying into effect, the intentions of the Testator. Acting with their advice, and with the concurrence of a nobleman immediately connected with the deceased, Mr. Davies Gilbert appointed the following eight gentlemen to write separate Treatises in the different branches of the subjects here stated:—

- I. The Adaptation of External Nature to the Moral and Intellectual Constitution of Man. By the Rev. THOMAS CHALMERS, D. D., Professor of Divinity in the University of Edinburgh.
- II. The Adaptation of External Nature to the Physical Condition of Man, principally with reference to the supply of his wants, and the exercise of his intellectual faculties. By JOHN KIDD, M. D., F. R. S., Regius Professor of Medicine in the University of Oxford.
- III. Astronomy and General Physics, considered with reference to Natural Theology. By the Rev. WM. WHEWELL, M. A., F. R. S., Fellow of Trinity College, Cambridge.
- IV. The Hand, its mechanism and vital endowments as evincing design, with numerous wood cuts. By SIR CHARLES BELL, K. H. F. R. S.
- V. Chemistry, Meteorology, and the Function of Digestion, considered with reference to Natural Theology. By WM. PROUT, M. D., F. R. S.
- VI. The History, Habits, and Instincts of Animals. By the Rev. WM. KIRBY, M. A., F. R. S., Illustrated by numerous engravings on copper.
- VII. Animal and Vegetable Physiology, considered with reference to Natural Theology. By PETER MARK ROJET, M. D. Illustrated with nearly 500 wood cuts.
- VIII. Geology and Mineralogy considered with reference to Natural Theology. By the Rev. WM. BUCKLAND, D. D., Canon of Christ Church, and Reader in Geology and Mineralogy in the University of Oxford. With eighty-nine copper-plates and large coloured maps.

The whole series forms a beautiful set of books, in seven volumes octavo, and to be had in handsome cloth or half bound with calf backs and corners.

* * The work of Chalmers, Bell, Whewell, Prout and Kidd, may be had separate each in one volume 12mo.

* * The work of Rojet, in 2 vols. 8vo. Buckland, in 2 vols. 8vo., and Kirby in one vol. 8vo., may also be had separate from the sets of the work.

HERSCHEL'S ASTRONOMY.

A TREATISE ON ASTRONOMY. By Sir John F. W. Herschel, F. R. S., &c.

A new Edition, with a Preface and a series of Questions for the examination of Students. By S. C. Walker. In 1 vol. 12mo.

"The present treatise is in no wise inferior to its predecessor: it is characterized by the same agreeable and elegant style, the same facility of illustration—added to which it possesses unrivalled precision and accuracy of demonstration. Avoiding, therefore, the abstruse niceties and the transcendental mathematics of the subject, the author has nevertheless produced a volume calculated, we are fully persuaded, to impress upon his readers the magnitude and importance of the science, and to initiate them in no mean degree into its mysteries."—*Literary Gazette*.

BREWSTER'S OPTICS.

TREATISE ON OPTICS. By Sir David Brewster. A new edition, with Notes and Additions, by A. D. Bache, Professor of Natural Philosophy and Chemistry in the University of Pennsylvania. 12mo.

"The author has given proof of his well-known industry, and extensive acquaintance with the results of science in every part of Europe."—*Monthly Magazine*.

"The subject is, as might be expected, ably treated, and clearly illustrated."—*U. S. Journal*.

EVAN'S MILLWRIGHT'S GUIDE.

MILLWRIGHT AND MILLER'S GUIDE. By Oliver Evans. The Tenth Edition with Additions and Corrections, by the Professor of Mechanics in the Franklin Institute of Pennsylvania; and a Description of an Improved Merchant Flour Mill. With Engravings. By C. & O. Evans, Engineers.

This is a practical work and has had a very extended sale.

BONAPARTE'S ORNITHOLOGY.

American Ornithology, or Natural History of Birds, Inhabiting the United States, by Charles Lucien Bonaparte; designed as a continuation of Wilson's Ornithology, Vols. I. II. III. and IV.

Gentlemen who possess Wilson, and are desirous of rendering the work complete, are informed that the edition of this work is very small, and that but a very limited number of copies remain unsold.

(Odd volumes can be supplied, to complete sets.)

BARTON'S FLORA.

A Flora of North America, with 108 coloured Plates. By W. P. C. Barton, M. D. In 3 vols. 4to.

ARNOTT'S PHYSICS.

Elements of Physics, or Natural Philosophy, General and Medical, explained independently of Technical Mathematics, and containing New Disquisitions and Practical Suggestions. By Neill Arnott, M. D. Fourth American from the fifth London edition, with Additions by Isaac Hays, M. D. In 2 vols. 8vo.

"Dr. Arnott's work has done for Physics as much as Locke's Essay did for the science of mind."—*London University Magazine*.

"We may venture to predict that it will not be surpassed."—*Times*.

"Dr. A. has not done less for Physics than Blackstone did for the Law."—*Morning Herald*.

"Dr. A. has made Natural Philosophy as attractive as Buffon made Natural History."—*French Critic*.

"A work of the highest class among the productions of mind."—*Courier*.

GRAY'S CHEMISTRY.

The Chemistry of the Arts on the basis of Gray's Operative Chemist, being an Exhibition of the Arts and Manufactures dependent on Chemical Principles, with numerous Engravings, by Arthur L. Porter, M. D. late Professor of Chemistry, &c. in the University of Vermont. In 8vo. With numerous Plates.

FARADAY'S MANIPULATIONS.

Chemical Manipulation. Instruction to Students on the Methods of performing Experiments of Demonstration or Research, with accuracy and success. By Michael Faraday, F. R. S. First American from the second London edition, with additions by J. K. Mitchell, M. D.

DELA BECHE'S GEOLOGY.

A GEOLOGICAL MANUEL. By Henry T. de la Beche, Esq., F. R. S., F. G. S., Member of the Geological Society of France, &c. In 1 vol. 8vo., with 104 wood cuts.

LEA'S GEOLOGY.

CONTRIBUTIONS TO GEOLOGY.

Illustrated by numerous coloured Engravings. By Isaac Lea, Member of the American Philosophical Society, &c. &c. In 1 vol. 8vo.

LEA'S NAIADES.

A SYNOPSIS OF THE FAMILY OF NAIADES. By Isaac Lea, Member of the American Philosophical Society, &c. &c. 1 vol. 8vo. with coloured plates.

LARDNER'S HYDROSTATICS.

A TREATISE ON HYDROSTATICS AND PNEUMATICS. By the Rev. Dionysius Lardner, LL. D. F. R. S. &c. A new American from the last London edition, with Notes by Benjamin F. Joslin, M. D., Professor of Natural Philosophy in Union College.

"It fully sustains the favourable opinion we have already expressed as to this valuable compendium of modern science."—*Lit. Gaz.*

"Dr. Lardner has made a good use of his acquaintance with the familiar facts which illustrate the principles of Science."—*Monthly Mag.*

"It is written with a full knowledge of the subject, and in a popular style, abounding in practical illustrations of the abstruse operations of these important sciences."—*U. S. Jour.*

LARDNER'S MECHANICS.

A TREATISE ON MECHANICS. By Captain Kater, and the Rev. Dionysius Lardner. With numerous Engravings. In 1 vol. 12mo.

"A work which contains an uncommon amount of useful information, exhibited in a plain and very intelligible form."—*Olmsted's Nat. Philosophy.*

"Admirable in development and clear in principles, and especially felicitous in illustration from familiar subjects."—*Monthly Mag.*

"Though replete with Philosophical information of the highest order in mechanics, adapted to ordinary capacities in a way to render it at once intelligible and popular."—*Lit. Gazette.*

"A work of great merit, full of valuable information, not only to the practical mechanic, but to the man of science."—*N. Y. Courier and Enquirer.*

RENWICK'S MECHANICS.

ELEMENTS OF MECHANICS. By James Renwick, Esq. Professor of Natural and Experimental Philosophy, Columbia College, N. Y. In 8vo. with numerous Engravings.

"We think this decidedly the best treatise on Mechanics, which has issued from the American press, that we have seen; one, too, that is alike creditable to the writer, and to the state of science in this country."—*American Quarterly Review.*

THE BOOK OF SCIENCE; in three parts, a familiar introduction to the Principles of Natural Philosophy, adapted to the comprehension of Young People. Illustrated by many curious and interesting Experiments and Observations, and including Notices of the most recent Discoveries. Embellished with numerous Engravings on wood.

- | | | |
|-------|---|-------------------|
| Part. | 1 | MECHANICS. |
| " | 2 | HYDROSTATICS, &c. |
| " | 3 | CHEMISTRY. |

REID ON CLOCK MAKING.

TREATISE ON CLOCK AND WATCH-MAKING. Theoretical and Practical. By Thomas Reid, Edinburgh Honorary Member of the Worshipful Company of Clock Makers, London. Royal Octavo. Illustrated by Numerous Plates.

THE NATIONAL SCHOOL MANUAL: a regular and connected Course of Elementary Studies, embracing the necessary and useful Branches of the Common Education. In Four Parts, with a quarto Atlas. Compiled from the latest and most approved Authors, by Mr. R. Bartlett.

The plan of this work was the suggestion of the late Governor Clinton, whose zeal and efforts in the cause of our Public Schools, will be cherished with grateful remembrance to the latest posterity; and this work so far as it had advanced up to the time of his lamented death, received his favourable regard and patronage.

The object of the National School Manual, is to furnish a System of instruction, for a thorough English education, in a plain, practical, and progressive Series of Lessons, collaterally arranged.

THE CLINTON PRIMER.

A
SERIES OF FIRST LESSONS.
WITH
NUMEROUS CUTS.

NEUMANS DICTIONARY.

A Pocket Dictionary of the Spanish and English Languages, Compiled from the last Improved Editions of Neuman and Barretti. In one small volume.

DONNEGAN'S LEXICON.

Greek and English Lexicon. By D. Donnegan. Abridged for the use of Schools. In 1 vol. royal 18mo., containing above 800 pages.

A SELECTION OF ONE HUNDRED PERIN'S FABLES, accompanied by a Key, containing the text, a literal and a free translation, arranged in such a manner as to point out the difference between the French and the English idiom, also a figured pronunciation of the French, according to the best French works extant on the subject; the whole preceded by a short treatise on the sounds of the French language, compared with those of the English. By A. Bolmar. In 1 vol. 12mo.

The expression 'figured pronunciation,' is above employed to express that the words in the Key to the French Fables are spelt and divided as they are pronounced. It is what Walker has done in his Critical Pronouncing Dictionary; for instance, he indicates the pronunciation of the word enough, by dividing and spelling it thus, e-nuf. In the same manner I indicate the pronunciation of the word compa-tient thus, kon-tè. As the understanding of the figured pronunciation of Walker requires the student to be acquainted with the primitive sounds of the English vowels, he must likewise, before he can understand the figured pronunciation of the French, make himself acquainted with the 20 primitive sounds of the French vowels. This any intelligent person can get from a native, or from any body who reads French well, in a few hours.

LES AVENTURES DE TELEMAQUE PAR FENELON, in 1 vol. 12mo., accompanied by a KEY to the EIGHT FIRST BOOKS; in 1 vol. 12mo., containing like the Fables—the Text a Literal and Free Translation; intended as a Sequel to the Fables.

BUTLER'S ANTIENT ATLAS.

ATLAS OF ANTIENT GEOGRAPHY, consisting of 21 Coloured Maps, with a complete Accentuated Index. By Samuel Butler, D. D., F. R. S. &c. Archdeacon of Derby.

A new edition revised and corrected, in one royal 8vo. volume.

Extract of a Letter from Professor Stuart of Andover.

"I have used Butler's Atlas Classica for 12 or 14 years, and prefer it on the score of convenience and correctness to any atlas within the compass of my knowledge. It is evidently a work of much care and taste, and most happily adapted to classical readers and indeed all others, who consult the history of past ages. I have long cherished a strong desire to see the work brought forward in this country, and I am exceedingly gratified that you have carried through this undertaking. The beautiful manner in which the specimen is executed that you have sent me does great credit to engravers and publishers. It cannot be that our schools and colleges will fail to adopt this work, and bring it into very general circulation. I know of none which in all respects would supply its place."

"The abridged but classical and excellent work of Butler, on Antient Geography, which you are printing, as an accompaniment to the maps, I consider one of the most attractive works of the kind, especially for young persons studying the classics, that has come under my notice. I wish you the most ample success in these highly useful publications."

A Collection of Colloquial Phrases on every Topic necessary to maintain Conversation, arranged under different heads, with numerous remarks on the peculiar pronunciation and use of various words—the whole so disposed as considerably to facilitate the acquisition of a correct pronunciation of the French. By A. Bolmar. 1 vol. 18mo.

A COMPLETE TREATISE ON THE GENDERS OF FRENCH NOUNS; in a small pamphlet of fourteen pages.

This little work, which is the most complete of the kind, is the fruit of great labour, and will prove of immense service to every learner.

ALL THE FRENCH VERBS BOTH REGULAR AND IRREGULAR, in a small volume.

The verbs être to be, avoir to have, parler to speak finir to finish, recevoir to receive, vendre to sell, se lever to rise, se bien porter to be well, s'en aller to go away, are here all conjugated through—affirmatively—negatively—interrogatively—and negatively and interrogatively—an arrangement which will greatly facilitate the scholar in his learning the French verbs, and which will save the master the trouble of explaining over and over again what may be much more easily learned from books, thus leaving him more time to give his pupil, during the lesson, that instruction which cannot be found in books, but which must be learned from a master.

BUTLER'S ANTIENT GEOGRAPHY.

GEOGRAPHIA CLASSICA, or the application of Antient Geography to the Classics. By Samuel Butler, &c. &c. Fourth American edition with questions. In 1 vol.

THE NOVELS OF JANE AUSTEN:

CONTAINING

PRIDE AND PREJUDICE,
MANSFIELD PARK,
PERSUASION,

SENSE AND SENSIBILITY,
EMMA, AND
NORTHANGER ABBEY.

COMPLETE IN ONE LARGE VOLUME.

Bound in Elegant Embossed Cloth, or neatly half bound with Calf backs and corners.

DR. BIRD'S NOVELS.

THE ADVENTURES

OF

ROBIN DAY.

IN 2 VOLS. 12MO.

WE have long been accustomed to regard the fictions of this author with peculiar respect for their energetic action. The dramatic power of the "novel," in its outlines, the writer of "Calavar" understands surpassingly well; and if in the detail he is sometimes untrue to taste, he is seldom if ever faithless to nature, in what he describes. This constitutes, in truth, a charm in what BIRD writes, that is easily appreciated and understood. It is his design to be simple in portraiture, and plain in expression! He succeeds, because he draws his transcripts, where colours cannot fail—from *life*. But in his gust for bold drawing and large effect, he sometimes rejects the *finish* of the master, though he never leaves the reader in doubt of his capacity.

NICK OF THE WOODS; or, The Jibbenainosay. A Tale of Kentucky. In two vols. 12mo.

PETER PILGRIM;

OR,

A RAMBLER'S RECOLLECTIONS.

In 2 vols. 12mo.

It is a sort of Decameron of Tales, of which the scenes are laid in various parts of our country, and it abounds in rich traditional lore and national characters.

THE HAWKS OF HAWK HOLLOW; A Tradition of Pennsylvania.—
In two vols. 12mo.

A Third Edition of

CALAVAR:

OR, THE KNIGHT OF THE CONQUEST.

In two vols. 12mo.

THE INFIDEL: or, The Knight of the Conquest. A Romance. By the Author of "Calavar." Two vols. 12mo.

A New and Handsome Edition of

PICCIOLA, THE PRISONER OF FENESTRELLA;

OR, CAPTIVITY CAPTIVE.

By M. D. Saintine. In one neat royal 18mo. volume.

In fancy paper binding.

For this beautiful little novel we may venture to predict a degree of popularity in the English world of letters equal to that enjoyed by "Paul and Virginia," "Elizabeth," and

one or two other favourites. It is a production of every class, and for every shade of taste.—*Court Journal*.

ROB OF THE BOWL;

OR,

A LEGEND OF ST. INIGOE.

By the Author of "Horse Shoe Robinson," &c. &c.
In two vols. 12mo.

"The hero, 'Rob of the Bowl,' is a finely executed conception. The perfect originality of the design, the bold colouring and elaborate finish of this character, stamp the author as an artist of no ordinary power. He has now produced an able work that will be read with pleasure wherever the English language is understood. His well known powers of description render his account of the eventful scenes of the early colonists most life-like and pleasingly exact.—*Gentleman's Magazine*.

HORSESHOE ROBINSON.

A TALE OF THE TORY ASCENDENCY.

By the Author of "SWALLOW BARN."
Fourth Edition. Two Volumes 12mo.

ALTHEA VERNON, the Embroidered Handkerchief, together with Henrietta Harrison, or the Blue Cotton Umbrella. By Miss Leslie.

PENCIL SKETCHES.

Or, Outlines of Character and Manners.
BY MISS LESLIE.

ISABEL, OR SICILY;

A Pilgrimage. By Henry T. Tuckerman. Author of "Italian Sketch Book." In 1 volume.

This is the production of a good scholar and a man of taste. It contains much delicacy of sentiment, prettiness of thought, and beauty of expression. The ideas and descriptions of Sicilian scenery, manners, history, &c. are conveyed in the form of graceful colloquy, by which many unimportant details are avoided, and the narrative of a connected pilgrimage preserved.—*Madisonian*.

THE COURT AND THE BASTILE,
by M. D. Saintine, Author of Picciola, &c.
(In preparation.)

SIMMS' LAST NOVEL.

The Damsel of Darien, a novel, by the author of 'The Yemassee.' 'Guy Rivers,' 'Mellichampe,' &c., in 2 vols. 12mo.

"Seldom have we read a tale which has so perfectly absorbed us by its interest. Mr. Simms has based his story upon the adventures of Vasco Nunes de Balboa, as some have called him, although our author calls him Vasco Nunes—Ojeda and Nicesa, as chiefly detailed in Irving's 'Companions of Columbus,' and from these with the aid of his vivid imagination to furnish the details of society and life, he has composed what may be considered with truth, the Romance of American History.

"Our author's descriptive powers are of the very first order, and in his sketches of character there is no writer that better understands the art of bringing out the salient features in high relief, and of impressing an individuality upon the portraiture that fixes the image perfectly in our mind, until it haunts us like the figure of one we have intimately known. In this novel, 'The Damsel of Darien.' Mr. Simms has opened a new and virgin mine in the treasures of Romance. An unploughed field, which only needs cultivation to produce the richest fruits.—*Evening Gazette and Galaxy*."

STANLEY;

Or the Recollections of a Man of the World.

By an unknown writer. In 2 volumes 12mo.

Stanley is far above the common class of novels. It is evidently the production of a ripe scholar and deep thinker; one who has seen much and meditated more, and who, from the affluence of his stores, sometimes becomes profuse in the distribution of his knowledge. The criticisms in which the work abounds, are original, vigorous, searching, and profound—and the characters of several of the most distinguished writers of the present century are drawn with singular felicity. Altogether it is the effort of a highly cultivated mind, rich alike in natural endowments and acquired treasures.—*Saturday News*.

SHELLEY'S PROSE WORKS.

ESSAYS.—LETTERS FROM ABROAD, Translations and Fragments of Percy Bysshe Shelley, edited by Mrs. Shelley. In 2 vols. 12mo.

SHAKSPEARE AND HIS FRIENDS;

OR,

THE GOLDEN AGE OF MERRY ENGLAND.

BY THE AUTHOR OF

"THE YOUTH OF SHAKSPEARE."

"We have been better pleased with this work than with any we have met with for a long time. The ground upon which the author has ventured is fearfully full of difficulties, but he has threaded his way with considerable skill. Shakspeare, Burbage, Ben Jonson, Sir W. Raleigh, Lord Burghley, Elizabeth, are brought in vivid portraiture before us; the generous disposition, ready wit, and noble bearing of the immortal poet being pictured with rare success."

To match the above, and by the same author.

"THE YOUTH OF SHAKSPEARE;"

THE GOVERNESS, a novel by the Countess of Blessington, in 2 vols. 12mo.

THE CONFESSIONS OF AN ELDERLY LADY AND GENTLEMAN, by the Countess of Blessington, in 2 vols. 12mo.

SOLOMON SEESAW. A Novel. By J. P. Robertson, author of "Letters on Paraguay." In two vols. 12mo.

TRIALS OF THE HEART; By Mrs. Bray, author of "Trelawney," "The Talba," &c. In two vols. 12mo.

THE NAVAL FOUNDLING: by the Old Sailor, author of "Tough Yarns," "Nights at Sea," "Greenwich Hospital," &c. In 3 vols. 12mo.

THE HUSBAND HUNTER: By Dennis Moriarty, Esq., author of "The Wife Hunter," &c. &c. In two vols. 12 mo.

THE DUKE.

A NOVEL.

BY MRS. GREY.

In two vols. 12mo.

HAMILTON KING; or, THE SMUGGLER AND THE DWARF. By the Old Sailor, author of "Tough Yarns," "Stories of Greenwich Hospital," &c. In two vols. 12mo.

WILHELM MEISTER.

A New Translation by Carlisle.

(In Preparation.)

GURNEY MARRIED,
A SEQUEL TO GILBERT GURNEY.

By the Author of

"SAYINGS AND DOINGS," "JACK BRAG,"
&c. &c.

In two volumes 12mo.

BIRTHS, DEATHS, & MARRIAGES.

By the Author of

"SAYINGS AND DOINGS," "JACK BRAG."
&c. &c.

In two volumes 12mo.

COUSIN GEOFFREY,
THE OLD BACHELOR.

BY THEODORE HOOK.

In two vols. 12mo.

(In Preparation.)

POE'S TALES.

Tales of the Grotesque and Arabesque, by Edgar A. Poe, in 2 vols. 12mo.

"To say we have read this production attentively is not enough. We have studied it. It is every way worthy of such a distinction, and whoever shall give it a careful study and a philosophical analysis, will find in it the evidences of an original, vigorous, and independent mind, stored with rich and various learning, and capable of successful application to a great variety of subjects."

NOVELS, TALES AND ROMANCES.

- Austin's Novels, Complete in one Royal Svo. volume.
- American Lounger, by the Author of La-fitte, 12mo.
- Althea Vernon, by Miss Leslie, 12mo.
- Adam Buff, or Other Men of Character.
- Boz's Sketches of Every-day Life and Every-day People, 1 vol. Various editions.
- Births, Deaths, and Marriages, by Theodore Hook, 2 vols. 12mo.
- Coopers Novels and Tales.
- " Spy.
 - " Pioneers.
 - " Pilot.
 - " Prairie.
 - " Lionel Lincoln.
 - " Last of the Mohicans.
 - " Red Rover.
 - " Wept of the Wish-Ton-Wish.
 - " Water-Witch.
 - " Bravo.
 - " Travelling Bachelor.
 - " Headsman.
 - " Heidenmauer.
 - " Homeward Bound.
 - " Home as Found.
 - " Pathfinder.
 - " Precaution.
- Calavar, by Dr. Bird, 2 vols.
- Country Stories, by Miss Mitford, 1 vol.
- Confessions of an Elderly Lady and Gentleman, by Lady Blessington, 2 vols.
- Concealment, 2 vols. 12mo.
- Collin Clink, 2 vols. 12mo.
- Clinton Bradshaw, 2 vols.
- Calderon, by Bulwer, 1 vol. 12mo.
- Cousin Geoffrey, by Hook.
- Damsel of Darien, by Simms, 2 vols. 12mo.
- East and West, 2 vols.
- Elvira, the Nabob's Wife, 2 vols. 12mo.
- Governess, by Lady Blessington, 12mo.
- Gurney Married, by Theodore Hook.
- Harry Austin, by an Officer, 2 vols. 12mo.
- Hawks of Hawk Hollow, by Dr. Bird,
- Horseshoe Robinson, by the Author of Swallow Barn, 2 vols. 12mo.
- Hawk Chief, by the Author of Indian Sketches, 2 vols. 12mo.
- Hamilton King, by the Old Sailor, 2 vols.
- Horace Vernon, or Fashionable Life.
- Husband Hunter, by the Author of the Wife Hunter, 2 vols. 12mo.
- Infidel, by Dr. Bird, 2 vols.
- Isabel or Sicily, by Tuckerman, 1 vol.
- Jack Sheppard, by Ainsworth, 2 vols. 12mo. With plates.
- James's Book of the Passions.
- Jane Lomax, by Horace Smith, 2 vols.
- Land Sharks and Sea Gulls, by Captain Glascock, 2 vols. 12mo.
- Little Frenchman and his Water Lots, by G. P. Morris, 12mo., with Illustrations.
- Mrs. Armytage, or Female Domination, by Mrs. Gore 2 vols.
- Mary Raymond, by Mrs. Gore, 2 vols.
- Naval Foundling, by the Old Sailor, 12mo.
- Nicholas Nickleby, by Boz. Various editions.
- Nick of the Woods, or the Jibbenainosay, a Tale of Kentucky, by Dr. Bird.
- Oliver Twist, by Boz. Various editions.
- Pascal Bruno, by Theodore Hook, 1 vol.
- Picciola, or Captivity Captive, 18mo.
- Pickwick Club, chronicled by Boz. Various editions.
- Peter Pilgrim, by Dr. Bird, 2 vols. 12mo.
- Pencil Sketches, by Miss Leslie, 2d series.
- Do. do. 3d. series.
- Rattlin the Reefer. In 2 vols. 12mo.
- Robin Day, by Dr. Bird, 2 vols. 12mo.
- Rob of the Bowl, by the Author of Horse-shoe Robinson, 2 vols, 12mo.
- Shakespeare and His Friends, 3 vols. 12mo.
- Soloman Seesaw, 2 vols. 12mo.
- Stanley, or the Adventures of a Gentleman, 2 vols. 12mo.
- Sedgewick's Tales and Sketches, 1 vol.
- Swallow Barn, by Kennedy. In 2 vols.
- The Duke, by Mrs. Grey. In 2 vols. 12mo.
- Trials of the Heart, by Mrs. Bray, 2 vols. 12mo.
- Tales of the Grotesque and Arabesque, by Edgar A. Poe, 2 vols. 12mo.
- Wood Leighton, by Mary Howitt, 12mo.
- Youth of Shakespeare. In 12mo.

PREPARING FOR PUBLICATION.

The Novelists Library, embracing in several large super royal volumes, such of the best novels and romances as may be deemed worthy of preservation in a form suitable for a Library of Standard Literature.

MEDICAL AND SURGICAL BOOKS.

PUBLISHED

BY

LEA & BLANCHARD,
PHILADELPHIA.

THE
AMERICAN

JOURNAL OF THE MEDICAL SCIENCES,

EDITED BY
ISAAC HAYS, M. D.

SURGEON TO WILLS HOSPITAL, &c. &c.

TERMS.

Each number contains 260 pages, or upwards, and is frequently illustrated by coloured engravings. It is published on the first of November, February, May and August. Price Five Dollars per annum, payable in advance.

Orders, enclosing the amount of one year's subscription, addressed to the publishers, or any of the agents, will receive prompt attention. The year of this work commences with the November number.

Persons sending Twenty Dollars will be entitled to five copies of the work, to be forwarded as they may direct. All persons desirous of advancing the interest of medical science, are requested to use their efforts to increase its circulation.

The postage per number is, within 100 miles, about 16 cents; over 100 miles, about 28 cents.

A few complete sets may be had at a large discount from the subscription price. Odd numbers can be furnished to complete sets.

The following Extracts show the estimation in which the Journal is held.

"Several of the American Journals are before us. *** Of these, the American Journal of the Medical Sciences is by far the better periodical; it is, indeed, the best of the trans-atlantic medical publications; and, to make a comparison nearer home, is in most respects superior to the great majority of European works of the same description."—*The London Lancet*.

"We need scarcely refer our esteemed and highly eminent contemporary, [The American Journal of Medical Sciences,] from whom we quote, to our critical remarks of the opinions of our own countrymen, or to the principles which influence us in the discharge

of our editorial duties."—"Our copious extracts from his unequalled publication, unnoticing multitudes of others which come before us, are the best proof of the esteem which we entertain for his talents and abilities."—*London Medical and Surgical Journal*!

"The Medical Journal of Medical Sciences is one of the most complete and best edited of the numerous periodical publications of the United States."—*Bulletin des Sciences Medicales*, tome xlv.

"The Medical Journal of Medical Sciences is conducted with distinguished ability. Published in one of the most literary cities in our country, and supported by a number of her most gifted and best educated physicians, its reputation is deservedly high as well abroad as at home."—*Pennsylvania Journal*.

MANUAL OF MATERIA MEDICA
AND PHARMACY, By H. M. EDWARDS,
M. D. and P. VASSEUR, M. D.

CHEMICAL MANIPULATION. In-
struction to Students on the Methods of
performing Experiments of Demonstration
or Research, with accuracy and success.
By MICHAEL FARRIDAY, F. R. S. First
American, from the second London edition,
with additions by J. K. MITCHELL, M. D.

A FLORA OF NORTH AMERICA,
with 108 coloured Plates. By W. P. C.
BARTON, M. D. In 3 vols. 4to.

A MEDICAL ACCOUNT OF
THE MINERAL SPRINGS
OF VIRGINIA.

By Professor Gibson. (In preparation.)

A MANUAL OF MEDICAL JU-
RISPRUDENCE. By Professor R.
E. Griffith. In one volume. (Now
preparing.)

THE PRINCIPLES AND PRACTICE
OF MEDICINE. By Professor Dungli-
son. In two volumes, octavo. (In prepa-
ration.)

A NEW DICTIONARY, OF MEDICAL SCIENCE AND LITERATURE.

A NEW EDITION,

Completely Revised, with Numerous Additions and Improvements,

OF

DUNGLISON'S DICTIONARY

OF

MEDICAL SCIENCE AND LITERATURE:

CONTAINING

A concise account of the various Subjects and Terms, with a vocabulary of Synonyms in different languages, and formulæ for various officinal and empirical preparations, &c.

IN ONE ROYAL 8vo. VOLUME.

"The present undertaking was suggested by the frequent complaints, made by the author's pupils, that they were unable to meet with information on numerous topics of professional inquiry,—especially of recent introduction,—in the medical dictionaries accessible to them.

It may, indeed, be correctly affirmed, that we have no dictionary of medical subjects and terms which can be looked upon as adapted to the state of the science. In proof of this the author need but to remark, that he has found occasion to add several thousand medical terms, which are not to be met with in the only medical lexicon at this time in circulation in the country.

The present edition will be found to contain many hundred terms more than the first, and to have experienced numerous additions and modifications.

The author's object has not been to make the work a mere lexicon or dictionary of terms, but to afford, under each, a condensed view of its various medical relations, and thus to render the work an epitome of the existing condition of medical science."

"To execute such a work requires great erudition, unwearied industry, and extensive research, and we know no one who could bring to the task higher qualifications of this description than Professor Dunglison."—*American Medical Journal*.

"This is an excellent compilation, and one that cannot fail to be very much referred to. It is the best medical lexicon in the English language that has yet appeared. We do not know any volume which contains so much information in a small compass. The Bibliographical notices, though so short, are very important and useful; and altogether we can recommend to every medical man to have this work by him, as the cheapest and best dictionary of re-

ference he can have."—*London Medical and Surgical Journal*.

"So far as we have been able to examine this Dictionary, it is exceedingly thorough and correct, not only in matters purely medical, but in whatever can fairly be arranged in the various branches of science, collateral or contributory to Medicine and Surgery."—*Medical Magazine*.

"So well known are the merits of this valuable work, that, in noticing a second edition of it, it will suffice to extract the remark of the author in the preface, 'that it will be found to contain many hundred terms more than the first, and to have experienced numerous additions and modifications.' It has been got up by the publishers in very handsome style, and must command, as it deserves, an extended circulation."—*Medical Examiner*.

"It is wholly unnecessary, we apprehend, to enter into a long or formal statement of the fact, that Dr. Dunglison's Dictionary, from the first day of its appearance, has been regarded with peculiar favour. And we have now a revised edition, constructed under the immediate eye of the author, who is most favourably circumstanced for adding to the previous edition whatever could give it additional claims on the score of accuracy. Here are eight hundred and twenty-one pages, large octavo, in double columns, distinct type, of which no one ought to complain. Finally, although most of our readers may be owners of the first edition, we cordially and conscientiously recommend to all future purchasers to procure this in preference to any medical lexicon extant. Its true and sterling value, as a key to medical science, and its moderate price, are so many common-sense recommendations which should not be forgotten."—*Boston Medical and Surgical Journal*.

A NEW AND VALUABLE WORK

FOR

PHYSICIANS, APOTHECARIES, AND STUDENTS.

NEW REMEDIES,*The Method of Preparing & Administering them;***THEIR EFFECTS**

UPON THE

HEALTHY AND DISEASED ECONOMY,
&c. &c.

BY ROBLEY DUNGLISON, M. D.

Professor of the Institutes of Medicine and Materia Medica in Jefferson Medical College of Philadelphia; Attending Physician to the Philadelphia Hospital, &c.

IN ONE VOLUME, OCTAVO.

"The value of this book is hardly to be estimated; to be without it, would be very much like obstinacy, and amount to the same thing as saying, like the Austrians in regard to their government, nothing can be improved, for we already live in a state of perfection. Dr. Dunglison, the author, has done an essential service to all classes of practitioners. It is creditable to the industry and wise discrimination of the author, and quite necessary to the libraries of those who feel the necessity of keeping pace with the improvements and discoveries in the broad but imperfectly explored domain of medicine."—*Boston Medical and Surgical Journal*.

*A Third Edition, Improved and Modified, of***DUNGLISON'S****HUMAN PHYSIOLOGY:***Illustrated With Numerous Engravings.*

IN TWO VOLUMES, OCTAVO.

"We are happy to believe that the rapid sale of the last edition of this valuable work may be regarded as an indication of the extending taste for sound physiological knowledge in the American schools: and what we then said of its merits, will show that we regarded it as deserving the reception it has experienced. Dr. Dunglison has, we are glad to perceive, anticipated the recommendation which we gave in regard to the addition of references, and has thereby not only added very considerably to the value of his work, but has shown an extent of reading which, we confess, we were not prepared by his former edition to expect. He has also availed himself of the additional materials supplied by the works that

have been published in the interval, especially those of Müller and Burdach. So that as a collection of details on human physiology alone, we do not think that it is surpassed by any work in our language: and we can recommend it to students in this country (England) as containing much with which they will not be likely to meet elsewhere."—*British and Foreign Medical Review*.

This work exhibits another admirable specimen of American industry and talent, and contains an account of every discovery in Europe up to the period of a few months prior to its publication. Many of the author's views are original and important."—*Dublin Journal of Medical Sciences*.

GENERAL THERAPEUTICS;

OR,

PRINCIPLES OF MEDICAL PRACTICE.

With Tables of the Chief Remedial Agents and their Preparations, and of the Different Poisons and their Antidotes.

By ROBERT DUNGLISON, M. D., &c., &c.

One Volume, large 8vo.

"There being at present before the public several American works on Therapeutics, written by physicians and teachers of distinction, it might be deemed unjust in us, and would certainly be invidious, to pronounce any of them superior to the others. We shall not, therefore, do so. If there be, however, in the English language, any work of the kind more valuable than that we have been examining, its title is unknown to us.

"We hope to be able to give such an account of the work as will strengthen the desire and determination of our readers to seek for a farther acquaintance with it, by a candid perusal of the volume itself. And, in so doing, we offer them an assurance that they will be amply rewarded for their time and labour."—*Transylvania Journal*, Vol. IX, No. 3.

THE MEDICAL STUDENT; or, *Aids to the Study of Medicine.* Including a Glossary of the Terms of the Science, and of the Mode of Prescribing; Bibliographical Notices of Medical Works; the Regulations of the Different Medical Colleges of the Union, &c. By Robley Dunglison, M. D., &c., &c. In one volume, 8vo.

ELEMENTS OF HYGIENE; on the Influence of Atmosphere and Locality; Change of Air and Climate, Seasons, Food, Clothing, Bathing, Sleep, Corporeal and Intellectual Pursuits, &c., on Human Health, Constituting Elements of Hygiene. By Robley Dunglison, M. D., &c., &c. In 1 vol. 8vo.

MEDICAL ESSAYS.

THE CYCLOPEDIA OF PRACTICAL MEDICINE AND SURGERY,

Or Essays on ASTHMA, APHTHÆ, ASPHYXIA, APOPLEXY, ARSENIC, ATROPA, AIR, ABORTION, ANGINA-PECTORIS, and other Subjects Embraced in the Articles from A to Azote, prepared for the Cyclopaedia of Practical Medicine by

Dr. Chapman,	Dr. Dewees,	Dr. Bache,	Dr. Griffith,
Dr. Jackson,	Dr. Hays,	Dr. Coates,	Dr. Harris,
Dr. Horner,	Dr. Dunglison,	Dr. Condie,	Dr. Warren,
Dr. Hodge,	Dr. Mitchell,	Dr. Emerson,	Dr. Patterson,
Dr. Wood,		Dr. Geddings,	

Each article is complete within itself, and embraces the practical experience of its author, and as they are only to be had in this collection will be found of great value to the profession.

** The two volumes are now offered at a price so low, as to place them within the reach of every practitioner and student.

GIBSON'S SURGERY.

A NEW EDITION OF GIBSON'S SURGERY.

THE INSTITUTES AND PRACTICE OF SURGERY; being the Outlines of a Course of Lectures. By William Gibson, M. D., Professor of Surgery in the University of Pennsylvania, &c. &c. Fifth edition, greatly enlarged. In 2 vols. 8vo. With thirty plates, several of which are coloured.

"The author has endeavoured to make this edition as complete as possible, by adapting it to the present condition of surgery, and to supply the deficiencies of former editions by adding chapters and sections on subjects not hitherto treated of. And, moreover, the arrangement of the work has been altered by transposing parts of the second volume to the first, and by changing entirely the order of the subject in the second volume. This has been done for the purpose of making the surgical course in the university correspond with the anatomical lectures, so that the account of surgical diseases may follow immediately the anatomy of the parts."

DEWEES'S WORKS.

A PRACTICE OF PHYSIC, comprising most of the diseases not treated of in Diseases of Females and Diseases of Children. By W. P. Dewees, M. D., formerly adjunct professor in the University of Pennsylvania. In one volume, octavo.

A COMPENDIOUS SYSTEM OF MIDWIFERY.

By DR. DEWEES.

Chiefly designed to facilitate the Inquiries of those who may be pursuing this branch of Study. Illustrated by occasional cases and with many plates. The ninth edition, with additions and improvements. In one vol. 8vo.

DEWEES ON THE DISEASES OF FEMALES.

The seventh edition, Revised and Corrected. With additions, and Numerous plates. In one vol. 8vo.

DEWEES ON THE PHYSICAL AND MEDICAL TREATMENT OF CHILDREN.

With Corrections and Improvements. The seventh ed. In one volume, 8vo.

The objects of this work are, 1st, to teach those who have the charge of children, either as parent or guardian, the most approved methods of securing and improving their physical powers. This is attempted by pointing out the duties which the parent or the guardian owes for this purpose, to this interesting but helpless class of beings, and the manner by which their duties shall be fulfilled. And 2d, to render available a long experience to these objects of our affection when they become diseased. In attempting this, the author has avoided as much as possible, "technicality;" and has given, if he does not flatter himself too much, to each disease of which he treats, its appropriate and designating characters, with a fidelity that will prevent any two being confounded together, with the best mode of treating them, that either his own experience or that of others has suggested.

HORNER'S SPECIAL ANATOMY.

A Treatise on Special and General Anatomy. By W. E. Horner, M. D., Professor of Anatomy in the University of Pennsylvania, &c. &c. Fifth edition, Revised, and much improved. In two volumes, 8vo.

ELLIS' MEDICAL FORMULARY.

The Medical Formulary, being a collection of prescriptions derived from the writings and practice of many of the most eminent Physicians in America and Europe. To which is added an appendix, containing the usual Dietetic preparations and Antidotes for Poisons, the whole accompanied with a few brief Pharmacuetic and Medical observations. By Benjamin Ellis, M. D., Fifth edition, with additions. In one vol.

Broussais on Inflammation, 2 vols. 8vo.

Broussais' Pathology, 1 vol. 8vo.

Colles' Surgical Anatomy, 1 vol. 8vo.

Costers' Physiological Practice, 1 vol. 8vo.

Greys' Chemistry applied to the Arts, 2 vols. with numerous plates.

ELEMENTS OF PHYSICS, OR NATURAL PHILOSOPHY, GENERAL AND MEDICAL, explained independently of TECHNICAL MATHEMATICS, and containing New Disquisitions and Practical Suggestions. By NEIL ARNOTT, M. D. In two volumes, octavo.

"Dr. Arnott's work has done for Physics as much as Locke's Essays did for the science of mind."—*London University Magazine*.

"We may venture to predict that it will not be surpassed."—*Times*.

"Dr. A. has not done less for Physics than Blackstone did for the Law."—*Morning Herald*.

"Dr. A. has made Natural Philosophy as attractive as Buffon made Natural History."—*French Critic*.

"A work of the highest class among the productions of mind."—*Courier*.

ROGET'S PHYSIOLOGY AND PHRENOLOGY.

OUTLINES OF PHYSIOLOGY;

WITH AN

APPENDIX ON PHRENOLOGY;

BY P. M. ROGET, M. D.

Professor of Physiology in the Royal Institute of Great Britain, &c. &c.

FIRST AMERICAN EDITION,

Revised, with numerous notes,

In one volume, 8vo.

From the American Preface.—"Of the Author's qualifications as a physiological writer it is scarcely requisite to speak. The fact of his having been selected to compose the Bridgewater Treatise on Animal and Vegetable Physiology, is sufficient evidence of the reputation which he then enjoyed; and the mode in which he executed the task amply evinces that his reputation rested on a solid basis.

"The present volume contains a concise, well-written epitome of the present state of Physiology—human and comparative—not, as a matter to be expected, the copious details and developments to be met with in the larger treatises on the subject; but enough to serve as an accompaniment and guide to the physiological student.

"The attention of the American Editor has been directed to the revision and correction of the text; to the supplying, in the form of notes, of omissions; to the rectification of some of the points that appeared

to him erroneous or doubtful, and to the furnishing of references to works in which the physiological inquirer might meet with more ample information.

"In Phrenology, the Author is a well-known unbeliever, and his published objections to the doctrine have been regarded as too cogent to be permitted to pass unheeded. It will be seen on farther examination in the interval of many years, which has elapsed since the publication of the sixth edition of the Encyclopædia, has not induced him to modify his sentiments on this head. On the contrary, he appears to be as satisfied at this time, of the fallacy of the positions of the Phrenologist, as he was at any former period."

✂ This work will be introduced into many of the Medical Colleges of the union as a Text Book, it being a cheap volume, and well fitted as an introduction to the larger works on Physiology.

COATES POPULAR MEDICINE:

POPULAR MEDICINE;

OR, FAMILY ADVISER.

Consisting of outlines of Anatomy, Physiology, and Hygiene, with such Hints on the Practice of Physic, Surgery, and the Diseases of Women and Children, as may prove useful in families when regular Physicians cannot be procured: Being a Companion and Guide for intelligent Principals of Manufactories, Plantations, and Boarding Schools; Heads of Families, Masters of Vessels, Missionaries, or Travellers, and a useful Sketch for Young Men about commencing the Study of Medicine.

BY REYNELL COATES, M. D.

Fellow of the College of Physicians of Philadelphia—Honorary Member of the Philadelphia Medical Society—Correspondent of the Lyceum of Natural History of New York—Member of the Academy of Natural Sciences of Philadelphia—Formerly Resident Surgeon of the Pennsylvania Hospital, &c.

Assisted by several Medical friends. In One Volume,

"It is with great satisfaction that we announce this truly valuable compilation, as the most complete and interesting treatise on Popular Medicine ever presented to the public. Simple and unambitious in its language, free from the technicalities, and embracing the most important facts on Anatomy, Physiology and Hygiene, or the art of preserving health; and the treatment of those affections which require immediate attention, or are of an acute character, this should be in the hands of every one, more particularly of those who, by their situations are prevented from resorting to the advice of a physician, nor would the careful perusal

of its pages fail to profit the inhabitants of our cities, by giving them a more accurate knowledge of the structure of the human frame, and the laws that govern its various functions; whose perfect integrity is absolutely essential to health, and even to existence; the various systems of medical charlatanry, daily imagined to take advantage of the credulity and ignorance of mankind, would be rendered far less prejudicial to the community than they now are. We would particularly direct attention to the Chapter on Hygiene, a science in itself of the utmost importance, and ably treated in the small space allowed to it in this volume."—*New York American*.

DR. CLARK ON CONSUMPTION.

A Treatise on Pulmonary Consumption, comprehending an inquiry into the Nature, Causes, Prevention, and Treatment of Tuberculous and Scrofulous Diseases in General. By James Clark, M. D., F. R. S.

As a text-book and guide to the inexperienced practitioner we know none equal to it in general soundness and practical utility—to the general as well as to the professional reader, the work will prove of the deepest interest, and its perusal of unequivocal advantage."—*British and Foreign Medical Review*.

"The work of Dr. Clark may be regarded as the most complete and instructive Treatise on Consumption in the English Language."—*Edinburgh Medical and Surgical Journal*.

CHITTY'S JURISPRUDENCE.

A Practical Treatise on Medical Jurisprudence, with so much of Anatomy, Physiology, Pathology, and the Practice of Medicine and Surgery, as are essential to be known by Members of the Bar and Private Gentlemen; and all the laws relating to Medical Practitioners; with explanatory plates. By J. Chitty, Esq. Second American edition: with Notes and Additions, adapted to American works and Judicial Decisions. 8vo.

A TREATISE ON THE PRACTICE OF MEDICINE, or a Systematic Digest of the Principles of General and Special Pathology and Therapeutics. By E. Geddings, (now preparing.)

SMITH ON FEVER.

A Treatise on Fever. By Southwood Smith, M. D., Physician to the London Fever Hospital. Fourth American edition. In 1 volume 8vo.

FITCH'S DENTAL SURGERY.

A Treatise on Dental Surgery. Second edition, revised, corrected, and improved, with new plates. By S. S. Fitch, M. D. 1 vol. 8vo.

ABERCROMBIE ON THE BRAIN.

Pathological and Practical Researches on Diseases of the Brain and Spinal Cord. Second American, from the third Edinburgh edition, enlarged. By John Abercrombie, M. D. In 1 volume 8vo.

ABERCROMBIE ON STOMACH.

Pathological and Practical Researches on Diseases of the Stomach, the Intestinal Canal, the Liver, and other Viscera of the Abdomen. By John Abercrombie M. D., third American from the second London edition enlarged. In 1 vol. 8vo.

EWELL'S MEDICAL COMPANION.

The Medical Companion or Family Physician: treating of the Diseases of the United States, with their symptoms, causes, cure, and means of prevention.

BERTIEN ON THE HEART.

A Treatise on Diseases of the Heart and Great Vessels. By J. R. Bertien. Edited by G. Bouillaud. Translated from the French. 8vo.

BOISSEAU ON FEVER.

Physiological Pyretology; or a Treatise on Fevers, according to the Principles of the New Medical Doctrine. By F. G. Boisseau, Doctor in Medicine of the Faculty of Paris, &c. &c. From the fourth French edition. Translated by J. R. Knox, M. D. 1 vol. 8vo.

HUTIN'S MANUAL.

Manual of the Physiology of Man; or a concise Description of the Phenomena of his Organization. By P. Hutin. Translated from the French, with notes, by J. Togno. In 12mo.

BELL ON THE TEETH.

The Anatomy, Physiology, and Diseases of the Teeth. By Thomas Bell, F. R. S., F. L. S. &c., third American edition. In 1 vol. 8vo. With numerous plates.

WILLIAMS ON THE LUNGS.

A Rational Exposition of the Physical Signs of Diseases of the Lungs and Pleura; Illustrating their Pathology and facilitating their Diagnosis. By Charles J. Williams, M. D. In 8vo. with plates.

THE BRIDGE WATER TREATISES, COMPLETE IN SEVEN VOLUMES, OCTAVO. Embracing.

I. The Adaptation of External Nature to the Moral and Intellectual Constitution of Man. By the Rev. Thomas Chalmers.

II. The Adaptation of External Nature to the Physical Condition of Man. By John Kidd, M. D., F. R. S.

III. Astronomy and General Physics, Considered with References to Natural Theology. By the Rev. Wm. Whewell.

IV. The Hand: Its Mechanism and Vital Endowments as Evincing Design. By Sir Charles Bell, K. H., F. R. S. With numerous wood cuts.

V. Chemistry, Meteorology, and the Function of Digestion. By Wm. Prout, M. D., F. R. S.

VI. The History, Habits and Instincts of Animals. By the Rev. Wm. Kirby, M. A., F. R. S. Illustrated by numerous Engravings on Copper.

VII. Anatomy and Vegetable Physiology Considered with Reference to Natural Theology. By Peter Mark Roget, M. D. Illustrated with nearly Five Hundred Wood Cuts.

VIII. Geology and Mineralogy, Considered with Reference to Natural Theology. By the Rev. Wm. Buckland, D. D. with numerous engravings on copper, and a large coloured map.

***The work of Buckland, Kirby and Roget may be had separate.

NATIONAL LIBRARY OF MEDICINE



NLM 03274532 7